

Goal: Derive the formula for the sum of a geometric series and explore the intuition behind this formula.

- (1) Consider coloring in a 1×1 square using the following step-by-step process. For the first step, we draw a line vertically down the middle of the square and color the right half:



Since the square has area 1, the area of the shaded region is $1/2$. For the next step, we draw a horizontal line through the remaining space and color in the top quarter:



At this point, the shaded region has area $\frac{1}{2} + \frac{1}{4} = \frac{3}{4}$. We continue this process, at each step coloring half of the remaining space.

- (a) (i). Draw the square after three steps. What is the area of the shaded region? Write this as both an expanded sum, and as a single fraction.
- (ii). Draw the square after four steps. What is the area of the shaded region? Write this as both an expanded sum, and as a single fraction.

(iii). Draw the square after five steps. What is the area of the shaded region? Write this as both an expanded sum, and as a single fraction.

(b) Now, let's think about the area after n steps, where n is an arbitrary number.

(i). Write down a sum that expresses the area after n steps. Write it in both expanded form and in sigma-notation.

(ii). In problem 1, you may have noticed a pattern in your final answers. Use this to guess a simple formula for the area after n steps (in other words, simplify so your answer is no longer a sum).

(iii). Based on this formula, what can you infer about the area of the shaded region as n tends toward infinity? Does this make geometric sense? Why or why not?

- (2) Now let's talk about general geometric sequences and series. A geometric sequence is defined by an *initial term* a and *constant ratio* r , and looks like this:

$$a, ar, ar^2, ar^3, \dots$$

Partial geometric sums are just the partial sums of geometric sequences., i.e., sums of the form

$$S_n = a + ar + \dots + ar^{n-1} = \sum_{i=1}^n ar^{i-1}.$$

Infinite geometric sums, on the other hand, are the limits of the partial sums (whenever these limits exist), and look like this:

$$\lim_{n \rightarrow \infty} S_n = \lim_{n \rightarrow \infty} (a + ar + \dots + ar^{n-1}) = \sum_{i=1}^{\infty} ar^{i-1}.$$

- (a) In problem 1, the areas that you calculated were geometric sums with a specific initial term and constant ratio. What was the initial term a in these sums? The constant ratio r ?

In problem (1), we came up with a simplified expression for the area of the shaded region after n steps. We can generalize this expression to arbitrary geometric sums in the following way. Let a and r be arbitrary numbers with $r \neq 1$.

- (b) Using the original definition of S_n , expand and then simplify $S_n - rS_n$ (Hint: Group like terms. Most of them should cancel.).

- (c) If you did the above calculation carefully, you should get an answer similar to $S_n - rS_n = a - ar^n$. Solve this equation to come up with a simple formula for S_n . Check that this formula agrees with your formula from problem 1.

Partial sum of a geometric series:

$$S_n = a + ar + \cdots + ar^{n-1} = \underline{\hspace{4cm}}$$

Recall that the value of a series is defined to be the limit of its partial sums, so that

$$\lim_{n \rightarrow \infty} (a + ar + \cdots + ar^{n-1}) = \sum_{i=1}^{\infty} ar^{i-1} = \lim_{n \rightarrow \infty} S_n = \lim_{n \rightarrow \infty} \underline{\hspace{2cm}}.$$

- (d) Based on your formula for S_n , what can you say about the convergence or divergence of S_n ?
- (i). Does it depend on a ?

 - (ii). On r ?

 - (iii). For what values of r does the series converge?

 - (iv). For what values does it diverge?

- (v). Using your formula for S_n from Problem 2(c) (on the previous page), take limits to come up with a formula for the value of the sum of a general infinite geometric series. Check that this formula agrees with the area of the square in problem 1.

Sum of an infinite geometric series:

$$\sum_{i=1}^{\infty} ar^{i-1} = \underline{\hspace{2cm}} \quad (\text{provided } |r| < 1)$$

- (3) Now that we know how to calculate finite and infinite geometric sums, let's get some practice with a few examples.

(a) $\sum_{n=1}^{\infty} 2 \cdot \left(\frac{1}{3}\right)^{n-1} =$

(b) $\frac{1}{4} + \frac{1}{8} + \frac{1}{16} + \dots =$

(c) $\sum_{n=1}^{\infty} \frac{e}{\pi^{n-1}} =$

(d) $0.9 + 0.09 + 0.009 + \dots =$

(e) $\sum_{n=1}^{10} 2 \left(\frac{2}{3}\right)^{n-1} =$

(f) $\sum_{n=1}^{\infty} (-2)^{n-1} =$