
Part I - Summary of higher-dimensional versions of the Fundamental Theorem of Calculus Fill in the blanks (assuming appropriate hypotheses are met for the integrands).

1. The theorem regarding the equation $\iint_{S} \mathbf{F} \cdot d\mathbf{S} = \iiint_{E} \operatorname{div} \mathbf{F} \, dV$ can be stated as follows:

Given a surface integral of a vector field \mathbf{F} over a surface S, if the surface S is _____, the surface integral is equal to a _____ over the region bounded by the surface. (Theorem)


- 2. Given a line integral of a vector field **F** over a curve C, if **F** is _____, then the value of the line integral is the difference between f evaluated at the start point and end point of the curve, where $_____f = ____$.
- 3. Given a line integral of a vector field \mathbf{F} along a curve C, if the curve C is _____, the line integral is equal to a ______ of _____ over any orientable surface that has the curve C as its boundary. (______)

4. Given a line integral of a vector field $\mathbf{F} = \langle P, Q \rangle$ over a planar closed curve C (oriented counter-clockwise), the line integral is equal to a ______ of _____ over the planar region bounded by C.



Part II - Practice problems

1. The figure below shows a surface S, which is a sphere of radius 5 centered at the origin, with the top cut off, so the upper edge of the surface lies at z = 4. Use one of the theorems from Chapter 13 to evaluate $\iint_{S} \operatorname{curl} \mathbf{F} \cdot d\mathbf{S}$, where $\mathbf{F} = \langle y, -x, z \rangle$. S is oriented outward.




2. Consider the surface integral $\iint_{S} \mathbf{F} \cdot d\mathbf{S}$, where S is the closed yurt-shaped surface shown below, and $\mathbf{F} = \langle 3x, 2y, z \rangle$. Notice that the surface comprises three separate pieces: the circular base, the cylinder walls,

 $\Gamma = \sqrt{3x}, 2y, z/2$. Notice that the surface comprises three separate pieces, the circular base, the cylinder wais, and the conical top. The cylinder has radius 3 and height 2, and the cone has radius 3 and height 3.

- (a) Discuss with your group the list of steps required to evaluate this surface integral directly.
- (b) Use one of the theorems from Chapter 13 to set up a different type of integral with the same value as the given surface integral.
- (c) Interpret the new integral geometrically to find its value without evaluating it.

S is oriented outward.



3. Consider the two integrals $\int_{C_1} \mathbf{F} \cdot d\mathbf{r}$ and $\int_{C_2} \mathbf{F} \cdot d\mathbf{r}$, where $\mathbf{F} = \langle y, -x, 0 \rangle$, and where C_1 is shown below (solid),

and C_2 is shown below (dashed). A top-view of the vector field **F** is also shown. Do the two line integrals give the same value, or not? Explain.

