

Math 3140 — Fall 2012

Exam #2

Work alone. No materials except pen (or pencil) and paper allowed. Write your solutions on a separate paper. Justify your answers. Giving incorrect or irrelevant justification will be penalized.

Problem 1. Suppose that A , B , and C are groups with their group operations all written multiplicatively. Suppose that $\varphi : A \rightarrow B$ and $\psi : B \rightarrow C$ are homomorphisms of groups. Let $\omega = \psi \circ \varphi$. Show that the function $\omega : A \rightarrow C$ is a homomorphism of groups.

Problem 2. Describe a group with 4 elements that is **not** isomorphic to $\mathbf{Z}/4\mathbf{Z}$. Make sure to explain how you know this group is not isomorphic to $\mathbf{Z}/4\mathbf{Z}$.

Problem 3. Let $\mathbf{R} \times \mathbf{R}$ be the group of pairs (a, b) where a and b are real numbers with group operation

$$(a, b) + (a', b') = (a + a', b + b').$$

Let $\varphi : \mathbf{R} \times \mathbf{R} \rightarrow \mathbf{R}$ be the function

$$\varphi(a, b) = a + b.$$

Show that φ is a homomorphism.

Problem 4. Let $A = \langle 210, 360, 756 \rangle$ be the subgroup of \mathbf{Z} containing $210 = 2 \cdot 3 \cdot 5 \cdot 7$ and $360 = 2^3 \cdot 3^2 \cdot 5$ and $756 = 2^2 \cdot 3^3 \cdot 7$. Recall that this means A consists of all integers of the form $210x + 360y + 756z$ with $x, y, z \in \mathbf{Z}$. List all numbers between 0 and 100 (inclusive) that are not in $\langle 210, 360 \rangle$, nor in $\langle 210, 756 \rangle$, nor in $\langle 360, 756 \rangle$. Remember to justify your answer.

Problem 5. Prove that every odd permutation in S_n has even order. (Recall that an odd permutation is one whose sign is -1 .)

Problem 6. Let $X = \{1, 2, 3, 4, 5\}$ and let Y be the set of subsets of X .

(a) How many elements does Y have?

For each $\sigma \in S_5$, let $T_\sigma : Y \rightarrow Y$ be the function defined by $T_\sigma(A) = \{\sigma(x) \mid x \in A\}$.

(b) Compute $T_{(15)(234)}(\{1, 3, 4\})$.

(c) Compute $T_{(23)}(\{1, 5\})$.

(d) List the elements of the stabilizer of $\{1, 2, 3\}$. Recall that the stabilizer of an element $y \in Y$ is the set of all $\sigma \in S_5$ such that $T_\sigma(y) = y$.

(e) How many elements will there be in the orbit of $\{1, 2, 3\}$ under the action of S_5 on Y ?

(f) (Extra credit) Prove that the stabilizer subgroup of $\{1, 2, 3\}$ is isomorphic to $S_3 \times S_2$. You may use the fact that $\varphi(\sigma) = T_\sigma$ is a homomorphism from S_5 into S_Y without proving it.

Definition 1. A **group** is a set G with an operation $*$: $G \times G \rightarrow G$ such that (i) $a * (b * c) = (a * b) * c$ for all $a, b, c \in G$, (ii) there is an $e \in G$ such that $e * a = a = a * e$ for all $a \in G$, and (iii) for any $a \in G$ there is an $a^{-1} \in G$ such that $aa^{-1} = e = a^{-1}a$. The group G is said to be **abelian** if $a * b = b * a$ for all $a, b \in G$.

A subset $H \subset G$ is called a **subgroup** if (i) for all $a, b \in H$ the element $a * b$ is in H , and (ii) H is a group with operation $*$.

A group is called **cyclic** if it is isomorphic \mathbf{Z} or it is isomorphic to $\mathbf{Z}/n\mathbf{Z}$ for some integer n .

The **order** of an element g of a group G (written multiplicatively with identity element 1) is the smallest positive integer n such that $g^n = 1$.

Definition 2. Suppose that G and H are groups with operations written multiplicatively. A **homomorphism** $\varphi : G \rightarrow H$ is a function $\varphi : G \rightarrow H$ such that $\varphi(xy) = \varphi(x)\varphi(y)$. A homomorphism is called an **isomorphism** if it is also a bijection.

The **kernel** of φ is the set $\ker(\varphi) = \{x \in G \mid \varphi(x) = 1\}$ where 1 is the identity in H .

The **image** of φ is the set $\text{im}(\varphi) = \{y \in H \mid \exists x \in G, y = \varphi(x)\}$.

Definition 3. An **action** of a group G on a set X is a homomorphism $\varphi : G \rightarrow S_X$. Use the notation $T_g = \varphi(g)$. The **stabilizer** of an element $x \in X$ is $\{g \in G \mid T_g(x) = x\}$. The **orbit** of $x \in X$ is $\{y \in X \mid \exists g \in G, T_g(x) = y\}$.

Notation

\mathbf{Z} is the set of integers and \mathbf{R} is the set of real numbers.

D_n is the set of rigid symmetries of a regular n -gon.

$\mathbf{Z}/n\mathbf{Z}$ is the set of equivalence classes of integers modulo n .

$\text{gcd}\{a_1, \dots, a_n\}$ denotes the greatest common divisor of integers a_1, \dots, a_n .

If X is a set, S_X is the set of bijections from X to itself. If $X = \{1, 2, \dots, n\}$ then S_X is also written S_n .

If $\sigma \in S_n$ the sign of σ is the expression $\text{sgn}(\sigma) = \prod_{1 \leq i < j \leq n} \frac{x_{\sigma(i)} - x_{\sigma(j)}}{x_i - x_j}$. An element of S_n is called a **transposition**

if it exchanges two numbers and leaves all others unchanged. An element of S_n is called **even** if its sign is 1 and **odd** if its sign is -1 . The set of even elements of S_n is denoted A_n .

Theorems

Proposition 1. The following are abelian groups: (i) \mathbf{Z} under addition, (ii) $\mathbf{Z}/n\mathbf{Z}$ under addition, (iii) \mathbf{R} under addition, (iv) \mathbf{R}^* under multiplication, (v) \mathbf{C}^* under multiplication, (vi) S_X if X is a set with 2 or fewer elements.

The following are non-abelian groups: (vii) D_n , (viii) S_X if X is a set with 3 or more elements.

Theorem 2 (Cayley's theorem). Every group is isomorphic to a subgroup of the group of symmetries of some set.

Proposition 3. Let G be a group. A subset $H \subset G$ is a subgroup if and only if both (i) $H \neq \emptyset$, and (ii) for all $a, b \in H$ the element ab^{-1} is in H .

Theorem 4. If x and y are integers with greatest common divisor d there are integers a and b such that $ax + by = d$.

Theorem 5. If G is a cyclic group then every subgroup of G is cyclic.

Proposition 6. Suppose that G and H are groups with operations written multiplicatively and identity elements both called 1. If $\varphi : G \rightarrow H$ is a homomorphism of groups then (i) $\varphi(1) = 1$, (ii) $\varphi(x^{-1}) = \varphi(x)^{-1}$ for all $x \in G$, (iii) $\ker(\varphi)$ is a subgroup of G , (iv) $\text{im}(\varphi)$ is a subgroup of H .

Proposition 7. If $\sigma \in S_n$ then $\text{sgn}(\sigma) \in \{\pm 1\}$ and the function $\text{sgn} : S_n \rightarrow \{\pm 1\}$ is a homomorphism. If τ is a transposition then $\text{sgn}(\tau) = -1$.

Proposition 8. For complex numbers z and w , we have $|zw| = |z||w|$.

Proposition 9. If $\varphi : G \rightarrow H$ is an isomorphism of groups then $\varphi^{-1} : H \rightarrow G$ is also an isomorphism.

Proposition 10. The inverse of a 2×2 matrix $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ is given by $\frac{1}{ad - bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$ provided $\frac{1}{ad - bc}$ exists.

Proposition 11. Let g be an element of a group G and suppose $g^n = 1$. Then $\text{ord}(g)$ is finite and divides n .

Theorem 12. Let $\varphi : G \rightarrow S_X$ be a group action. For $x \in X$, let G_x be the stabilizer of x and let O_x be the orbit of x . If G is finite then $\#G_x \cdot \#O_x = \#G$.