



Objectives:

- Find derivatives of logarithmic functions.

Review of logarithmic functions:



Inverse: $y = \ln(x)$
 Domain: $(0, \infty)$
 Range: $(-\infty, \infty)$



Important facts:

$y = e^x$ means $\ln y = x$ and $y = a^x$ means $\log_a y = x$

$\log_2 8 = 3$	is equivalent to	$2^3 = 8$
$\log_{10} 100 = 2$	is equivalent to	$10^2 = 100$
$\log_{10} 0.001 = -3$	is equivalent to	$10^{-3} = \frac{1}{1000} = 0.001$
$\ln \sqrt{e} = \frac{1}{2}$	is equivalent to	$e^{1/2} = \sqrt{e}$
$y = \log_a x$	is equivalent to	$a^y = x$
$\ln y = x$	is equivalent to	$y = e^x$

Solving Equations:

Solve $5e^{0.34t} = 6$ for t .

$$e^{0.34t} = \frac{6}{5}$$

$$\ln(e^{0.34t}) = \ln\left(\frac{6}{5}\right)$$

$$0.34t = \ln \frac{6}{5}$$

$$t = \frac{\ln \frac{6}{5}}{0.34}$$

Solve $\log_2(x) + \log_2(x - 1) = 1$.
 (Hint: use laws of logs)

$$\log_2(x(x - 1)) = 1$$

$$x(x - 1) = 2^1$$

$$x^2 - x = 2$$

$$x^2 - x - 2 = 0$$



$$(x - 2)(x + 1) = 0$$

So $x = 2$ or $x = -1$ but substituting shows $x = -1$ is not in the domain. So $x = 2$ is the solution.

Derivatives of logarithmic functions:

$$1. \frac{d}{dx} (\ln x) = \frac{1}{x}$$

$$2. \frac{d}{dx} (\log_a x) = \frac{1}{\ln a} \cdot \frac{1}{x}$$



Proof: (of 1.)

Step 1: $y = \ln x$

Step 2: $e^y = x$ (use inverses)

Step 3: $e^y \cdot y' = 1$ (differentiate)

Step 4: $y' = \frac{1}{e^y}$ (solve for y')

Step 5: $y' = \frac{1}{x}$ (substitute to write in terms of x)

Try the proof for 2 at home!

Examples: Find the derivatives of the following functions using our new formulas.

1. $f(x) = x \ln(x)$

$$f'(x) = x \cdot \frac{1}{x} + \ln x = 1 + \ln x$$

2. $f(x) = \ln(\arctan(x))$

$$f'(x) = \frac{1}{\arctan x} \cdot \frac{1}{1+x^2}$$

3. $f(x) = \sqrt{\log_2(x)} = (\log_2 x)^{1/2}$

$$f'(x) = \frac{1}{2}(\log_2 x)^{-1/2} \cdot \frac{1}{x \ln 2}$$

4. $f(x) = \arcsin(e^{\tan(x^2)})$

$$f'(x) = \frac{1}{\sqrt{1 - e^{2 \tan(x^2)}}} \cdot e^{\tan(x^2)} \cdot \sec^2(x^2) \cdot 2x$$

5. $f(x) = \ln\left(\frac{x^2 \sqrt{x-1}}{(x+3)^4}\right) = \ln(x^2) + \ln(\sqrt{x-1}) - \ln((x+3)^4) = 2 \ln x + \frac{1}{2} \ln(x-1) - 4 \ln(x+3)$

$$f'(x) = \frac{2}{x} + \frac{1}{2(x-1)} - \frac{4}{x+3}$$

Logarithmic Differentiation: Why would we take logarithms to take derivatives?

- Use it with functions that have products/quotients/powers, like $f(x) = \frac{x^4\sqrt{x^2+1}}{(3x+4)^2}$
- Use if there is a variable in the base and the exponent, like $f(x) = x^{\sin x}$, since neither the power rule nor the exponential rule apply.

Examples Find $f'(x)$ for the following functions.

1. $f(x) = \frac{x^4\sqrt{x^2+1}}{(3x+4)^2}$.

Step 1: Use y for $f(x)$

$$y = \frac{x^4\sqrt{x^2+1}}{(3x+4)^2}$$

Step 2: Take \ln of both sides

$$\ln y = \ln \left(\frac{x^4\sqrt{x^2+1}}{(3x+4)^2} \right)$$

Step 3: Use laws of logs to simplify

$$\ln y = 4 \ln x + \frac{1}{2} \ln(x^2+1) - 2 \ln(3x+4)$$

Step 4: Differentiate implicitly

$$\frac{1}{y} \cdot y' = \frac{4}{x} + \frac{1}{2} \cdot \frac{1}{x^2+1} \cdot 2x - \frac{2}{3x+4} \cdot 3$$

Step 5: Solve for y'

$$y' = y \left(\frac{4}{x} + \frac{x}{x^2+1} - \frac{6}{3x+4} \right) = \left(\frac{x^4\sqrt{x^2+1}}{(3x+4)^2} \right) \left(\frac{4}{x} + \frac{x}{x^2+1} - \frac{6}{3x+4} \right)$$

2. $f(x) = x^{\sin x}$

$$y = x^{\sin x}$$

$$\ln y = \ln(x^{\sin(x)}) = \sin x \cdot \ln x$$

$$\frac{1}{y} y' = \frac{\sin(x)}{x} + \ln x \cos x$$

$$\begin{aligned} y' &= y \left(\frac{\sin(x)}{x} + \ln x \cos x \right) \\ &= x^{\sin x} \left(\frac{\sin(x)}{x} + \ln x \cos x \right) \end{aligned}$$