

1. Consider a 3-dimensional solid S where the base of S is the region enclosed by the parabola $y = 1 - x^2$ and the x -axis and cross-sections perpendicular to the x -axis are squares.
 - (a) Sketch the region of the base of S and include a cross-section of S .
 - (b) Find the area $A(x)$ of the cross section of S at x . (This is what we call an arbitrary cross-section.)
 - (c) By integrating $A(x)$ over an appropriate interval, compute the volume of the solid S .

2. Consider a 3-dimensional solid S whose base is the triangular region with vertices $(0, 0)$, $(1, 0)$, and $(0, 2)$ and the cross-sections perpendicular to the y -axis are equilateral triangles.
 - (a) Sketch the region of the base of S and include a cross-section of S .
 - (b) Find a formula for the area of an equilateral triangle with side-length a .
 - (c) Using the formula from part (b), find the area $A(y)$ of the cross section of S at y .
 - (d) By integrating $A(y)$ over an appropriate interval, compute the volume of the solid S .

3. Consider a 3-dimensional solid S whose base is the triangular region with vertices $(0, 0)$, $(2, 0)$, and $(0, 1)$ and the cross-section perpendicular to the x -axis are squares. Find the volume of S .

4. Find the volume of the solid S which is a right circular cone with height h and base radius r .

5. Find the volume of the solid S whose base is a circular disk with radius r and the parallel cross-sections perpendicular to the base are squares.