Math 3130 - Assignment 6

Due February 26, 2016 Markus Steindl

- (46) Let $\mathbf{v}_1, \dots, \mathbf{v}_n$ be vectors in a vector space V. Show that $H := \operatorname{Span}\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ is a subspace of V.
- (47) Let A be an $m \times n$ matrix. Prove that the Nullspace of A is a subspace of \mathbb{R}^n .
- (48) Let $M_{2\times 2}$ be the set of all 2×2 matrices. Let + be the sum of matrices and \cdot be the multiplication of a matrix by a scalar.
 - (a) Show that $M_{2\times 2}$ forms is a vector space.
 - (b) Let H be the set of invertible 2×2 matrices. Show that H is not a subspace of $M_{2\times 2}$.
- (49) Show that $V := \{ \begin{bmatrix} a \\ b \end{bmatrix} \mid a, b \in \mathbb{R}, \ a \ge 0 \}$ is no subspace of \mathbb{R}^2 .
- (50) (a) Which of the vectors $\mathbf{u}, \mathbf{v}, \mathbf{w}, \mathbf{x}$ are in Nul A?
 - (b) Which of the vectors $\mathbf{u}, \mathbf{v}, \mathbf{w}, \mathbf{x}$ are in Col A?

$$\mathbf{u} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \quad \mathbf{v} = \begin{bmatrix} -1 \\ 0 \\ 2 \\ -1 \end{bmatrix}, \quad \mathbf{w} = \begin{bmatrix} 2 \\ -1 \\ 5 \end{bmatrix}, \quad \mathbf{x} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}, \quad A = \begin{bmatrix} 0 & 0 & 2 & 4 \\ 2 & -4 & 1 & 0 \\ -3 & 6 & 2 & 7 \end{bmatrix}$$

- (51) Let A be the matrix from (50).
 - (a) Solve $A\mathbf{x} = \mathbf{0}$ and give the solution in parametric vector form.
 - (b) Find vectors $\mathbf{u}, \mathbf{v} \in \mathbb{R}^4$ such that

$$\operatorname{Nul} A = \{ r\mathbf{u} + s\mathbf{v} \mid r, s \in \mathbb{R} \}.$$

- (52) Let $V := \{f \colon \mathbb{R} \to \mathbb{R}\}$ be the vector space of functions on \mathbb{R} .
 - (a) Is $\{f : \mathbb{R} \to \mathbb{R} \mid f(0) = 1\}$ a subspace of V?
 - (b) Is $\{f \colon \mathbb{R} \to \mathbb{R} \mid f(1) = 0\}$ a subspace of V?
 - (c) Is $\{f : \mathbb{R} \to \mathbb{R} \mid f \text{ is continuous}\}\$ a subspace of V?
- (53) Is $\{0\}$ a subspace of \mathbb{R}^n ?
- (54) Are the vectors $\mathbf{u} = 1$, $\mathbf{v} = t$, $\mathbf{w} = t^2$ in the vector space $V := \{f : \mathbb{R} \to \mathbb{R}\}$ linearly independent?