

# Math 2001 - Assignment 4

Due February 14, 2018

- (1) (a) How many different truth tables (Boolean functions) are there for 2 statements  $x_1, x_2$ ? How many for  $k$  statements  $x_1, \dots, x_k$ ?  
(b) Let  $f(x_1, x_2, x_3)$  be a Boolean function that is true for the following assignments and false otherwise.

$x_1$	$x_2$	$x_3$	$f(x_1, x_2, x_3)$
$T$	$T$	$F$	$T$
$T$	$F$	$T$	$T$
$F$	$T$	$T$	$T$

Write an expression for  $f(x_1, x_2, x_3)$  using only  $\wedge, \vee, \sim$ .

## Solution

- (a) A truth table for 2 statements  $x_1, x_2$  has  $2^2 = 4$  rows. For each row we can choose either true or false. Hence we have  $2^4$  options to make a truth table.  
A truth table for  $k$  statements  $x_1, \dots, x_k$  has  $2^k$  rows. Hence there are  $2^{2^k}$  such tables.  
(b)  $f(x_1, x_2, x_3)$  and  $(x_1 \wedge x_2 \wedge \sim x_3) \vee (x_1 \wedge \sim x_2 \wedge x_3) \vee (\sim x_1 \wedge x_2 \wedge x_3)$  are true at exactly the same assignments. Hence they are equal.  
(2) [1, Section 2.7]: Exercises 4,6,7,9,10. Also give the negation of the corresponding statements.

## Solution:

4. For all elements  $X$  in the powerset of  $\mathbb{N}$ , we have that  $X$  is a subset of  $\mathbb{R}$ .  
Every subset of  $\mathbb{N}$  is a subset of  $\mathbb{R}$ .

**True since**  $\mathbb{N} \subseteq \mathbb{R}$

**Negation:**  $\exists X \in P(\mathbb{N}), X \not\subseteq \mathbb{R}$

6. There exists a natural number  $n$  such that every subset of  $\mathbb{N}$  has less than  $n$  elements.

**False,  $\{1, \dots, n\}$  is a subset with  $n$  elements**

**Negation:**  $\forall n \in \mathbb{N} \exists X \in P(\mathbb{N}), |X| \geq n$

7. For every subset  $X$  of  $\mathbb{N}$  there exists an integer  $n$  such that  $X$  has size  $n$ .

**False,  $X = \mathbb{N}$  is a subset of  $\mathbb{N}$  of infinite size.**

**Negation:**  $\exists X \subseteq \mathbb{N} \forall n \in \mathbb{Z} : |X| \neq n$

9. For every integer  $n$  there exists an integer  $m$  such that  $m = n + 5$ .

**True,  $m = n + 5$  is the integer we are looking for.**

**Negation:**  $\exists n \in \mathbb{Z} \forall m \in \mathbb{Z} : m \neq n + 5$

10. There exists an integer  $m$  for every  $n$  such that  $m = n + 5$ .

**False, the same number  $m$  cannot work for  $n = 0$  and  $n = 1$ .**

**Negation:**  $\forall m \in \mathbb{Z} \forall \exists n \in \mathbb{Z} : m \neq n + 5$

- (3) Formulate the following sentences using quantifiers and logical operations. Are they true? Negate them.

- (a) For all integers  $n$  we have that  $n(n+1)$  is even.

**Solution**  $\forall n \in \mathbb{Z} n(n+1)$  is even.

True because one of  $n$  or  $n+1$  is even.

Negation:  $\exists n \in \mathbb{Z} n(n+1)$  is odd.

- (b) There exists a real number  $z$  such that  $x+z=x$  for every real  $x$ .

**Solution**  $\exists z \in \mathbb{R} \forall x \in \mathbb{R} x+z=x$

True for  $z=0$ .

Negation:  $\forall z \in \mathbb{R} \exists x \in \mathbb{R} x+z \neq x$

- (c) Every real number is smaller than some integer.

**Solution**  $\forall x \in \mathbb{R} \exists z \in \mathbb{Z} x < z$

True.

Negation:  $\exists x \in \mathbb{R} \forall z \in \mathbb{Z} x \geq z$

- (4) Negate the following sentences. Are they true?

- (a) If  $x^2$  is rational, then so is  $x$ .

**Solution**  $x^2 \in \mathbb{Q} \Rightarrow x \in \mathbb{Q}$ .

False, e.g.  $2 \in \mathbb{Q}$  and  $\sqrt{2} \notin \mathbb{Q}$

Negation:  $x^2 \in \mathbb{Q}$  and  $x \notin \mathbb{Q}$ .

- (b)  $xy=0$  iff  $x=0$  or  $y=0$

**True, Negation**  $xy \neq 0$  iff  $x=0$  or  $y=0$

$xy=0$  iff  $x \neq 0$  and  $y \neq 0$

- (c) The derivative of a polynomial function  $f$  is 0 iff  $f$  is constant.

**True, Negation:** The derivative of a polynomial function  $f$  is 0 iff  $f$  is not constant.

- (d)  $\exists x \in \mathbb{R} : x^2 = -1$

**False, Negation:**  $\forall x \in \mathbb{R} : x^2 \neq -1$

- (e)  $\forall r \in \mathbb{R} : \sin(r\pi) = 0 \Leftrightarrow r$  is an integer

**Negation:**  $\exists r \in \mathbb{R} : \sin(r\pi) = 0$  iff  $r$  is not an integer

- (5) True or false? Give a proof or a counter-example:

- (a)  $\forall x \in \mathbb{R} \forall y \in \mathbb{R} x+y=1$

**False, counter-example:**  $x=y=0$

- (b)  $\forall x \in \mathbb{R} \exists y \in \mathbb{R} x+y=1$

**True:** for  $x \in \mathbb{R}$  we have  $y=1-x$  such that  $x+y=1$

- (c)  $\exists x \in \mathbb{R} \forall y \in \mathbb{R} x+y=1$

**False:** Suppose we have such a fixed  $x \in \mathbb{R}$ . Then for  $y=-x$  we'd have  $x+y=0 \neq 1$ . Hence there cannot be a single  $x$  that makes  $x+y=1$  true for all  $y \in \mathbb{R}$ .

- (d)  $\exists x \in \mathbb{R} \exists y \in \mathbb{R} x+y=1$

**True:** e.g.  $x=0, y=1$

- (6) Write as complete English sentences. True or false? Negate:

- (a)  $\forall a \in \mathbb{R} \exists b \in \mathbb{R} \forall c \in \mathbb{R} : a < b \Rightarrow c < b$

**Solution:** For all  $a \in \mathbb{R}$  there exists  $b \in \mathbb{R}$  such that for all  $c \in \mathbb{R}$ , if  $a < b$  then  $c < b$ .

True: Let  $a \in \mathbb{R}$  arbitrary. Next choose  $b \in \mathbb{R}$  such that  $a \not< b$ , say  $b=a$ . Then  $\forall c \in \mathbb{R} : a < b \Rightarrow c < b$  is true. (Note that by the choice of  $b$ , the assumption  $a < b$  of the implication is false. Hence FALSE  $\Rightarrow c < b$  is true.)

Negation:  $\exists a \in \mathbb{R} \forall b \in \mathbb{R} \exists c \in \mathbb{R} : a < b \wedge b \leq c$ .

Note that the negation is clearly false. Hence again the original statement must be true!

(b)  $\forall \text{ set } A \forall \text{ set } B \exists \text{ set } C : A \cup B = C.$

**Solution:** For all sets  $A$  and  $B$  there exists a set  $C$  that is the union of  $A$  and  $B$ .

True by Axiom of Unions in Zermelo-Fraenkel Set Theory.

Negation:  $\exists \text{ set } A \exists \text{ set } B \forall \text{ set } C : A \cup B \neq C.$

There exist sets  $A$  and  $B$  whose union is not a set.

(c)  $\forall x, y \in \mathbb{R} \forall \epsilon > 0 \exists \delta > 0 : |x - y| < \epsilon \Rightarrow |2x - 2y| < \delta$

**Solution:** For all reals  $x, y$  and every  $\epsilon > 0$  there exists a  $\delta > 0$  such that  $|x - y| < \epsilon$  implies  $|2x - 2y| < \delta$ .

True for  $\delta = 2\epsilon$ .

Negation:  $\exists x, y \in \mathbb{R} \exists \epsilon > 0 \forall \delta > 0 : |x - y| < \epsilon \wedge |2x - 2y| \geq \delta.$

#### REFERENCES

- [1] Richard Hammack. The Book of Proof. Creative Commons, 2nd edition, 2013. Available for free: <http://www.people.vcu.edu/~rhammack/BookOfProof/>