

# Math 2130 - Assignment 4

Due September 24, 2021

(1) Let  $T: \mathbb{R}^2 \rightarrow \mathbb{R}^3$  be a linear map such that

$$T\left(\begin{bmatrix} 1 \\ 2 \end{bmatrix}\right) = \begin{bmatrix} 2 \\ 0 \\ -3 \end{bmatrix}, \quad T\left(\begin{bmatrix} 3 \\ 2 \end{bmatrix}\right) = \begin{bmatrix} -2 \\ 2 \\ 1 \end{bmatrix}.$$

(a) Use linearity to find  $T(e_1)$  and  $T(e_2)$  for the unit vectors  $e_1, e_2$  in  $\mathbb{R}^2$ .

(b) Give the standard matrix for  $T$  and determine  $T\left(\begin{bmatrix} x \\ y \end{bmatrix}\right)$  for arbitrary  $x, y \in \mathbb{R}$ .

**Solution:**

(a) First write the unit vectors as linear combinations of  $\begin{bmatrix} 1 \\ 2 \end{bmatrix}$  and  $\begin{bmatrix} 3 \\ 2 \end{bmatrix}$ .

Solve

$$x \begin{bmatrix} 1 \\ 2 \end{bmatrix} + y \begin{bmatrix} 3 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

to get  $x = -\frac{1}{2}$  and  $y = \frac{1}{2}$ . By the linearity of  $T$  we obtain

$$\begin{aligned} T\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) &= T\left(-\frac{1}{2} \begin{bmatrix} 1 \\ 2 \end{bmatrix} + \frac{1}{2} \begin{bmatrix} 3 \\ 2 \end{bmatrix}\right) \\ &= -\frac{1}{2} T\left(\begin{bmatrix} 1 \\ 2 \end{bmatrix}\right) + \frac{1}{2} T\left(\begin{bmatrix} 3 \\ 2 \end{bmatrix}\right) \\ &= -\frac{1}{2} \begin{bmatrix} 2 \\ 0 \\ -3 \end{bmatrix} + \frac{1}{2} \begin{bmatrix} -2 \\ 2 \\ 1 \end{bmatrix} \\ &= \begin{bmatrix} -2 \\ 1 \\ 2 \end{bmatrix} \end{aligned}$$

Similarly we compute that

$$\begin{bmatrix} 0 \\ 1 \end{bmatrix} = \frac{3}{4} \begin{bmatrix} 1 \\ 2 \end{bmatrix} - \frac{1}{4} \begin{bmatrix} 3 \\ 2 \end{bmatrix}$$

and hence obtain

$$T\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = \frac{3}{4} \begin{bmatrix} 2 \\ 0 \\ -3 \end{bmatrix} - \frac{1}{4} \begin{bmatrix} -2 \\ 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ -1/2 \\ -5/2 \end{bmatrix}$$

(b) By (a) we know the standard matrix of  $T$  is

$$A = \begin{bmatrix} -2 & 2 \\ 1 & -1/2 \\ 2 & -5/2 \end{bmatrix}.$$

$$\text{Thus } T\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = A \cdot \begin{bmatrix} x \\ y \end{bmatrix}.$$

□

- (2) Is the following injective, surjective, bijective? What is its range?

$$T : \mathbb{R}^3 \rightarrow \mathbb{R}^2, x \mapsto \begin{bmatrix} 0 & 2 & -1 \\ 0 & 0 & 3 \end{bmatrix} \cdot x$$

**Solution:** Not injective because  $x_1$  is free in  $A \cdot x = \mathbf{0}$ . Alternatively, the columns of  $A$  are linearly dependent. So  $T$  is not injective (Theorem 12 of Section 1.9).

Surjective because  $A$  is in row echelon form and has no 0-rows (Theorem 12 of Section 1.9). Hence its range is just its codomain  $\mathbb{R}^2$ .

Bijective means injective and surjective. Hence  $T$  is not bijective because it is not injective.  $\square$

- (3) Is the following injective, surjective, bijective?

$$T : \mathbb{R}^3 \rightarrow \mathbb{R}^3, x \mapsto \begin{bmatrix} 1 & -1 & 2 \\ -2 & 0 & 1 \\ 3 & -1 & 1 \end{bmatrix} \cdot x$$

**Solution:** Row reduce the standard matrix of  $T$  to get

$$\begin{array}{ccc|ccc} 1 & -1 & 2 & & & \\ -2 & 0 & 1 & & & \\ 3 & -1 & 1 & & & \\ \hline 1 & -1 & 2 & & & \\ 0 & -2 & 5 & 2I + II & & \\ 0 & 2 & -5 & -3I + III & & \\ \hline 1 & -1 & 2 & & & \\ 0 & -2 & 5 & & & \\ 0 & 0 & 0 & -II + III & & \end{array}$$

$T$  is not injective because not every column of the echelon form of  $A$  has a pivot. In particular  $x_3$  is free in  $A \cdot x = \mathbf{0}$ .

$T$  is not surjective because the echelon form of  $A$  has a zero row. Hence  $Ax = y$  is not consistent for every  $y \in \mathbb{R}^3$ .

Since  $T$  is neither injective nor surjective, it is certainly not bijective.  $\square$

- (4) True or False? Explain why and correct the false statements to make them true.
- A linear transformation  $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$  is completely determined by the images of the unit vectors in  $\mathbb{R}^n$ .
  - Not every linear transformation  $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$  can be written as  $T(x) = Ax$  for some matrix  $A$ .
  - The composition of any two linear transformations is linear as well.

**Solution:**

- True because every vector is a linear combination of unit vectors.
- False. Every linear  $T: \mathbb{R}^n \rightarrow \mathbb{R}^m$  can be written as  $T(x) = Ax$  for its standard matrix  $A$ .
- True. If  $S, T$  are linear, then

$$S(T(u + v)) = S(T(u) + T(v)) = S(T(u)) + S(T(v))$$

for all  $u, v$  in the domain of  $T$  and

$$S(T(cu)) = S(cT(u)) = cS(T(u))$$

for  $c \in \mathbb{R}$ . Hence  $S$  composed with  $T$  is linear.  $\square$

- (5) True or False? Explain why and correct the false statements to make them true.

- (a)  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is onto  $\mathbb{R}^m$  if every vector  $x \in \mathbb{R}^n$  is mapped onto some vector in  $\mathbb{R}^m$ .
- (b)  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is one-to-one if every vector  $x \in \mathbb{R}^n$  is mapped onto a unique vector in  $\mathbb{R}^m$ .
- (c) A linear map  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$  cannot be one-to-one.
- (d) There is a surjective linear transformation  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^4$ .

**Solution:**

- (a) False. Any function  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  maps every vector  $x \in \mathbb{R}^n$  onto some vector  $T(x)$  in  $\mathbb{R}^m$ .  
The correct statement is:  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is onto  $\mathbb{R}^m$  if for every vector  $y \in \mathbb{R}^m$  there is some vector  $x \in \mathbb{R}^n$  such that  $T(x) = y$ .
- (b) False. Any function  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  maps every vector  $x \in \mathbb{R}^n$  onto the unique vector  $T(x)$  in  $\mathbb{R}^m$ .  
The correct statement is:  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is one-to-one if any 2 distinct vectors  $x_1, x_2 \in \mathbb{R}^n$  are mapped to distinct vectors  $T(x_1), T(x_2)$ .
- (c) True. If  $A$  is the  $2 \times 3$  standard matrix of  $T$ , then solving  $A \cdot x = \mathbf{0}$  will always yield at least one free variable.
- (d) False. If  $A$  is the  $3 \times 4$  standard matrix of  $T$ , then  $Ax = y$  cannot have a solution for every  $y \in \mathbb{R}^4$  since the echelon form of  $A$  has at least one zero row.

□

- (6) If defined, compute the following for the matrices

$$A = \begin{bmatrix} 2 & 1 & -4 \\ 3 & -1 & 1 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & -1 \\ -3 & 4 \\ 2 & 0 \end{bmatrix}, \quad C = \begin{bmatrix} -2 & 1 \\ 2 & -3 \end{bmatrix}$$

Else explain why the computation is not defined.

- (a)  $AB$                       (b)  $BA$                       (c)  $AC$                       (d)  $A + C$                       (e)  $AB + 2C$

**Solution:**

$$AB = \begin{bmatrix} 2 \cdot 1 + 1 \cdot (-3) - 4 \cdot 2 & 2 \cdot (-1) + 1 \cdot 4 - 4 \cdot 0 \\ 3 \cdot 1 - 1 \cdot (-3) + 1 \cdot 2 & 3 \cdot (-1) - 1 \cdot 4 + 1 \cdot 0 \end{bmatrix} = \begin{bmatrix} -9 & 2 \\ 8 & -7 \end{bmatrix}$$

$$BA = \begin{bmatrix} 1 \cdot 2 - 1 \cdot 3 & 1 \cdot 1 - 1 \cdot (-1) & 1 \cdot (-4) - 1 \cdot 1 \\ * & * & * \\ * & * & * \end{bmatrix} = \begin{bmatrix} -1 & 2 & -5 \\ 6 & -7 & 16 \\ 4 & 2 & -8 \end{bmatrix}$$

$AC$  is undefined since the length of  $A$ 's rows and the length of  $C$ 's columns are not the same.

$A + C$  is undefined since the sizes of  $A$  and  $C$  don't match.

$$AB + 2C = \begin{bmatrix} -9 & 2 \\ 8 & -7 \end{bmatrix} + \begin{bmatrix} -4 & 2 \\ 4 & -6 \end{bmatrix} = \begin{bmatrix} -13 & 4 \\ 12 & -13 \end{bmatrix}$$

□

- (7) Let  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  first rotate points around the origin by  $60^\circ$  counter clockwise and then reflect points at the line with equation  $y = x$ . Give the standard matrix for  $T$ .
- (a) Recall the standard matrix  $A$  for the rotation  $R$  by  $60^\circ$  from class.
- (b) Determine the standard matrix  $B$  for the reflection  $S$  at the line with equation  $y = x$  (a sketch will help).
- (c) Since  $T$  is the composition of  $S$  and  $R$ , compute the standard matrix  $C$  of  $T$  as the product of  $B$  and  $A$ . Careful about the order!

**Solution:**

(a) The standard matrix for the rotation  $R$  by  $\alpha = 60^\circ$  is

$$A = \begin{bmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{bmatrix}$$

(b) The reflection  $S$  at the diagonal flips the unit vectors, i.e.,  $T(e_1) = e_2$  and  $T(e_2) = e_1$ . Hence the standard matrix of  $S$  is

$$B = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$$

(c) Since  $T = S \circ R$  ( $S$  after  $R$ ), its standard matrix is

$$C = BA = \begin{bmatrix} \sin \alpha & \cos \alpha \\ \cos \alpha & -\sin \alpha \end{bmatrix}$$

□

(8) Continuation of (7): What is the standard matrix for  $U: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  which first reflects points at the line with equation  $y = x$  and then rotates points around the origin by  $60^\circ$  counter clockwise? Compare  $T$  and  $U$ .

**Solution:** Since  $U = R \circ S$  ( $R$  after  $S$ ), its standard matrix is

$$AB = \begin{bmatrix} -\sin \alpha & \cos \alpha \\ \cos \alpha & \sin \alpha \end{bmatrix}$$

Comparing the result with the standard matrix of  $S \circ R$ , we see that the result is not the same. Order of function composition and matrix multiplication matters!

□