

Math 2130 - Assignment 12

Due Nov 19, 2021

- (1) Are the matrices A, B, C, D in (3), (4), (5) of assignment 11 diagonalizable? How?

Solution:

A is not diagonalizable because its eigenvalue -3 has multiplicity 2 but the corresponding eigenspace only dimension 1.

B is diagonalizable because it has 3 distinct eigenvalues, so

$$B = P \begin{bmatrix} 2 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 3 \end{bmatrix} P^{-1} \text{ for } P = \begin{bmatrix} 2 & 0 & 0 \\ 1 & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix}$$

C is diagonalizable because it has 2 distinct eigenvalues, so

$$C = P \begin{bmatrix} 1 + \sqrt{6} & 0 \\ 0 & 1 - \sqrt{6} \end{bmatrix} P^{-1} \text{ for } P = \begin{bmatrix} 2 & -2 \\ \sqrt{6} & \sqrt{6} \end{bmatrix}$$

D is not diagonalizable because its eigenvalue -3 has multiplicity 2 but the corresponding eigenspace only dimension 1. \square

- (2) Let A be an $n \times n$ -matrix. Are the following true or false? Explain why:
- If A has n eigenvectors, then A is diagonalizable.
 - If a 4×4 -matrix A has two eigenvalues with eigenspaces of dimension 3 and 1, respectively, then A is diagonalizable.
 - A is diagonalizable iff A has n eigenvalues (counting multiplicities).
 - If \mathbb{R}^n has a basis of eigenvectors of A , then A is diagonalizable.
 - Every triangular matrix is diagonalizable.

Solution:

- False.** You need n linearly independent eigenvectors.
- True.**
- False.** See for example A in problem 92.
- True.** A basis of \mathbb{R}^n of eigenvectors consists of n linearly independent eigenvectors.
- False.** See example A in the previous problem. \square

- (3) Let A be the standard matrix for the reflection t of \mathbb{R}^2 on some line g through the origin. What are the eigenvalues and eigenvectors of A ? Can A be diagonalized? Hint: Consider what a reflection does to specific vectors.

Solution:

Let v_1 be a non-zero vector on the line g , that is, v_1 spans g . Then $t(v_1) = Av_1 = v_1$. Hence v_1 is an eigenvector for A (equivalently for t) with eigenvalue 1.

Let v_w be a non-zero vector orthogonal to g . Then $t(v_w) = Av_w = -v_w$. Hence v_w is an eigenvector for A (equivalently for t) with eigenvalue -1 .

Since A is a 2×2 -matrix and has at most 2 eigenvalues we found all of them. Since v_1 and v_w are non-zero and orthogonal, they form a basis $B = (v_1, v_w)$ of \mathbb{R}^2 . For P the matrix with columns v_1, v_w , we then have

$$A = P \cdot \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \cdot P^{-1}.$$

Note that P is the change of coordinates matrix $[id]_{B,E}$ and $[t]_{B,B} = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$. So that's exactly how we computed $[t]_{E,E} = A$ earlier. \square

- (4) Consider a population of owls feeding on a population of squirrels. In month k , let o_k denote the number of owls and s_k the number of squirrels. Assume that the populations change every month as follows:

$$\begin{aligned} o_{k+1} &= 0.3o_k + 0.4s_k \\ s_{k+1} &= -0.4o_k + 1.3s_k \end{aligned}$$

That is, if there would be no squirrels to hunt, only 30% of the owls would survive to the next month; if there were no owls that hunted squirrels, then the squirrel population would grow by factor 1.3 every month.

Let $x_k = \begin{bmatrix} o_k \\ s_k \end{bmatrix}$. Express the population change from x_k to x_{k+1} using a matrix A . Diagonalize A .

Solution:

$$x_{k+1} = \underbrace{\begin{bmatrix} 0.3 & 0.4 \\ -0.4 & 1.3 \end{bmatrix}}_A x_k$$

We diagonalize A . The characteristic equation is

$$0 = \det(A - \lambda I) = (0.3 - \lambda)(1.3 - \lambda) + 0.4^2 = \lambda^2 - 1.6\lambda + 0.55,$$

the eigenvalues are $\lambda = \frac{1}{2}(1.6 \pm \sqrt{1.6^2 - 4 \cdot 0.55}) = 0.8 \pm 0.3 \in \{0.5, 1.1\}$. We compute a basis for each eigenspace.

$$\lambda = 0.5 : \quad \text{Nul}(A - 0.5I) = \text{Nul} \begin{bmatrix} -0.2 & 0.4 \\ -0.4 & 0.8 \end{bmatrix} = \text{Nul} \begin{bmatrix} 1 & -2 \\ 0 & 0 \end{bmatrix} = \text{Span} \left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix} \right\}.$$

$$\lambda = 1.1 : \quad \text{Nul}(A - 1.1I) = \text{Nul} \begin{bmatrix} -0.8 & 0.4 \\ -0.4 & 0.2 \end{bmatrix} = \text{Nul} \begin{bmatrix} 1 & -1/2 \\ 0 & 0 \end{bmatrix} = \text{Span} \left\{ \begin{bmatrix} 1/2 \\ 1 \end{bmatrix} \right\}.$$

We write the eigenvectors in a matrix P and compute P^{-1} :

$$P = \begin{bmatrix} 2 & 1/2 \\ 1 & 1 \end{bmatrix}, \quad P^{-1} = \frac{1}{3/2} \begin{bmatrix} 1 & -1/2 \\ -1 & 2 \end{bmatrix} = \frac{1}{3} \begin{bmatrix} 2 & -1 \\ -2 & 4 \end{bmatrix}.$$

We obtain a diagonalization

$$A = \underbrace{\begin{bmatrix} 2 & 1/2 \\ 1 & 1 \end{bmatrix}}_P \underbrace{\begin{bmatrix} 0.5 & 0 \\ 0 & 1.1 \end{bmatrix}}_D \underbrace{\frac{1}{3} \begin{bmatrix} 2 & -1 \\ -2 & 4 \end{bmatrix}}_{P^{-1}}.$$

\square

- (5) Continue the previous problem: Let the starting population be $x_1 = \begin{bmatrix} o_1 \\ s_1 \end{bmatrix} = \begin{bmatrix} 20 \\ 100 \end{bmatrix}$.
- Give an explicit formula for the populations in month $k + 1$.
 - Are the populations growing or decreasing over time? By which factor?

- (c) What is ratio of owls to squirrels after 12 months? After 24 months? Can you explain why?

Solution:

- (a) (2 points)

$$\begin{aligned} x_{k+1} = A^k x_1 = PD^k P^{-1} x_1 &= \begin{bmatrix} 2 & 1/2 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 0.5^k & 0 \\ 0 & 1.1^k \end{bmatrix} \frac{1}{3} \begin{bmatrix} 2 & -1 \\ -2 & 4 \end{bmatrix} \begin{bmatrix} 20 \\ 100 \end{bmatrix} \\ &= \begin{bmatrix} 60 \cdot 1.1^k - 40 \cdot 0.5^k \\ 120 \cdot 1.1^k - 20 \cdot 0.5^k \end{bmatrix} \end{aligned}$$

- (b) (2 points) Both populations are growing. For large k , the term 0.5^k can be neglected (e.g. for $k \geq 12$ we have $1.1^k \geq 3.138$ and $0.5^k \leq 0.00025$). We can approximate the populations by

$$x_{k+1} \approx \begin{bmatrix} 60 \cdot 1.1^k \\ 120 \cdot 1.1^k \end{bmatrix} = 1.1^k \begin{bmatrix} 60 \\ 120 \end{bmatrix} \quad \text{for large } k.$$

After a large number of months, both populations grow by a factor of 1.1 every month.

- (c) (1 point) The populations are $x_{13} = \begin{bmatrix} 188.3 \\ 376.6 \end{bmatrix}$ after 12 months and $x_{25} = \begin{bmatrix} 591.0 \\ 1182.0 \end{bmatrix}$ after 24 months. After a large number of months, the ratio of owls to squirrels is always about 1 : 2 by the approximation formula for x_{k+1} . □

- (6) (a) Give 3 vectors of length 1 in \mathbb{R}^3 that are orthogonal to $u = \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix}$.
 (b) Which of the following sets are orthogonal? Orthonormal?

$$A = \left\{ \begin{bmatrix} 0.6 \\ 0.8 \end{bmatrix}, \begin{bmatrix} 0.8 \\ -0.6 \end{bmatrix} \right\}, \quad B = \left\{ \frac{1}{3} \begin{bmatrix} 1 \\ -2 \\ 2 \end{bmatrix}, \frac{1}{\sqrt{18}} \begin{bmatrix} 4 \\ 1 \\ -1 \end{bmatrix} \right\}$$

Solution:

- (a) Switch any 2 components of u , change one sign and set the remaining component 0 to get orthogonal vectors, e.g., $\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$, $\begin{bmatrix} 0 \\ 2 \\ -1 \end{bmatrix}$, $\begin{bmatrix} 0 \\ 2 \\ 1 \end{bmatrix}$.

- (b) A is orthonormal since its vectors are pairwise orthogonal and all have length 1. Same for B . □