

Math 2135 - Assignment 13

Due April 26, 2019

- (1) Are the matrices A, B, C, D in (3), (4), (5) of assignment 12 diagonalizable? How?
- (2) Let A be an $n \times n$ -matrix. Are the following true or false? Explain why:
 - (a) If A has n eigenvectors, then A is diagonalizable.
 - (b) If a 4×4 -matrix A has two eigenvalues with eigenspaces of dimension 3 and 1, respectively, then A is diagonalizable.
 - (c) A is diagonalizable iff A has n eigenvalues (counting multiplicities).
 - (d) If \mathbb{R}^n has a basis of eigenvectors of A , then A is diagonalizable.
- (3) Let A be the standard matrix for the reflection t of \mathbb{R}^2 on some line g through the origin. What are the eigenvalues and eigenvectors of A ? Can A be diagonalized? Hint: Consider what a reflection does to specific vectors.
- (4) Consider a population of owls feeding on a population of squirrels. In month k , let o_k denote the number of owls and s_k the number of squirrels. Assume that the populations change every month as follows:

$$\begin{aligned}o_{k+1} &= 0.3o_k + 0.4s_k \\s_{k+1} &= -0.4o_k + 1.3s_k\end{aligned}$$

That is, if there would be no squirrels to hunt, only 30% of the owls would survive to the next month; if there were no owls that hunted squirrels, then the squirrel population would grow by factor 1.3 every month.

Let $x_k = \begin{bmatrix} o_k \\ s_k \end{bmatrix}$. Express the population change from x_k to x_{k+1} using a matrix A . Diagonalize A .

- (5) Continue the previous problem: Let the starting population be $x_1 = \begin{bmatrix} o_1 \\ s_1 \end{bmatrix} = \begin{bmatrix} 20 \\ 100 \end{bmatrix}$.
 - (a) Give an explicit formula for the populations in month $k + 1$.
 - (b) Are the populations growing or decreasing over time? By which factor?
 - (c) What is ratio of owls to squirrels after 12 months? After 24 months? Can you explain why?
- (6) (a) Give 3 vectors of length 1 in \mathbb{R}^3 that are orthogonal to $u = \begin{bmatrix} 1 \\ -1 \\ 2 \end{bmatrix}$.
(b) Which of the following are orthonormal sets?

$$A = \left\{ \begin{bmatrix} 0.6 \\ 0.8 \end{bmatrix}, \begin{bmatrix} 0.8 \\ -0.6 \end{bmatrix} \right\}, \quad B = \left\{ \frac{1}{3} \begin{bmatrix} 1 \\ -2 \\ 2 \end{bmatrix}, \frac{1}{\sqrt{18}} \begin{bmatrix} 4 \\ 1 \\ -1 \end{bmatrix} \right\}$$

- (7) (a) Let W be the subspace of \mathbb{R}^3 with orthonormal basis $B = \left(\frac{1}{3} \begin{bmatrix} 2 \\ -1 \\ 2 \end{bmatrix}, \frac{1}{\sqrt{5}} \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix} \right)$.
Compute the coordinates $[x]_B$ for $x = \begin{bmatrix} 7 \\ 4 \\ 4 \end{bmatrix}$ using dot products.
 - (b) Find the closest point to $y = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$ in W . What is the distance from y to W ?
- (8) True or false. Explain your answers.
 - (a) Every orthogonal set is also orthonormal.
 - (b) Not every orthonormal set in \mathbb{R}^n is linearly independent.
 - (c) For each x and each subspace W , the vector $x - \text{proj}_W(x)$ is orthogonal to W .