

Due Tuesday, September 5th at the beginning of class. Show all work and write solution on separate paper, treating this as a cover sheet.

Use integration by parts to find the following:

- $\int_4^9 \frac{\ln y}{\sqrt{y}} dy$. With $u = \ln y$, $dv = y^{-1/2} dy$, we have $du = \frac{dy}{y}$, $v = 2y^{1/2}$ and the integral becomes

$$2\sqrt{y} \ln y \Big|_4^9 - \int_4^9 2\sqrt{y} \frac{1}{y} dy = 2\sqrt{y} \ln y - 4\sqrt{y} \Big|_4^9 = (6 \ln 9 - 12) - (4 \ln 4 - 8) = \ln \left(\frac{531441}{256} \right) - 4.$$

- $\int \frac{x}{e^{2x}} dx$. With $u = x$, $dv = e^{-2x} dx$, we have $du = dx$, $v = -\frac{1}{2}e^{-2x}$ and the integral becomes

$$-\frac{x}{2}e^{-2x} + \frac{1}{2} \int e^{-2x} dx = -\frac{e^{-2x}}{2}(x + 1/2).$$

- $\int t^2 \sin(\pi t) dt$. With $u = t^2$, $dv = \sin(\pi t) dt$, we have $du = 2t dt$, $v = -\frac{1}{\pi} \cos(\pi t)$ and the integral becomes

$$-\frac{t^2}{\pi} \cos(\pi t) + \frac{2}{\pi} \int t \cos(\pi t) dt.$$

Using integration by parts again on the new integral ($u = t$, $dv = \cos(\pi t) dt$, $du = dt$, $v = \sin(\pi t)/\pi$) we get

$$-\frac{t^2}{\pi} \cos(\pi t) + \frac{2}{\pi} \left[\frac{t}{\pi} \sin(\pi t) - \frac{1}{\pi} \int \sin(\pi t) dt \right] = -\frac{t^2}{\pi} \cos(\pi t) + \frac{2t}{\pi^2} \sin(\pi t) + \frac{2}{\pi^3} \cos(\pi t).$$

- $\int \cos(\sqrt{x}) dx$ (Make a substitution first.) With the substitution

$$y = \sqrt{x}, \quad dy = \frac{dx}{2\sqrt{x}}, \quad dx = 2\sqrt{x} dy = 2y dy,$$

the integral becomes $2 \int y \cos y dy$. Using integration by parts with $u = y$, $dv = \cos y dy$, $du = dy$, $v = \sin y$, this integral is

$$2y \sin y - 2 \int \sin y dy = 2y \sin y + 2 \cos y = 2\sqrt{x} \sin(\sqrt{x}) + 2 \cos(\sqrt{x}).$$

- $\int e^{-x} \cos(\pi x) dx$. With $u = e^{-x}$, $dv = \cos(\pi x) dx$, we have $du = -e^{-x} dx$, $v = \frac{1}{\pi} \sin(\pi x)$ and the integral becomes

$$\frac{e^{-x}}{\pi} \sin(\pi x) + \frac{1}{\pi} \int e^{-x} \sin(\pi x) dx.$$

Integrating by parts again allows us to solve algebraically for the integral with which we started. Letting $u = e^{-x}$, $dv = \sin(\pi x) dx$, $du = -e^{-x}$, $v = -\cos(\pi x)/\pi$ we have

$$\int e^{-x} \cos(\pi x) dx = \frac{e^{-x}}{\pi} \sin(\pi x) + \frac{1}{\pi} \left[-\frac{e^{-x}}{\pi} \cos(\pi x) - \frac{1}{\pi} \int e^{-x} \cos(\pi x) dx \right],$$

so that

$$\int e^{-x} \cos(\pi x) dx = \frac{e^{-x}}{1 + \pi^2} (\pi \sin(\pi x) - \cos(\pi x)).$$

- $\int (\ln x)^2 dx$. With $u = (\ln x)^2$, $dv = dx$, $du = 2 \ln x \frac{dx}{x}$, $v = x$, the integral is

$$x(\ln x)^2 - 2 \int \ln x dx = x(\ln x)^2 - 2(x \ln x - x),$$

the second equality coming from memorization or integration by parts again.

- $\int \arccos z dz$. Trying $u = \arccos z$, $dv = dz$, we get $du = \frac{-dz}{\sqrt{1-z^2}}$, $v = z$ and the integral becomes

$$z \arccos z + \int \frac{z dz}{\sqrt{1-z^2}}.$$

We now make a substitution, $u = 1 - z^2$, $du = -2z dz$ to get

$$z \arccos z - \frac{1}{2} \int u^{-1/2} du = z \arccos z - \sqrt{1-z^2}.$$

- $\int \sin(\ln w) dw$ (Make a substitution first.) With the substitution

$$z = \ln w, dz = dw/w, dw = w dz = e^z dz,$$

the integral becomes

$$\int e^z \sin z dz = \frac{e^z}{2} (\sin z - \cos z) = \frac{w}{2} (\sin(\ln w) - \cos(\ln w)),$$

the first equality coming from the trick used above (integration by parts twice, solve algebraically for the integral).

- $\int_1^{\sqrt{3}} \arctan(1/x) dx$. With $u = \arctan(1/x)$, $dv = dx$, $du = \frac{-dx}{1+x^2}$, $v = x$, the integral becomes

$$x \arctan(1/x) \Big|_1^{\sqrt{3}} + \int \frac{x dx}{1+x^2}.$$

Now make the substitution $u = 1 + x^2$, $du = 2x dx$, to get

$$x \arctan(1/x) \Big|_1^{\sqrt{3}} + \frac{1}{2} \int_2^4 \frac{du}{u} = \sqrt{3} \arctan(1/\sqrt{3}) - \arctan(1) + \frac{1}{2} (\ln 2) = \frac{\pi}{2\sqrt{3}} - \frac{\pi}{4} + \frac{1}{2} \ln 2.$$

- $\int x^3 \sqrt{1+x^2} dx$. [Solution 1.] With $u = x^2$, $dv = x\sqrt{1+x^2}$, $du = 2xdx$,
 $v = \int x\sqrt{1+x^2} dx = \frac{1}{3}(1+x^2)^{3/2}$ (using a substitution), we get

$$\frac{x^2}{3}(1+x^2)^{3/2} - \frac{1}{3} \int 2x(1+x^2)^{3/2} dx = \frac{x^2}{3}(1+x^2)^{3/2} - \frac{2}{15}(1+x^2)^{5/2},$$

using a substitution in the second integral. [Solution 2.] Instead of doing integration by parts, one can directly make the substitution

$$u = 1 + x^2, \quad du = 2xdx, \quad x^2 = u - 1,$$

to get

$$\frac{1}{2} \int (u-1)\sqrt{u} du = \frac{u^{5/2}}{5} - \frac{u^{3/2}}{3} = \frac{1}{5}(1+x^2)^{5/2} - \frac{1}{3}(1+x^2)^{3/2}.$$

The solutions don't immediately look the same, but they are both equal to

$$(1+x^2)^{3/2} \frac{3x^2 - 2}{15}.$$

- $\int \sec^3 \theta d\theta$. With $u = \sec \theta$, $dv = \sec^2 \theta d\theta$, $du = \sec \theta \tan \theta d\theta$, $v = \tan \theta$, we get

$$\sec \theta \tan \theta - \int \tan^2 \theta \sec \theta d\theta.$$

Using the Pythagorean identity $1 + \tan^2 \theta = \sec^2 \theta$ in the integrand, we get

$$\int \sec^3 \theta d\theta = \sec \theta \tan \theta - \int (\sec^2 \theta - 1) \sec \theta d\theta = \sec \theta \tan \theta - \int \sec^3 \theta d\theta + \int \sec \theta d\theta.$$

Solving algebraically for $\int \sec^3 \theta d\theta$ and recalling

$$\int \sec \theta d\theta = \int \sec \theta \frac{\sec \theta + \tan \theta}{\sec \theta + \tan \theta} = \ln |\sec \theta + \tan \theta|$$

(using the substitution $u = \sec \theta + \tan \theta$) gives

$$\int \sec^3 \theta d\theta = \frac{1}{2} [\sec \theta \tan \theta + \ln |\sec \theta + \tan \theta|].$$

- Show that

$$\int (\ln x)^n dx = x(\ln x)^n - n \int (\ln x)^{n-1} dx.$$

With $u = (\ln x)^n$, $dv = dx$, $du = n(\ln x)^{n-1} \frac{dx}{x}$, $v = x$, we get

$$x(\ln x)^n - n \int (\ln x)^{n-1} dx.$$