

1. What is a sequence? What does it mean for a sequence to converge? Give an example of a convergent sequence and of a divergent sequence.

A sequence $(a_n)_n$ is a list of (real) numbers (in a particular order) indexed by natural numbers. A sequence converges,

$$\lim_{n \rightarrow \infty} a_n = L,$$

if there is a real number L which the a_n are approaching as n increases, i.e. a_n becomes arbitrarily close to L for n sufficiently large.

Here are some convergent sequences and their limits:

$$(a_n)_{n=1}^{\infty} = \left(\frac{1}{n} \right)_{n=1}^{\infty} = \left(1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \frac{1}{5}, \dots \right),$$

$$\lim_{n \rightarrow \infty} \frac{1}{n} = 0$$

$$(b_n)_{n=0}^{\infty} = \left(\cos \left(\frac{\pi}{2^n} \right) \right)_{n=0}^{\infty} = \left(-1, 0, \frac{1}{\sqrt{2}}, \frac{\sqrt{2+\sqrt{2}}}{2}, \dots \right),$$

$$\lim_{n \rightarrow \infty} \cos \left(\frac{\pi}{2^n} \right) = \cos \left(\lim_{n \rightarrow \infty} \frac{\pi}{2^n} \right) = \cos(0) = 1,$$

$$(c_n)_{n=1}^{\infty} = \left(1 + \frac{(-1)^n}{n} \right)_{n=1}^{\infty} = \left(0, \frac{3}{2}, \frac{2}{3}, \frac{5}{4}, \frac{4}{5}, \dots \right),$$

$$\lim_{n \rightarrow \infty} 1 + \frac{(-1)^n}{n} = 1.$$

Here are some divergent sequences:

$$(a_n)_{n=0}^{\infty} = (2^n)_{n=0}^{\infty} = (1, 2, 4, 8, 16, \dots)$$

exponential growth,

$$(b_n)_{n=0}^{\infty} = \left(\sin \left(\frac{n\pi}{3} \right) \right)_{n=0}^{\infty} = \left(0, \frac{\sqrt{3}}{2}, \frac{\sqrt{3}}{2}, 0, -\frac{\sqrt{3}}{2}, -\frac{\sqrt{3}}{2}, \dots \right)$$

periodic sequence,

$$(c_n)_{n=0}^{\infty} = ((-1)^n(n^2 + 1))_{n=0}^{\infty} = (1, -2, 5, -10, 17, \dots)$$

grows in absolute value, alternating in sign.

2. What is a series? What does it mean for a series to converge? Give an example of a convergent series and of a divergent series.

A series $\sum_{n=0}^{\infty} a_n$ is (a formal expression for) an attempt to add infinitely many numbers a_n , the terms of the series. A series is convergent if the limit of the partial sums

$$\lim_{N \rightarrow \infty} \sum_{n=1}^N a_n = \lim_{N \rightarrow \infty} (a_0 + a_1 + \dots + a_{N-1} + a_N)$$

exists. In other words, we add up finitely many terms (in order) from a given sequence a_n and see if the partial sums approach a limiting value.

Here are some convergent series:

$$\sum_{n=0}^{\infty} \frac{1}{2^n} = \frac{1}{1 - 1/2} = 2, \text{ a convergent geometric series, } r = 1/2, |r| < 1,$$

$$\sum_{n=1}^{\infty} \frac{1}{n(n+1)} = 1, \text{ a telescoping series, or convergent by comparison to } \sum_n \frac{1}{n^2},$$

$$\sum_{n=1}^{\infty} \frac{1}{n^2}, \text{ a convergent } p\text{-series, } p = 2 > 1.$$

Here are some divergent series:

$$\sum_{n=0}^{\infty} 2^n, \text{ a divergent geometric series, } r = 2, |r| \geq 1,$$

$$\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}}, \text{ a divergent } p\text{-series, } p = 1/2 \leq 1,$$

$$\sum_{n=0}^{\infty} \frac{2n^2 - 1}{3n^3 + 1}, \text{ divergent by comparison to the harmonic series } \sum_n \frac{1}{n}.$$

3. Use the integral test to determine the convergence or divergence of the series

$$\sum_{n=2}^{\infty} \frac{1}{n(\ln n)^2}.$$

The function $f(x) = \frac{1}{x(\ln x)^2}$ is positive and decreasing on the interval $[2, \infty)$ (positive because x and $(\ln x)^2$ are positive there, decreasing because the numerator is fixed and the denominator is increasing). The series therefore behaves like the improper integral

$$\begin{aligned} \int_2^{\infty} \frac{dx}{x(\ln x)^2} &= \lim_{T \rightarrow \infty} \int_2^T \frac{dx}{x(\ln x)^2} = \lim_{T \rightarrow \infty} \int_{\ln 2}^{\ln T} \frac{du}{u^2} \\ &= \lim_{T \rightarrow \infty} -\frac{1}{u} \Big|_{\ln 2}^{\ln T} = \lim_{T \rightarrow \infty} -\frac{1}{\ln T} + \frac{1}{\ln 2} = \frac{1}{\ln 2} < \infty. \end{aligned}$$

The improper integral converges, therefore the series converges as well.

4. Use the comparison test to determine the convergence or divergence of the series

$$\sum_{n=0}^{\infty} \frac{3^n - n^3}{5^n + n^5}.$$

We can use either the direct comparison test or the limit comparison test, in both cases comparing to the convergent geometric series $\sum_{n=0}^{\infty} (3/5)^n$.

For the direct comparison test, we have

$$0 \leq \frac{3^n - n^3}{5^n + n^5} \leq \left(\frac{3}{5}\right)^n,$$

so the series in the problem statement is dominated by the convergent geometric series $\sum_n (3/5)^n$.

For the limit comparison test, we have

$$\lim_{n \rightarrow \infty} \frac{\frac{3^n - n^3}{5^n + n^5}}{(3/5)^n} = \lim_{n \rightarrow \infty} \frac{1 - \frac{n^3}{3^n}}{1 + \frac{n^5}{5^n}} = 1$$

since

$$\lim_{n \rightarrow \infty} \frac{n^5}{5^n} = \lim_{n \rightarrow \infty} \frac{n^3}{3^n} = 0$$

e.g. using L'Hôpital's rule a few times. Therefore the series in the problem statement has the same convergence behavior as the convergent geometric series $\sum_n (3/5)^n$.