

Lecture 9: Section 2.8: What does f' say about f ?*Lecturer: Sarah Arpin***Today's Goal: What do f' , f'' tell us about f ?**


Logistics: Should be starting this on a Wednesday and finishing it on Friday.
We have a check-in on Friday that will cover 2.6 and 2.7.

Warm-Up 9.1 *True or False: If $f(x)$ is continuous at $x = a$, then $f(x)$ is differentiable at $x = a$.*

9.1 $f'(x)$


$f'(a)$ specifies the slope of the tangent line to $f(x)$ at $x = a$. If you're trying to "eyeball" the slope of a tangent line, it's hard to tell the difference between a slope of 2 and a slope of 2.5. But it's *not* difficult to tell the difference between a slope of 2 and a slope of -1 .

- If $f'(a) > 0$, then $f(x)$ is **increasing** at $x = a$.
- if $f'(x) > 0$ on an interval, then $f(x)$ is **increasing** on that interval.
- If $f'(a) < 0$, then $f(x)$ is **decreasing** at $x = a$.
- if $f'(x) < 0$ on an interval, then $f(x)$ is **decreasing** on that interval.



Definition 9.2 *A point where $f'(x) = 0$ or $f'(x)$ DNE is a potential spot where f could be changing from increasing to decreasing or vice versa! An x -value c such that $f'(c) = 0$ or $f'(c)$ DNE is called a **critical point**.*

9.1.1 Local Minima and Maxima




Definition 9.3 We say that $f(x)$ has a **local maximum** at $x = c$ if $f(x) \leq f(c)$ for every x in an open interval around $x = c$. This is also referred to as a *relative maximum*.

A local maximum can also be characterized by considering $f'(x)$:

Definition 9.4 We say that $f(x)$ has a **local minimum** at $x = c$ if $f(x) \geq f(c)$ for every x in an open interval around $x = c$. This is also referred to as a *relative minimum*.


A local minimum can also be characterized by considering $f'(x)$:

Example 9.5 The graph of $f'(x)$ is shown here.



1. On what interval(s) is $f(x)$ increasing?
2. On what interval(s) is $f(x)$ decreasing?
3. At what values of x does $f(x)$ have local maxima and minima?

9.2 $f''(x)$



In a previous project, we noticed that $f(x)$ is **concave up** when $f'(x)$ is increasing, and $f(x)$ is **concave down** when $f'(x)$ is decreasing. We can also characterize this using $f''(x)$:

- If $f''(a) > 0$, then $f(x)$ is **concave up** at $x = a$.
- If $f''(x) > 0$ on an interval, then $f(x)$ is **concave up** on that interval.
- If $f''(a) < 0$, then $f(x)$ is **concave down** at $x = a$.
- If $f''(x) < 0$ on an interval, then $f(x)$ is **concave down** on that interval.

Remark 9.6 $f''(x) > 0$ is saying the same thing as ‘ $f'(x)$ is increasing’!
Likewise, $f''(x) < 0$ is the same thing as saying ‘ $f'(x)$ is decreasing’!


Definition 9.7 An **inflection point** is a point (x, y) where the function changes concavity.

Inflection points can only occur at x -values where $f''(x) = 0$ or DNE, but we don't use the word inflection point unless there is a *change* in concavity. For example, if $f''(x)$ goes from positive, to zero, to positive – there is no change in concavity, so this is not an inflection point. However, if $f''(x)$ goes from positive to zero to negative, then we have passed through an inflection point. In addition, it needs to be a *point* on the graph of $f(x)$, so the x -value of the location of change in concavity must be in the domain of $f(x)$.

Example 9.8 (Example due to Dr. Patrick Newberry) Sketch a graph with the given properties:

- $f'(x) > 0$ for $|x| < 2$
- $f'(x) < 0$ for $|x| > 2$
- $f'(2) = 0$
- $f''(x) < 0$ for $0 < x < 3$
- $\lim_{x \rightarrow \infty} f(x) = 1$
- $f(-x) = -f(x)$ for all x

Hint: Translate each of these bullet points to a statement about f ! Use sign lines for $f'(x)$, $f''(x)$ to keep track of the information.



9.3 Antiderivatives

Definition 9.9 A function $F(x)$ is an **antiderivative** of a function $f(x)$ if $F'(x) = f(x)$.

Big idea: If we are given some function which we *know* is a derivative, can we recover the original function? How close can we get?

Example 9.10 Suppose $f(x) = x^2 - 3x + 4$, and $F(x)$ is an antiderivative of $f(x)$.

1. On what interval(s) is $F(x)$ increasing?
2. On what interval(s) is $F(x)$ decreasing?
3. At what x -values does $F(x)$ have any inflection points, if any?
4. What is $F(0)$?