

Objectives:

- Define relationships between $f(x)$, $f'(x)$ and $f''(x)$.
- Use information from $f(x)$ to graph $f'(x)$.

What does $f(x)$ tell us about $f'(x)$?


If $f(x)$ is _____ at $x = a$, then $f'(a)$ is _____ .

If $f(x)$ is _____ at $x = a$, then $f'(a)$ is _____ .


If $f(x)$ has a _____ at $x = a$, then $f'(a) =$ _____ .

Note: If $f(x)$ is discontinuous at a , has a corner/cusp at a , or has a vertical tangent line at a , then $f'(a)$ is undefined.

What does $f(x)$ tell us about $f''(x)$?



Concave _____



Concave _____

If $f(x)$ is _____ , $f'(x)$ is _____ , so $f''(x)$ is _____ .

If $f(x)$ is _____ , $f'(x)$ is _____ , so $f''(x)$ is _____ .


Summary

First, look for points where the derivative or second derivative is zero. Then consider where $f'(x)$ and $f''(x)$ are positive or negative, according to the following patterns:

$f(x)$		
$f'(x)$		

$f(x)$		
$f'(x)$		
$f''(x)$		

Example:



	$x \in (1, 3)$	$x = 3$	$x \in (3, 6)$	$x = 6$	$x \in (6, 7)$
$f(x)$	increasing				
$f'(x)$	+				

	$x \in (1, 3)$	$x \in (3, 5)$	$x \in (5, 7)$
$f(x)$	concave down		
$f'(x)$	decreasing		
$f''(x)$			

We'll use these basic rules in today's class activity. The solutions to the activity will be posted on the course website - I would recommend adding at least some of those examples to your notes.