

Math 2300, Midterm 3

November 16, 2015

PRINT YOUR NAME: _____

PRINT INSTRUCTOR'S NAME: _____

Mark your section/instructor:

<input type="checkbox"/>	Section 001	Albert Bronstein	9:00 - 9:50
<input type="checkbox"/>	Section 002	Andrew Healy	10:00 - 10:50
<input type="checkbox"/>	Section 003	Joshua Frinak	11:00 - 11:50
<input type="checkbox"/>	Section 004	Kevin Berg	12:00 - 12:50
<input type="checkbox"/>	Section 005	Jeffrey Shriner	2:00 - 2:50
<input type="checkbox"/>	Section 006	Megan Ly	3:00 - 3:50
<input type="checkbox"/>	Section 007	Albert Bronstein	8:00 - 8:50
<input type="checkbox"/>	Section 008	Jonathan Lamar	1:00 - 1:50
<input type="checkbox"/>	Section 009	Keli Parker	3:00 - 3:50
<input type="checkbox"/>	Section 010	Steven Weinell	4:00 - 4:50
<input type="checkbox"/>	Section 011	Benjamin Cooper	8:00 - 8:50
<input type="checkbox"/>	Section 880	Jordan Watts	8:00 - 8:50

Question	Points	Score
1	8	
2	12	
3	12	
4	8	
5	12	
6	12	
7	12	
8	8	
9	6	
10	10	
Total:	100	

- No calculators or cell phones or other electronic devices allowed at any time.
- Show all your reasoning and work for full credit. Use full mathematical or English sentences.
- You have 90 minutes and the exam is 100 points.
- You do not need to simplify numerical expressions. For example leave fractions like $100/7$ or expressions like $\ln(3)/2$ as is.
- When done, give your exam to your instructor, who will mark your name off on a photo roster.
- We hope you show us your best work!

1. (8 points) Match the following functions with their corresponding Maclaurin series:

(a) $e^{x^2/2} = \underline{\hspace{2cm}}$

I. $\sum_{n=0}^{\infty} x^{2n}$

(b) $\cos\left(\frac{x}{2}\right) = \underline{\hspace{2cm}}$

II. $\sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{2^{2n} (2n)!}$

(c) $\frac{1}{(1-x)^2} = \underline{\hspace{2cm}}$

IV. $\sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+2}}{2n+1}$

(d) $x \arctan(x) = \underline{\hspace{2cm}}$

V. $\sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+2}}{(2n+1)!}$

VI. $\sum_{n=0}^{\infty} \frac{x^{2n}}{2^n n!}$

2. (12 points) (a) Find the radius of convergence of the power series $\sum_{n=1}^{\infty} \frac{(x-5)^n}{2^n n^2}$. Show all work in justifying your answer.

(b) Find the interval of convergence. Show all work in justifying your answer.

3. (12 points) Find the solution of the differential equation

$$y(x + 1) + y' = 0$$

that satisfies the initial condition $y(-2) = 1$. **Show all your work and write the solution on the line given below.**

Solution: $y =$ _____

4. (8 points) Given the following power series $\sum_{n=0}^{\infty} a_n(x-2)^n$ we know that at $x = 0$ the series converges and at $x = 8$ the series diverges. What do we know about the following values?

(a) At $x = 3$ the series $\sum_{n=0}^{\infty} a_n(x-2)^n$ is:

(i) convergent

(ii) divergent

(iii) We cannot determine its convergence/divergence with the information given.

(b) At $x = -4$ the series $\sum_{n=0}^{\infty} a_n(x-2)^n$ is:

(i) convergent

(ii) divergent

(iii) We cannot determine its convergence/divergence with the information given.

(c) At $x = 9$ the series $\sum_{n=0}^{\infty} a_n(x-2)^n$ is:

(i) convergent

(ii) divergent

(iii) We cannot determine its convergence/divergence with the information given.

(d) The following series $\sum_{n=0}^{\infty} a_n$ is:

(i) convergent

(ii) divergent

(iii) We cannot determine its convergence/divergence with the information given.

5. (12 points) (a) Write the definition for the n^{th} degree Taylor polynomial of $f(x)$ centered at $x = a$.

(b) Find the second degree Taylor polynomial for $f(x) = \ln(\sec(x))$ centered at $\frac{\pi}{4}$.

6. (12 points) (a) Express the function $f(x) = \ln(1 + x^3)$ as a power series centered about $x = 0$.

(b) Express the definite integral as an infinite series.

$$\int_0^1 \ln(x^3 + 1) dx$$

7. (12 points) (a) Fill in the boxes to complete the statement of **Taylor's Inequality**:

If $\leq M$ on the interval between the center, a , and the point of approximation, x , then the remainder, $R_n(x)$, of the n^{th} degree Taylor polynomial, $T_n(x)$, satisfies the inequality:

$$|R_n(x)| \leq$$

(b) Use Taylor's Inequality to determine the number of terms of the Maclaurin series for e^x that should be used to estimate the number e with an error less than 0.6. Clearly justify your choice of M .





8. (8 points) Each of the following slope fields represents one of the following differential equations. Match each slope field to the corresponding differential equation.

(a) $\frac{dy}{dx} = \frac{xy}{2}$ _____

(b) $\frac{dy}{dx} = y - x - 2$ _____

(c) $\frac{dy}{dx} = x + 2$ _____

(d) $\frac{dy}{dx} = e^{x^2}$ _____



9. (6 points) Find the sum of the series.

$$\sum_{n=0}^{\infty} \frac{(-1)^n}{2n+1} = 1 - \frac{1}{3} + \frac{1}{5} - \frac{1}{7} + \frac{1}{9} - \cdots$$

10. (10 points) Assume we approximate the sum of the series

$$\sum_{n=0}^{\infty} \frac{2}{n^2}$$

by using the first 3 terms. Give an upper bound for the error involved in this approximation by using the Remainder Estimate for the Integral Test.