

## Exercise 26.21

### Abstract Algebra 1 MATH 3140

SEBASTIAN CASALAINA

ABSTRACT. This is Exercise 26.21 from Fraleigh [Fra03, §26]:

**Exercise 26.21.** Let  $R$  and  $R'$  be rings and let  $\phi : R \rightarrow R'$  be a ring homomorphism such that  $\phi[R] \neq \{0'\}$ . Show that if  $R$  has unity 1 and  $R'$  has no 0 divisors, then  $\phi(1)$  is unity for  $R'$ .

*Solution.* Let  $R$  and  $R'$  be rings and let  $\phi : R \rightarrow R'$  be a ring homomorphism such that  $\phi[R] \neq \{0'\}$ . Suppose further that  $R$  has unity 1, and *assume only the weaker hypothesis* that  $\phi(1)$  is not a zero divisor in  $R'$  (which of course is true if  $R'$  has no 0 divisors). We will show that  $\phi(1)$  is unity for  $R'$ .

The first observation is that

$$(0.1) \quad \phi(1) = \phi(1 \cdot 1) = \phi(1)\phi(1),$$

since  $\phi$  is a homomorphism of rings. The second observation is that  $\phi(1) \neq 0'$ , since otherwise, for any  $r \in R$  we would have  $\phi(r) = \phi(1 \cdot r) = \phi(1)\phi(r) = 0' \cdot r = 0'$ , contradicting the assumption that  $\phi[R] \neq \{0'\}$ .

Now, with these observations, to show that  $\phi(1)$  is a left multiplicative identity, we want to show that for all  $r' \in R'$ , we have

$$(0.2) \quad \phi(1)r' = r'.$$

This is equivalent to showing

$$(0.3) \quad \phi(1)r' - r' = 0,$$

which, since  $\phi(1) \neq 0'$  and is not a zero divisor, is equivalent to showing

$$(0.4) \quad \phi(1)(\phi(1)r' - r') = 0.$$

In other words, to show that  $\phi(1)$  is a left multiplicative identity for  $R'$ , it suffices to show that (0.4) holds for all  $r' \in R$ . We now prove this:

$$\begin{aligned} \phi(1)(\phi(1)r' - r') &= \phi(1)\phi(1)r' - \phi(1)r' \\ &= \phi(1)r' - \phi(1)r' && \text{(using (0.1))} \\ &= 0. \end{aligned}$$

This completes the proof that  $\phi(1)$  is a left multiplicative identity. A similar argument shows that it is also a right multiplicative identity, and therefore unity in  $R'$ .  $\square$

*Remark 0.1.* We also have the related result: if  $R$  is a ring with unity 1 and  $\phi : R \rightarrow R'$  is a surjective ring homomorphism, then  $\phi(1)$  is unity for  $R'$ . Indeed, for any  $r' \in R'$ , there exists  $r \in R$  such that  $\phi(r) = r'$ , and then we have  $\phi(1)r' = \phi(1)\phi(r) = \phi(1 \cdot r) = \phi(r) = r'$ , and similarly, one can show  $r'\phi(1) = r'$ .

## REFERENCES

[Fra03] John Fraleigh, *A First Course in Abstract Algebra*, Seventh edition, Addison Wesley, Pearson, 2003.

UNIVERSITY OF COLORADO, DEPARTMENT OF MATHEMATICS, CAMPUS BOX 395, BOULDER, CO 80309

*Email address:* casa@math.colorado.edu