

Midterm I

Intro to Discrete Math

MATH 2001

Spring 2022

Friday February 11, 2022

NAME: _____

PRACTICE EXAM

SOLUTIONS

Question:	1	2	3	4	5	Total
Points:	25	25	25	5	20	100
Score:						

- The exam is closed book. You **may not use any resources** whatsoever, other than paper, pencil, and pen, to complete this exam.
- You **may not discuss the exam** with anyone except me, in any way, under any circumstances.
- You **must explain your answers**, and you will be **graded on the clarity of your solutions**.
- You must upload your exam as a single **.pdf** to **Canvas**, with the questions in the correct order, etc.
- You have 45 minutes to complete the exam. **We will spend the last 5 minutes of class to upload your exam to Canvas.**

1. • Consider the sets $A = \{3, 9\}$ and $B = \{2, 3, 5\}$.

For this problem **you do not need to justify your answer.**

- (a) (5 points) List the elements of the power set $\mathcal{P}(A)$.

SOLUTION:

$\emptyset, \{3\}, \{9\}, \{3, 9\}$

- (b) (5 points) List the elements of the set $A \times B$.

SOLUTION:

$(3, 2), (3, 3), (3, 5), (9, 2), (9, 3), (9, 5)$.

- (c) (5 points) List the elements of the set $A \cup B$.

SOLUTION:

$2, 3, 5, 9$

- (d) (5 points) Is it true that $2 \in (A \cap B)$?

SOLUTION:

No

- (e) (5 points) List the elements of the set $B - A$.

SOLUTION:

$2, 5$

1
25 points

2. (25 points) • Suppose that A and B are finite sets. What is $|A \times B|$ in terms of $|A|$ and $|B|$? Explain.

SOLUTION:

Solution.

$$|A \times B| = |A||B|$$

Indeed, recall that the product $A \times B$ is the set of ordered pairs:

$$A \times B := \{(a, b) : a \in A, b \in B\}.$$

There are $|A|$ choices for the first entry of an ordered pair (a, b) in the product $A \times B$, and then for each choice of first entry there are $|B|$ choices for the second entry, so that all together, we have $|A||B|$ ordered pairs in the product $A \times B$. □

You can find a similar explanation on p.9 of Hammack, where this is Fact 1.1.

Also, as an example to illustrate the formula, suppose that $A = \{a_1, a_2, a_3\}$ and $B = \{b_1, b_2\}$. Then we have

$$A \times B = \{(a_1, b_1), (a_1, b_2), (a_2, b_1), (a_2, b_2), (a_3, b_1), (a_3, b_2)\},$$

and so we can count that $|A \times B| = 6 = 3 \cdot 2 = |A||B|$, confirming the general formula in this special case. (See also Problem 1(b) for a similar example.)

2
25 points

3. (25 points) • For each t in the interval $[0, 1] \subseteq \mathbb{R}$, consider the set

$$A_t = \{(x, y) \in \mathbb{R}^2 : t \leq x \leq t + 1 \text{ and } y = t\}.$$


Describe the union

$$\bigcup_{t \in [0, 1]} A_t$$

as a geometric object in the plane \mathbb{R}^2 . A good picture and a brief explanation of your solution is sufficient for this problem.

SOLUTION:

Solution. For each $t \in [0, 1]$, the set A_t is the horizontal line segment joining the points (t, t) and $(t + 1, t)$.



Therefore, the union of the A_t over all t in the interval $[0, 1]$ is the region bounded by the parallelogram with vertices $(0, 0)$, $(0, 1)$, $(2, 1)$, and $(1, 1)$. □

3
25 points

4. (5 points) • Write the \LaTeX code that will produce the following:

$$A \in \mathcal{P}(X) \iff A \subseteq X$$

SOLUTION:

Solution. `A \in \mathscr{P}(X) \iff A \subseteq X`

□

4

5 points

5. • **True or False.** For this problem you do not need to justify your answer.

- (a) (4 points) **True or False** (circle one). If A and B are finite sets and $A \cap B = \emptyset$, then $|A \cup B| = |A| + |B|$.

SOLUTION: TRUE. More generally, for finite sets A and B , we have $|A \cup B| = |A| + |B| - |A \cap B|$.

- (b) (4 points) **True or False** (circle one). If A and B are subsets of a set X , then, regarding complements of subsets of X , we have $(A \cap B)^C = A^C \cup B^C$.

SOLUTION: TRUE. We explained this in class.

- (c) (4 points) **True or False** (circle one). The negation of the statement $\forall x \in X, \exists y \in Y, p(x, y)$ is logically equivalent to the statement $\exists x \in X, \exists y \in Y, \sim p(x, y)$.

SOLUTION: FALSE. The negation is logically equivalent to $\exists x \in X, \forall y \in Y, \sim p(x, y)$. To see that $\exists x \in X, \exists y \in Y, \sim p(x, y)$ is NOT the correct negation, consider the example where $X = \mathbb{Z} - \{0\}$, $Y = \mathbb{Q}$, and for $x \in X$ and $y \in Y$, we take $p(x, y)$ to be the statement $xy = 1$ (then both the statement $\forall x \in X, \exists y \in Y, p(x, y)$ and the claimed negation, $\exists x \in X, \exists y \in Y, \sim p(x, y)$, would be true, which is not possible).

- (d) (4 points) **True or False** (circle one). The truth table for the statement $p \wedge q$ is

p	q	$p \wedge q$
T	T	T
T	F	F
F	T	T
F	F	F

SOLUTION: FALSE. The third row is not correct: if p is false and q is true, then $p \wedge q$ is false.

- (e) (4 points) **True or False** (circle one). $(\mathbb{R} - \mathbb{Z}) \times \mathbb{N} = (\mathbb{R} \times \mathbb{N}) - (\mathbb{Z} \times \mathbb{N})$.

SOLUTION: TRUE. If A , B , and C are sets, we have $(A - B) \times C = (A \times C) - (B \times C)$.

5

20 points
