

Exact Sequences, Hom Functors, Tensor Product, II

Commutative Algebra

Sep 21, 2009

Background: Adjoints

Defn. A pair of functors $\mathcal{C} \begin{smallmatrix} F \\ \rightleftarrows \\ G \end{smallmatrix} \mathcal{D}$ is an *adjoint pair* if there is a natural isomorphism

$$\mathrm{Hom}_{\mathcal{C}}(F(X), Y) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{D}}(X, G(Y)).$$

F is a *left adjoint* and G is a *right adjoint*.

Background: Adjoint

Defn. A pair of functors $\mathcal{C} \begin{smallmatrix} F \\ \rightleftarrows \\ G \end{smallmatrix} \mathcal{D}$ is an *adjoint pair* if there is a natural isomorphism

$$\mathrm{Hom}_{\mathcal{C}}(F(X), Y) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{D}}(X, G(Y)).$$

F is a *left adjoint* and G is a *right adjoint*.

Example. $G(Y) = \mathrm{Hom}_R(M, Y)$ and $F(X) = _ \otimes_R M$.

This follows from the definition of \otimes :

$$\mathrm{Hom}_R(L \otimes_R M, N) \xrightarrow{\sim} \mathrm{Hom}_R(L, \mathrm{Hom}_R(M, N)).$$

Background: Adjoints

Defn. A pair of functors $\mathcal{C} \xrightleftharpoons[G]{F} \mathcal{D}$ is an *adjoint pair* if there is a natural isomorphism

$$\mathrm{Hom}_{\mathcal{C}}(F(X), Y) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{D}}(X, G(Y)).$$

F is a *left adjoint* and G is a *right adjoint*.

Example. $G(Y) = \mathrm{Hom}_R(M, Y)$ and $F(X) = _ \otimes_R M$.

This follows from the definition of \otimes :

$$\mathrm{Hom}_R(L \otimes_R M, N) \xrightarrow{\sim} \mathrm{Hom}_R(L, \mathrm{Hom}_R(M, N)).$$

Background Info.

Background: Adjoints

Defn. A pair of functors $\mathcal{C} \xrightleftharpoons[G]{F} \mathcal{D}$ is an *adjoint pair* if there is a natural isomorphism

$$\mathrm{Hom}_{\mathcal{C}}(F(X), Y) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{D}}(X, G(Y)).$$

F is a *left adjoint* and G is a *right adjoint*.

Example. $G(Y) = \mathrm{Hom}_R(M, Y)$ and $F(X) = _ \otimes_R M$.

This follows from the definition of \otimes :

$$\mathrm{Hom}_R(L \otimes_R M, N) \xrightarrow{\sim} \mathrm{Hom}_R(L, \mathrm{Hom}_R(M, N)).$$

Background Info.

- A functor between preadditive categories that has an adjoint must be additive.

Background: Adjoint

Defn. A pair of functors $\mathcal{C} \xrightleftharpoons[G]{F} \mathcal{D}$ is an *adjoint pair* if there is a natural isomorphism

$$\mathrm{Hom}_{\mathcal{C}}(F(X), Y) \xrightarrow{\sim} \mathrm{Hom}_{\mathcal{D}}(X, G(Y)).$$

F is a *left adjoint* and G is a *right adjoint*.

Example. $G(Y) = \mathrm{Hom}_R(M, Y)$ and $F(X) = _ \otimes_R M$.

This follows from the definition of \otimes :

$$\mathrm{Hom}_R(L \otimes_R M, N) \xrightarrow{\sim} \mathrm{Hom}_R(L, \mathrm{Hom}_R(M, N)).$$

Background Info.

- A functor between preadditive categories that has an adjoint must be additive.
- A functor with a left adjoint must be left exact. A functor with a right adjoint must be right exact.

\otimes as a Functor

If $\varphi: A \rightarrow B$ and $\psi: C \rightarrow D$ are R -module homomorphisms, then

$$A \times C \xrightarrow{\varphi \times \psi} B \times D \rightarrow B \otimes D : (a, c) \mapsto \varphi(a) \otimes \psi(c)$$

is R -bilinear, hence induces an R -module homomorphism

$$\varphi \otimes \psi: A \otimes C \rightarrow B \otimes D : a \otimes c \mapsto \varphi(a) \otimes \psi(c).$$

\otimes as a Functor

If $\varphi: A \rightarrow B$ and $\psi: C \rightarrow D$ are R -module homomorphisms, then

$$A \times C \xrightarrow{\varphi \times \psi} B \times D \rightarrow B \otimes D : (a, c) \mapsto \varphi(a) \otimes \psi(c)$$

is R -bilinear, hence induces an R -module homomorphism

$$\varphi \otimes \psi: A \otimes C \rightarrow B \otimes D : a \otimes c \mapsto \varphi(a) \otimes \psi(c).$$

$_ \otimes M$ becomes a functor by defining $\varphi \otimes M = \varphi \otimes \text{id}_M$ whenever $A \xrightarrow{\varphi} B$.

Some Isomorphisms Involving \otimes

Some Isomorphisms Involving \otimes

$$(i) \operatorname{Hom}_R(L_R, \operatorname{Hom}_S({}_R M_S, N_S)) \cong \operatorname{Hom}_S(L \otimes_R M_S, N_S)$$

Some Isomorphisms Involving \otimes

(i) $\text{Hom}_R(L_R, \text{Hom}_S({}_R M_S, N_S)) \cong \text{Hom}_S(L \otimes_R M_S, N_S)$
 $\varphi(x)(y) = \varphi(x, y) \mapsto \bar{\varphi}$ (give by the universal property of \otimes).

Some Isomorphisms Involving \otimes

(i) $\text{Hom}_R(L_R, \text{Hom}_S({}_R M_S, N_S)) \cong \text{Hom}_S(L \otimes_R M_S, N_S)$
 $\varphi(x)(y) = \varphi(x, y) \mapsto \overline{\varphi}$ (give by the universal property of \otimes).

(ii) $M \otimes_R N \cong N \otimes_R M$

Some Isomorphisms Involving \otimes

(i) $\text{Hom}_R(L_R, \text{Hom}_S({}_R M_S, N_S)) \cong \text{Hom}_S(L \otimes_R M_S, N_S)$
 $\varphi(x)(y) = \varphi(x, y) \mapsto \bar{\varphi}$ (give by the universal property of \otimes).

(ii) $M \otimes_R N \cong N \otimes_R M$

$m \otimes n \mapsto n \otimes m$

Some Isomorphisms Involving \otimes

(i) $\text{Hom}_R(L_R, \text{Hom}_S({}_R M_S, N_S)) \cong \text{Hom}_S(L \otimes_R M_S, N_S)$
 $\varphi(x)(y) = \varphi(x, y) \mapsto \bar{\varphi}$ (give by the universal property of \otimes).

(ii) $M \otimes_R N \cong N \otimes_R M$

$m \otimes n \mapsto n \otimes m$

(iii) $L \otimes_R (M \otimes_S N) \cong (L \otimes_R M) \otimes_S N$

Some Isomorphisms Involving \otimes

(i) $\text{Hom}_R(L_R, \text{Hom}_S({}_R M_S, N_S)) \cong \text{Hom}_S(L \otimes_R M_S, N_S)$
 $\varphi(x)(y) = \varphi(x, y) \mapsto \bar{\varphi}$ (give by the universal property of \otimes).

(ii) $M \otimes_R N \cong N \otimes_R M$

$$m \otimes n \mapsto n \otimes m$$

(iii) $L \otimes_R (M \otimes_S N) \cong (L \otimes_R M) \otimes_S N$

$$l \otimes (m \otimes n) \mapsto (l \otimes m) \otimes n$$

Some Isomorphisms Involving \otimes

(i) $\text{Hom}_R(L_R, \text{Hom}_S({}_R M_S, N_S)) \cong \text{Hom}_S(L \otimes_R M_S, N_S)$
 $\varphi(x)(y) = \varphi(x, y) \mapsto \bar{\varphi}$ (give by the universal property of \otimes).

(ii) $M \otimes_R N \cong N \otimes_R M$

$$m \otimes n \mapsto n \otimes m$$

(iii) $L \otimes_R (M \otimes_S N) \cong (L \otimes_R M) \otimes_S N$

$$l \otimes (m \otimes n) \mapsto (l \otimes m) \otimes n$$

(iv) $M \otimes_R \left(\bigoplus_{i \in I} N_i \right) \cong \bigoplus_{i \in I} (M \otimes N_i)$

Some Isomorphisms Involving \otimes

(i) $\text{Hom}_R(L_R, \text{Hom}_S({}_R M_S, N_S)) \cong \text{Hom}_S(L \otimes_R M_S, N_S)$
 $\varphi(x)(y) = \varphi(x, y) \mapsto \bar{\varphi}$ (give by the universal property of \otimes).

(ii) $M \otimes_R N \cong N \otimes_R M$

$$m \otimes n \mapsto n \otimes m$$

(iii) $L \otimes_R (M \otimes_S N) \cong (L \otimes_R M) \otimes_S N$

$$l \otimes (m \otimes n) \mapsto (l \otimes m) \otimes n$$

(iv) $M \otimes_R (\bigoplus_{i \in I} N_i) \cong \bigoplus_{i \in I} (M \otimes N_i)$

$$m \otimes (n_i)_{i \in I} \mapsto (m \otimes n_i)_{i \in I}$$

Some Isomorphisms Involving \otimes

(i) $\text{Hom}_R(L_R, \text{Hom}_S({}_R M_S, N_S)) \cong \text{Hom}_S(L \otimes_R M_S, N_S)$
 $\varphi(x)(y) = \varphi(x, y) \mapsto \bar{\varphi}$ (give by the universal property of \otimes).

(ii) $M \otimes_R N \cong N \otimes_R M$

$$m \otimes n \mapsto n \otimes m$$

(iii) $L \otimes_R (M \otimes_S N) \cong (L \otimes_R M) \otimes_S N$

$$l \otimes (m \otimes n) \mapsto (l \otimes m) \otimes n$$

(iv) $M \otimes_R (\bigoplus_{i \in I} N_i) \cong \bigoplus_{i \in I} (M \otimes N_i)$

$$m \otimes (n_i)_{i \in I} \mapsto (m \otimes n_i)_{i \in I}$$

(v) $R \otimes_R M \cong M$

Some Isomorphisms Involving \otimes

(i) $\text{Hom}_R(L_R, \text{Hom}_S({}_R M_S, N_S)) \cong \text{Hom}_S(L \otimes_R M_S, N_S)$
 $\varphi(x)(y) = \varphi(x, y) \mapsto \bar{\varphi}$ (give by the universal property of \otimes).

(ii) $M \otimes_R N \cong N \otimes_R M$

$$m \otimes n \mapsto n \otimes m$$

(iii) $L \otimes_R (M \otimes_S N) \cong (L \otimes_R M) \otimes_S N$

$$l \otimes (m \otimes n) \mapsto (l \otimes m) \otimes n$$

(iv) $M \otimes_R (\bigoplus_{i \in I} N_i) \cong \bigoplus_{i \in I} (M \otimes N_i)$

$$m \otimes (n_i)_{i \in I} \mapsto (m \otimes n_i)_{i \in I}$$

(v) $R \otimes_R M \cong M$

$$r \otimes m \mapsto rm$$

Some Isomorphisms Involving \otimes

(i) $\text{Hom}_R(L_R, \text{Hom}_S({}_R M_S, N_S)) \cong \text{Hom}_S(L \otimes_R M_S, N_S)$
 $\varphi(x)(y) = \varphi(x, y) \mapsto \bar{\varphi}$ (give by the universal property of \otimes).

(ii) $M \otimes_R N \cong N \otimes_R M$

$$m \otimes n \mapsto n \otimes m$$

(iii) $L \otimes_R (M \otimes_S N) \cong (L \otimes_R M) \otimes_S N$

$$l \otimes (m \otimes n) \mapsto (l \otimes m) \otimes n$$

(iv) $M \otimes_R (\bigoplus_{i \in I} N_i) \cong \bigoplus_{i \in I} (M \otimes N_i)$

$$m \otimes (n_i)_{i \in I} \mapsto (m \otimes n_i)_{i \in I}$$

(v) $R \otimes_R M \cong M$

$$r \otimes m \mapsto rm$$

(v)' If $I \triangleleft R$, then $R/I \otimes_R M \cong M/IM$

Some Isomorphisms Involving \otimes

(i) $\text{Hom}_R(L_R, \text{Hom}_S({}_R M_S, N_S)) \cong \text{Hom}_S(L \otimes_R M_S, N_S)$
 $\varphi(x)(y) = \varphi(x, y) \mapsto \bar{\varphi}$ (give by the universal property of \otimes).

(ii) $M \otimes_R N \cong N \otimes_R M$

$$m \otimes n \mapsto n \otimes m$$

(iii) $L \otimes_R (M \otimes_S N) \cong (L \otimes_R M) \otimes_S N$

$$l \otimes (m \otimes n) \mapsto (l \otimes m) \otimes n$$

(iv) $M \otimes_R (\bigoplus_{i \in I} N_i) \cong \bigoplus_{i \in I} (M \otimes N_i)$

$$m \otimes (n_i)_{i \in I} \mapsto (m \otimes n_i)_{i \in I}$$

(v) $R \otimes_R M \cong M$

$$r \otimes m \mapsto rm$$

(v)' If $I \triangleleft R$, then $R/I \otimes_R M \cong M/IM$

$$(r + I) \otimes m \mapsto rm + IM$$

Right Exactness of \otimes

Right Exactness of \otimes

Thm. Tensor functors are right exact.

Right Exactness of \otimes

Thm. Tensor functors are right exact.

Proof:

Right Exactness of \otimes

Thm. Tensor functors are right exact.

Proof: Assume $L \xrightarrow{\alpha} M \xrightarrow{\beta} N \longrightarrow 0$ is exact, and consider

$$L \otimes A \xrightarrow{\alpha \otimes \text{id}} M \otimes A \xrightarrow{\beta \otimes \text{id}} N \otimes A \longrightarrow 0.$$

Right Exactness of \otimes

Thm. Tensor functors are right exact.

Proof: Assume $L \xrightarrow{\alpha} M \xrightarrow{\beta} N \longrightarrow 0$ is exact, and consider

$$L \otimes A \xrightarrow{\alpha \otimes \text{id}} M \otimes A \xrightarrow{\beta \otimes \text{id}} N \otimes A \longrightarrow 0.$$

- $\beta \otimes \text{id}$ maps generators onto generators, so it is surjective.

Right Exactness of \otimes

Thm. Tensor functors are right exact.

Proof: Assume $L \xrightarrow{\alpha} M \xrightarrow{\beta} N \longrightarrow 0$ is exact, and consider

$$L \otimes A \xrightarrow{\alpha \otimes \text{id}} M \otimes A \xrightarrow{\beta \otimes \text{id}} N \otimes A \longrightarrow 0.$$

- $\beta \otimes \text{id}$ maps generators onto generators, so it is surjective.
- To show exactness at $M \otimes A$ it suffices to prove that the second map in

$$M \otimes A \xrightarrow{\nu} (M \otimes A)/I \xrightarrow{\overline{\beta \otimes \text{id}}} N \otimes A$$

is invertible, $I = \text{im}(\alpha \otimes \text{id})$. To construct an inverse $\overline{\psi}$, define

$\psi: N \times A \rightarrow (M \otimes A)/I$ by $\psi(n \otimes a) = m \otimes a + I$ for any $m \in \beta^{-1}(n)$. ψ is well-defined because $\ker(\beta) = \text{im}(\alpha)$ and $\text{im}(\alpha) \otimes A = I$. ψ is bilinear, hence extends to a map $\overline{\psi}$ of $N \otimes A$. Since $\overline{\psi} \circ \overline{\beta \otimes \text{id}}$ is the identity on elements $m \otimes a + I$, the map $\overline{\beta \otimes \text{id}}$ is 1-1.

Nonexactness of \otimes

Example. $0 \longrightarrow \mathbb{Z} \xrightarrow{\subseteq} \mathbb{Q} \xrightarrow{\nu} \mathbb{Q}/\mathbb{Z} \longrightarrow 0$ is an exact sequence of \mathbb{Z} -modules.

Nonexactness of \otimes

Example. $0 \longrightarrow \mathbb{Z} \xrightarrow{\subseteq} \mathbb{Q} \xrightarrow{\nu} \mathbb{Q}/\mathbb{Z} \longrightarrow 0$ is an exact sequence of \mathbb{Z} -modules.

If you apply $_ \otimes \mathbb{Z}_2$ you obtain $0 \longrightarrow \mathbb{Z}_2 \longrightarrow 0 \longrightarrow 0 \longrightarrow 0$.

Nonexactness of \otimes

Example. $0 \longrightarrow \mathbb{Z} \xrightarrow{\subseteq} \mathbb{Q} \xrightarrow{\nu} \mathbb{Q}/\mathbb{Z} \longrightarrow 0$ is an exact sequence of \mathbb{Z} -modules.

If you apply $_ \otimes \mathbb{Z}_2$ you obtain $0 \longrightarrow \mathbb{Z}_2 \longrightarrow 0 \longrightarrow 0 \longrightarrow 0$.

Defn. An R -module M is *flat* if $_ \otimes M$ is exact.

Nonexactness of \otimes

Example. $0 \longrightarrow \mathbb{Z} \xrightarrow{\subseteq} \mathbb{Q} \xrightarrow{\nu} \mathbb{Q}/\mathbb{Z} \longrightarrow 0$ is an exact sequence of \mathbb{Z} -modules.

If you apply $_ \otimes \mathbb{Z}_2$ you obtain $0 \longrightarrow \mathbb{Z}_2 \longrightarrow 0 \longrightarrow 0 \longrightarrow 0$.

Defn. An R -module M is *flat* if $_ \otimes M$ is exact. (equivalently if $_ \otimes M$ maps injections to injections).

Consequences

Consequences

- If F is free of rank κ , then $F \otimes M \cong \bigoplus^{\kappa} M$.

Consequences

- If F is free of rank κ , then $F \otimes M \cong \bigoplus^{\kappa} M$.
- A tensor product of two free modules is free.

Consequences

- If F is free of rank κ , then $F \otimes M \cong \bigoplus^{\kappa} M$.
- A tensor product of two free modules is free.
- a tensor product of two projective modules is projective.

Consequences

- If F is free of rank κ , then $F \otimes M \cong \bigoplus^{\kappa} M$.
- A tensor product of two free modules is free.
- a tensor product of two projective modules is projective.
- Free \implies projective \implies flat, and none of the arrows can be reversed in general.

Tensor Product of R -algebras

Defn. If A and B are commutative R -algebras, then $A \otimes_R B$ has an R -module structure: define $(a \otimes b)(a' \otimes b') = aa' \otimes bb'$.

Tensor Product of R -algebras

Defn. If A and B are commutative R -algebras, then $A \otimes_R B$ has an R -module structure: define $(a \otimes b)(a' \otimes b') = aa' \otimes bb'$.

- The maps $\iota_A: A \rightarrow A \otimes_R B: a \mapsto a \otimes 1$ and $\iota_B: B \rightarrow A \otimes_R B: b \mapsto 1 \otimes b$ are R -algebra homomorphisms.

Tensor Product of R -algebras

Defn. If A and B are commutative R -algebras, then $A \otimes_R B$ has an R -module structure: define $(a \otimes b)(a' \otimes b') = aa' \otimes bb'$.

- The maps $\iota_A: A \rightarrow A \otimes_R B: a \mapsto a \otimes 1$ and $\iota_B: B \rightarrow A \otimes_R B: b \mapsto 1 \otimes b$ are R -algebra homomorphisms.
- $(A \otimes_R B, \iota_A, \iota_B)$ is a coproduct of A and B in the category of (commutative) R -algebras.

Tensor Product of R -algebras

Defn. If A and B are commutative R -algebras, then $A \otimes_R B$ has an R -module structure: define $(a \otimes b)(a' \otimes b') = aa' \otimes bb'$.

- The maps $\iota_A: A \rightarrow A \otimes_R B: a \mapsto a \otimes 1$ and $\iota_B: B \rightarrow A \otimes_R B: b \mapsto 1 \otimes b$ are R -algebra homomorphisms.
- $(A \otimes_R B, \iota_A, \iota_B)$ is a coproduct of A and B in the category of (commutative) R -algebras.

Example. If X and Y are disjoint sets of indeterminates, then $R[X] \otimes_R R[Y] \cong R[X \cup Y]$.