

Quantifiers.

In these notes we will discuss how to determine the truth of formulas of the form

$$(\dagger) \quad \varphi = \forall x_1 \cdots \exists x_n F(x_1, \dots, x_n),$$

which consist of a string $\forall x_1 \cdots \exists x_n$ of quantified variables followed by a quantifier-free formula $F(x_1, \dots, x_n)$ whose variables are among those that have been quantified. So, we will discuss formulas like

$$\exists x (x = x^2) \quad \text{or} \quad \forall x \exists y (x < y) \quad \text{or} \quad \forall x \forall y \exists z ((x^2 + y^2 = z^2 + 1) \vee (x^2 + y^2 = z^2)),$$

but postpone the discussion of formulas like

$$\underbrace{\exists x (x + y = 0)}_{\text{unquantified variable } y} \quad \text{or} \quad \underbrace{\forall x ((0 < x) \rightarrow \exists y ((0 < y) \wedge (y < x)))}_{\text{one quantifier is not at the front}}.$$

When applied to ordered sets, the formula $\exists t \forall x (x \leq t)$ expresses the fact that there is a top element. Some ordered sets have a top element and others do not, so the truth of this formula depends on which structure is referred to. Let's define "structure":

Definition 1. A *structure* for a language L is a set equipped with a fixed interpretation of each of the nonlogical symbols of L .

Here are some examples of structures for different languages:

$$V = \langle \{\text{a universe of sets}\}; \in \rangle.$$

$$\mathbb{N} = \langle \{\text{natural numbers}\}; +, -, 0, \cdot, 1, S \rangle.$$

$$\mathbb{R} = \langle \{\text{real numbers}\}; +, -, 0, \cdot, 1, < \rangle.$$

Quantifier games. Given a formula φ of type (\dagger) and a structure $\mathbb{A} = \langle A; +, -, <, \dots \rangle$ in the same language as φ there is a game between the quantifiers \exists and \forall which decides the truth of φ in \mathbb{A} . In this game, \exists (also known as Eloise, or The Prover) tries to prove that φ is true in \mathbb{A} while \forall (also known as Abélard, or The Refuter) tries to prove that φ is false in \mathbb{A} .

The sequence of play is determined by the sequence of quantifiers in the formula. For example, if $\varphi = \forall x \exists y \exists z ((x < y) \wedge (z < x))$, then one play of the game involves one turn by \forall followed by two turns by \exists . A player takes a turn by selecting an element of the structure to substitute for the quantified variable. For example if $\mathbb{A} = \langle \mathbb{Z}; < \rangle$, then \forall may select $1 \in \mathbb{Z}$ to substitute for x , then \exists may select $0 \in \mathbb{Z}$ to substitute for y , and finally \exists may select $7 \in \mathbb{Z}$ to substitute for z . Now we delete the quantifiers from φ and make the substitutions: $((1 < 0) \wedge (7 < 1))$. This is false in the structure $\langle \mathbb{Z}; < \rangle$, so it counts as a win for The Refuter, \forall . (Note: each time a quantifier chooses a value to substitute for its variable, it is allowed to know all choices made previously by either player.)

The formula φ is false in \mathbb{A} if \forall has a *winning strategy*, which means a strategy to force a win every time the game is played. The formula is true in \mathbb{A} if \exists has a winning strategy. Although \forall won the instance of the game described in the previous paragraph, \forall does not

have a winning strategy; rather, \exists does. One strategy for \exists is: “whatever \forall chooses for the value of x , choose $y = x + 1$ and $z = x - 1$ ”.

Zermelo’s Theorem (from Game Theory) guarantees that one of \exists or \forall must have a winning strategy. It may not be easy to find a winning strategy. Indeed, a winning strategy for \exists is a proof that φ is true, while a winning strategy for \forall is a proof that φ is false, so finding winning strategies is the same thing as finding proofs for formal statements. This means that discovering winning strategies requires creativity and effort.

Exercises. Decide the truth or falsity of the given formula in the given structure. In each example, describe a winning strategy.

(1) $\forall x \exists y (x < y)$ in $\langle \mathbb{R}; < \rangle$.

(2) $\exists y \forall x (x < y)$ in $\langle \mathbb{R}; < \rangle$.

(3) $\forall x \forall y \exists z ((x < y) \rightarrow ((x < z) \wedge (z < y)))$ in $\langle \mathbb{R}; < \rangle$. What about $\langle \mathbb{N}; < \rangle$?

(4) $\forall w \exists x \forall y \exists z (wxyz = w + x + y + z)$, in \mathbb{R} .

(5) $\forall u \exists v \forall w \exists x \forall y \exists z ((u - v)(w - x)(y - z) = 1)$, in \mathbb{R} .

(6) Show that $f(x) = 2x$ is continuous at $x = 2$. That is, show that

$$\forall \varepsilon > 0 \exists \delta > 0 \forall x ((0 < |x - 2| < \delta) \rightarrow (|2x - 4| < \varepsilon))$$

is true in \mathbb{R} .

(7) Show that $f(x) = -3x$ is continuous at $x = 1$.

(8) Show that $f(x) = |x|$ is continuous at $x = 1$.

(9) Show that $f(x) = |x|$ is continuous at $x = 0$.

(10) Show that $f(x) = x^2$ is continuous at $x = 1$.