

# Examples, Properties, Applications

Math 6270

$G$	1	$k_2$	$\cdots$	$k_r$
	1	$g_2$	$\cdots$	$g_r$
$\chi_1$	1	1	$\cdots$	1
$\chi_2$	$d_2$	$\chi_2(g_2)$	$\cdots$	$\chi_2(g_r)$
$\vdots$	$\vdots$	$\vdots$	$\ddots$	$\vdots$
$\chi_r$	$d_r$	$\chi_r(g_2)$	$\cdots$	$\chi_r(g_r)$

Each  $d_i$  divides  $|G| = 1^2 + d_2^2 + \cdots + d_r^2$

Each  $k_j$  divides  $|G| = 1 + k_2 + \cdots + k_r$

# Character tables of abelian groups

# Character tables of abelian groups

$\mathbb{Z}_2$	1	1
	0	1
$\xi_1$	1	1
$\xi_2$	1	-1

# Character tables of abelian groups

$\mathbb{Z}_2$	1	1
	0	1
$\xi_1$	1	1
$\xi_2$	1	-1

$\mathbb{Z}_3$	1	1	1
	0	1	2
$\chi_1$	1	1	1
$\chi_2$	1	$\omega$	$\omega^2$
$\chi_2^2$	1	$\omega^2$	$\omega$

# Character tables of abelian groups

$\mathbb{Z}_2$	1	1
	0	1
$\xi_1$	1	1
$\xi_2$	1	-1

$\mathbb{Z}_3$	1	1	1
	0	1	2
$\chi_1$	1	1	1
$\chi_2$	1	$\omega$	$\omega^2$
$\chi_2^2$	1	$\omega^2$	$\omega$

$\mathbb{Z}_4$	1	1	1	1
	0	1	2	3
$\chi_1$	1	1	1	1
$\chi_2$	1	$i$	-1	$-i$
$\chi_2^2$	1	-1	1	-1
$\chi_2^3$	1	$-i$	-1	$i$

# Character tables of abelian groups

$\mathbb{Z}_2$	1	1
	0	1
$\xi_1$	1	1
$\xi_2$	1	-1

$\mathbb{Z}_3$	1	1	1
	0	1	2
$\chi_1$	1	1	1
$\chi_2$	1	$\omega$	$\omega^2$
$\chi_2^2$	1	$\omega^2$	$\omega$

$\mathbb{Z}_4$	1	1	1	1
	0	1	2	3
$\chi_1$	1	1	1	1
$\chi_2$	1	$i$	-1	$-i$
$\chi_2^2$	1	-1	1	-1
$\chi_2^3$	1	$-i$	-1	$i$

$\mathbb{Z}_2 \times \mathbb{Z}_2$	1	1	1	1
	(0, 0)	(0, 1)	(1, 0)	(1, 1)
$\xi_1(x)\xi_1(y)$	1	1	1	1
$\xi_2(x)\xi_1(y)$	1	1	-1	-1
$\xi_1(x)\xi_2(y)$	1	-1	1	-1
$\xi_2(x)\xi_2(y)$	1	-1	-1	1

# Completing a table from partial information

# Completing a table from partial information

Let  $G$  be a nonabelian group of order 8. Necessarily  $G/Z(G) \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ , so inflation gives us partial information about the character table.



# Completing a table from partial information

Let  $G$  be a nonabelian group of order 8. Necessarily  $G/Z(G) \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ , so inflation gives us partial information about the character table.

$G$	1	1	2	2	2
	1	$g_2$	$g_3$	$g_4$	$g_5$
$\chi_1$	1	1	1	1	1
$\chi_2$	1	1	1	-1	-1
$\chi_3$	1	1	-1	1	-1
$\chi_4$	1	1	-1	-1	1
$\chi_5$	?	?	?	?	?

# Completing a table from partial information

Let  $G$  be a nonabelian group of order 8. Necessarily  $G/Z(G) \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ , so inflation gives us partial information about the character table.

$G$	1	1	2	2	2
	1	$g_2$	$g_3$	$g_4$	$g_5$
$\chi_1$	1	1	1	1	1
$\chi_2$	1	1	1	-1	-1
$\chi_3$	1	1	-1	1	-1
$\chi_4$	1	1	-1	-1	1
$\chi_5$	?	?	?	?	?

The entries in the last row can be determined by:

# Completing a table from partial information

Let  $G$  be a nonabelian group of order 8. Necessarily  $G/Z(G) \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ , so inflation gives us partial information about the character table.

$G$	1	1	2	2	2
	1	$g_2$	$g_3$	$g_4$	$g_5$
$\chi_1$	1	1	1	1	1
$\chi_2$	1	1	1	-1	-1
$\chi_3$	1	1	-1	1	-1
$\chi_4$	1	1	-1	-1	1
$\chi_5$	?	?	?	?	?

The entries in the last row can be determined by:

(i) finding the missing irrep,

# Completing a table from partial information

Let  $G$  be a nonabelian group of order 8. Necessarily  $G/Z(G) \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ , so inflation gives us partial information about the character table.

$G$	1	1	2	2	2
	1	$g_2$	$g_3$	$g_4$	$g_5$
$\chi_1$	1	1	1	1	1
$\chi_2$	1	1	1	-1	-1
$\chi_3$	1	1	-1	1	-1
$\chi_4$	1	1	-1	-1	1
$\chi_5$	?	?	?	?	?

The entries in the last row can be determined by:

(i) finding the missing irrep, (ii) column orthogonality,

# Completing a table from partial information

Let  $G$  be a nonabelian group of order 8. Necessarily  $G/Z(G) \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ , so inflation gives us partial information about the character table.

$G$	1	1	2	2	2
	1	$g_2$	$g_3$	$g_4$	$g_5$
$\chi_1$	1	1	1	1	1
$\chi_2$	1	1	1	-1	-1
$\chi_3$	1	1	-1	1	-1
$\chi_4$	1	1	-1	-1	1
$\chi_5$	?	?	?	?	?

The entries in the last row can be determined by:

(i) finding the missing irrep, (ii) column orthogonality, (iii) row orthogonality,

# Completing a table from partial information

Let  $G$  be a nonabelian group of order 8. Necessarily  $G/Z(G) \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ , so inflation gives us partial information about the character table.

$G$	1	1	2	2	2
	1	$g_2$	$g_3$	$g_4$	$g_5$
$\chi_1$	1	1	1	1	1
$\chi_2$	1	1	1	-1	-1
$\chi_3$	1	1	-1	1	-1
$\chi_4$	1	1	-1	-1	1
$\chi_5$	?	?	?	?	?

The entries in the last row can be determined by:

- (i) finding the missing irrep, (ii) column orthogonality, (iii) row orthogonality,
- (iv)  $\sum_i d_i \chi_i(h) = \chi_{\text{reg}}(h)$ ,

# Completing a table from partial information

Let  $G$  be a nonabelian group of order 8. Necessarily  $G/Z(G) \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ , so inflation gives us partial information about the character table.

$G$	1	1	2	2	2
	1	$g_2$	$g_3$	$g_4$	$g_5$
$\chi_1$	1	1	1	1	1
$\chi_2$	1	1	1	-1	-1
$\chi_3$	1	1	-1	1	-1
$\chi_4$	1	1	-1	-1	1
$\chi_5$	?	?	?	?	?

The entries in the last row can be determined by:

- (i) finding the missing irrep, (ii) column orthogonality, (iii) row orthogonality,
- (iv)  $\sum_i d_i \chi_i(h) = \chi_{\text{reg}}(h)$ , ETC.

# Completing a table from partial information, $Q_8, D_4$



# Completing a table from partial information, $Q_8, D_4$

The table for either of  $D_4(D_8)$  or  $Q_8$  is:

# Completing a table from partial information, $Q_8, D_4$

The table for either of  $D_4(D_8)$  or  $Q_8$  is:

$G$	1	1	2	2	2
	1	$g_2$	$g_3$	$g_4$	$g_5$
$\chi_1$	1	1	1	1	1
$\chi_2$	1	1	1	-1	-1
$\chi_3$	1	1	-1	1	-1
$\chi_4$	1	1	-1	-1	1
$\chi_5$	2	-2	0	0	0

# Completing a table from partial information, $Q_8, D_4$

The table for either of  $D_4(D_8)$  or  $Q_8$  is:

$G$	1	1	2	2	2
	1	$g_2$	$g_3$	$g_4$	$g_5$
$\chi_1$	1	1	1	1	1
$\chi_2$	1	1	1	-1	-1
$\chi_3$	1	1	-1	1	-1
$\chi_4$	1	1	-1	-1	1
$\chi_5$	2	-2	0	0	0

The groups  $D_4$  and  $Q_8$  can be distinguished by the fact that  $\det(\chi_5^{Q_8}) = \chi_1 \neq \det(\chi_5^{D_4})$ .



$A_4/K \cong \mathbb{Z}_3$ , so 3 linear characters arise from inflation.

$A_4/K \cong \mathbb{Z}_3$ , so 3 linear characters arise from inflation. Also,  $A_4$  acts 2-transitively on  $S = \{1, 2, 3, 4\}$ , so  $\chi_4 = \chi_S - \chi_1$  is an irrep of degree 3.

$A_4/K \cong \mathbb{Z}_3$ , so 3 linear characters arise from inflation. Also,  $A_4$  acts 2-transitively on  $S = \{1, 2, 3, 4\}$ , so  $\chi_4 = \chi_S - \chi_1$  is an irrep of degree 3. This must be all.

$A_4/K \cong \mathbb{Z}_3$ , so 3 linear characters arise from inflation. Also,  $A_4$  acts 2-transitively on  $S = \{1, 2, 3, 4\}$ , so  $\chi_4 = \chi_S - \chi_1$  is an irrep of degree 3. This must be all.

$A_4$	1	3	4	4
	1	(1 2)(3 4)	(1 2 3)	(1 3 2)
$\chi_1$	1	1	1	1
$\chi_2$	1	1	$\omega$	$\omega^2$
$\chi_3$	1	1	$\omega^2$	$\omega$
$\chi_4$	3	-1	0	0



$A_4/K \cong \mathbb{Z}_3$ , so 3 linear characters arise from inflation. Also,  $A_4$  acts 2-transitively on  $S = \{1, 2, 3, 4\}$ , so  $\chi_4 = \chi_S - \chi_1$  is an irrep of degree 3. This must be all.

$A_4$	1	3	4	4
	1	(1 2)(3 4)	(1 2 3)	(1 3 2)
$\chi_1$	1	1	1	1
$\chi_2$	1	1	$\omega$	$\omega^2$
$\chi_3$	1	1	$\omega^2$	$\omega$
$\chi_4$	3	-1	0	0

Another way to produce  $\chi_4$  is to realize  $A_4$  as the rotation group of the tetrahedron.

$A_4/K \cong \mathbb{Z}_3$ , so 3 linear characters arise from inflation. Also,  $A_4$  acts 2-transitively on  $S = \{1, 2, 3, 4\}$ , so  $\chi_4 = \chi_S - \chi_1$  is an irrep of degree 3. This must be all.

$A_4$	1	3	4	4
	1	(1 2)(3 4)	(1 2 3)	(1 3 2)
$\chi_1$	1	1	1	1
$\chi_2$	1	1	$\omega$	$\omega^2$
$\chi_3$	1	1	$\omega^2$	$\omega$
$\chi_4$	3	-1	0	0

Another way to produce  $\chi_4$  is to realize  $A_4$  as the rotation group of the tetrahedron. Or use orthogonality.

$A_4/K \cong \mathbb{Z}_3$ , so 3 linear characters arise from inflation. Also,  $A_4$  acts 2-transitively on  $S = \{1, 2, 3, 4\}$ , so  $\chi_4 = \chi_S - \chi_1$  is an irrep of degree 3. This must be all.

$A_4$	1	3	4	4
	1	(1 2)(3 4)	(1 2 3)	(1 3 2)
$\chi_1$	1	1	1	1
$\chi_2$	1	1	$\omega$	$\omega^2$
$\chi_3$	1	1	$\omega^2$	$\omega$
$\chi_4$	3	-1	0	0

Another way to produce  $\chi_4$  is to realize  $A_4$  as the rotation group of the tetrahedron. Or use orthogonality. Or use the regular representation.



$S_4/K \cong S_3$ , so get 3 irreps by inflation.

$S_4/K \cong S_3$ , so get 3 irreps by inflation.  $S_4$  acts 2-transitively on  $\{1, 2, 3, 4\}$ , so get a degree 3 irrep from that,  $\chi_4$ .

$S_4/K \cong S_3$ , so get 3 irreps by inflation.  $S_4$  acts 2-transitively on  $\{1, 2, 3, 4\}$ , so get a degree 3 irrep from that,  $\chi_4$ . By orthogonality (or by realizing  $S_4$  as the rotation group of the cube) we get another degree 3 irrep,  $\chi_5$ .

$S_4/K \cong S_3$ , so get 3 irreps by inflation.  $S_4$  acts 2-transitively on  $\{1, 2, 3, 4\}$ , so get a degree 3 irrep from that,  $\chi_4$ . By orthogonality (or by realizing  $S_4$  as the rotation group of the cube) we get another degree 3 irrep,  $\chi_5$ .

$S_4$	1	3	8	4	4
	1	(1 2)(3 4)	(1 2 3)	(1 2)	(1 2 3 4)
$\chi_1$	1	1	1	1	1
$\chi_2$	1	1	1	-1	-1
$\chi_3$	2	2	-1	0	0
$\chi_4$	3	-1	0	1	-1
$\chi_5$	3	-1	0	-1	1





$A_5$  is the smallest nonabelian simple group, which makes it an interesting example.

$A_5$  is the smallest nonabelian simple group, which makes it an interesting example.  $\chi_1$  is the only irrep of degree 1.

$A_5$  is the smallest nonabelian simple group, which makes it an interesting example.  $\chi_1$  is the only irrep of degree 1.  $A_5$  acts 2-transitively on  $\{1, 2, 3, 4, 5\}$ , yielding an irrep of degree 4.

$A_5$  is the smallest nonabelian simple group, which makes it an interesting example.  $\chi_1$  is the only irrep of degree 1.  $A_5$  acts 2-transitively on  $\{1, 2, 3, 4, 5\}$ , yielding an irrep of degree 4.  $A_5$  acts 2-transitively by conjugation on its six Sylow 5-subgroups, yielding an irrep of degree 5.

$A_5$  is the smallest nonabelian simple group, which makes it an interesting example.  $\chi_1$  is the only irrep of degree 1.  $A_5$  acts 2-transitively on  $\{1, 2, 3, 4, 5\}$ , yielding an irrep of degree 4.  $A_5$  acts 2-transitively by conjugation on its six Sylow 5-subgroups, yielding an irrep of degree 5.  $A_5$  may be realized as the rotation group of the dodecahedron, yielding an irrep of degree 3.

$A_5$  is the smallest nonabelian simple group, which makes it an interesting example.  $\chi_1$  is the only irrep of degree 1.  $A_5$  acts 2-transitively on  $\{1, 2, 3, 4, 5\}$ , yielding an irrep of degree 4.  $A_5$  acts 2-transitively by conjugation on its six Sylow 5-subgroups, yielding an irrep of degree 5.  $A_5$  may be realized as the rotation group of the dodecahedron, yielding an irrep of degree 3. (Char value of a rotation in  $\mathbb{R}^3$  through  $2\pi/n$  is  $1 + 2\cos(2\pi/n)$ .)

$A_5$  is the smallest nonabelian simple group, which makes it an interesting example.  $\chi_1$  is the only irrep of degree 1.  $A_5$  acts 2-transitively on  $\{1, 2, 3, 4, 5\}$ , yielding an irrep of degree 4.  $A_5$  acts 2-transitively by conjugation on its six Sylow 5-subgroups, yielding an irrep of degree 5.  $A_5$  may be realized as the rotation group of the dodecahedron, yielding an irrep of degree 3. (Char value of a rotation in  $\mathbb{R}^3$  through  $2\pi/n$  is  $1 + 2\cos(2\pi/n)$ .)

$A_5$	1	15	20	12	12
	1	(1 2)(3 4)	(1 2 3)	(1 2 3 4 5)	(1 2 3 5 4)
$\chi_1$	1	1	1	1	1
$\chi_2$	3	-1	0	$\phi$	$-\phi^{-1}$
$\chi_3$	3	-1	0	$-\phi^{-1}$	$\phi$
$\chi_4$	4	0	1	-1	-1
$\chi_5$	5	1	-1	0	0



$A_5$  is the smallest nonabelian simple group, which makes it an interesting example.  $\chi_1$  is the only irrep of degree 1.  $A_5$  acts 2-transitively on  $\{1, 2, 3, 4, 5\}$ , yielding an irrep of degree 4.  $A_5$  acts 2-transitively by conjugation on its six Sylow 5-subgroups, yielding an irrep of degree 5.  $A_5$  may be realized as the rotation group of the dodecahedron, yielding an irrep of degree 3. (Char value of a rotation in  $\mathbb{R}^3$  through  $2\pi/n$  is  $1 + 2\cos(2\pi/n)$ .)

$A_5$	1	15	20	12	12
	1	(1 2)(3 4)	(1 2 3)	(1 2 3 4 5)	(1 2 3 5 4)
$\chi_1$	1	1	1	1	1
$\chi_2$	3	-1	0	$\phi$	$-\phi^{-1}$
$\chi_3$	3	-1	0	$-\phi^{-1}$	$\phi$
$\chi_4$	4	0	1	-1	-1
$\chi_5$	5	1	-1	0	0

$\chi_2$	$1 + 2\cos(0)$	$1 + 2\cos(\pi)$	$1 + 2\cos(2\pi/3)$	$1 + 2\cos(2\pi/5)$	$1 + 2\cos(4\pi/5)$
----------	----------------	------------------	---------------------	---------------------	---------------------

## Theorem

*If  $|G| = p$ ,  $p$  prime, then  $G$  is abelian.*

# Silly applications

## Theorem

*If  $|G| = p$ ,  $p$  prime, then  $G$  is abelian.*

## Proof.

# Silly applications

## Theorem

*If  $|G| = p$ ,  $p$  prime, then  $G$  is abelian.*

## Proof.

# Silly applications

## Theorem

*If  $|G| = p$ ,  $p$  prime, then  $G$  is abelian.*

## Proof.

The first column of the character table for  $G$  consists of at most  $p$  integers  $d_1, \dots, d_r$  such that (i)  $d_1 = 1$ , (ii)  $d_j | p$  for all  $j$ , and (iii)  $d_1^2 + \dots + d_r^2 = p$ .

# Silly applications

## Theorem

*If  $|G| = p$ ,  $p$  prime, then  $G$  is abelian.*

## Proof.

The first column of the character table for  $G$  consists of at most  $p$  integers  $d_1, \dots, d_r$  such that (i)  $d_1 = 1$ , (ii)  $d_j | p$  for all  $j$ , and (iii)  $d_1^2 + \dots + d_r^2 = p$ . It must be that  $d_1 = \dots = d_r = 1$ , so  $G$  is abelian. □

# Silly applications

## Theorem

*If  $|G| = p$ ,  $p$  prime, then  $G$  is abelian.*

## Proof.

The first column of the character table for  $G$  consists of at most  $p$  integers  $d_1, \dots, d_r$  such that (i)  $d_1 = 1$ , (ii)  $d_j | p$  for all  $j$ , and (iii)  $d_1^2 + \dots + d_r^2 = p$ . It must be that  $d_1 = \dots = d_r = 1$ , so  $G$  is abelian. □

## Theorem

*Any group of order  $< 6$  is abelian.*

# Silly applications

## Theorem

*If  $|G| = p$ ,  $p$  prime, then  $G$  is abelian.*

## Proof.

The first column of the character table for  $G$  consists of at most  $p$  integers  $d_1, \dots, d_r$  such that (i)  $d_1 = 1$ , (ii)  $d_j | p$  for all  $j$ , and (iii)  $d_1^2 + \dots + d_r^2 = p$ . It must be that  $d_1 = \dots = d_r = 1$ , so  $G$  is abelian.  $\square$

## Theorem

*Any group of order  $< 6$  is abelian.*

## Proof.

Same idea.



# Silly applications

## Theorem

*If  $|G| = p$ ,  $p$  prime, then  $G$  is abelian.*

## Proof.

The first column of the character table for  $G$  consists of at most  $p$  integers  $d_1, \dots, d_r$  such that (i)  $d_1 = 1$ , (ii)  $d_j | p$  for all  $j$ , and (iii)  $d_1^2 + \dots + d_r^2 = p$ . It must be that  $d_1 = \dots = d_r = 1$ , so  $G$  is abelian.  $\square$

## Theorem

*Any group of order  $< 6$  is abelian.*

## Proof.

Same idea.

# Silly applications

## Theorem

*If  $|G| = p$ ,  $p$  prime, then  $G$  is abelian.*

## Proof.

The first column of the character table for  $G$  consists of at most  $p$  integers  $d_1, \dots, d_r$  such that (i)  $d_1 = 1$ , (ii)  $d_j | p$  for all  $j$ , and (iii)  $d_1^2 + \dots + d_r^2 = p$ . It must be that  $d_1 = \dots = d_r = 1$ , so  $G$  is abelian.  $\square$

## Theorem

*Any group of order  $< 6$  is abelian.*

## Proof.

Same idea. The smallest number  $n$  that is a sum of squares  $d_1^2 + \dots + d_r^2$  where  $d_1 = 1$ , some  $d_j > 1$ , and all  $d_j$  divide  $n$  is 6.  $\square$

# More silly applications

## Theorem

*If  $|G| = p^2$ ,  $p$  prime, then  $G$  is abelian.*

# More silly applications

## Theorem

*If  $|G| = p^2$ ,  $p$  prime, then  $G$  is abelian.*

## Proof.

# More silly applications

## Theorem

*If  $|G| = p^2$ ,  $p$  prime, then  $G$  is abelian.*

## Proof.

# More silly applications

## Theorem

*If  $|G| = p^2$ ,  $p$  prime, then  $G$  is abelian.*

## Proof.

Each degree  $d_j$  must be  $1, p$  or  $p^2$ .

# More silly applications

## Theorem

*If  $|G| = p^2$ ,  $p$  prime, then  $G$  is abelian.*

## Proof.

Each degree  $d_j$  must be 1,  $p$  or  $p^2$ . The choices  $d_j = p$  or  $p^2$  are too big if  $1^2 + d_2^2 + \cdots + d_r^2 = p^2$ . □

# More silly applications

## Theorem

*If  $|G| = p^2$ ,  $p$  prime, then  $G$  is abelian.*

## Proof.

Each degree  $d_j$  must be 1,  $p$  or  $p^2$ . The choices  $d_j = p$  or  $p^2$  are too big if  $1^2 + d_2^2 + \cdots + d_r^2 = p^2$ . □

## Theorem

*If  $|G| = pq$ ,  $q < p$  primes, and  $G$  is nonabelian, then  $p \equiv 1 \pmod{q}$ .*



# More silly applications

## Theorem

*If  $|G| = p^2$ ,  $p$  prime, then  $G$  is abelian.*

## Proof.

Each degree  $d_j$  must be 1,  $p$  or  $p^2$ . The choices  $d_j = p$  or  $p^2$  are too big if  $1^2 + d_2^2 + \cdots + d_r^2 = p^2$ . □

## Theorem

*If  $|G| = pq$ ,  $q < p$  primes, and  $G$  is nonabelian, then  $p \equiv 1 \pmod{q}$ .*

## Proof.

# More silly applications

## Theorem

*If  $|G| = p^2$ ,  $p$  prime, then  $G$  is abelian.*

## Proof.

Each degree  $d_j$  must be 1,  $p$  or  $p^2$ . The choices  $d_j = p$  or  $p^2$  are too big if  $1^2 + d_2^2 + \cdots + d_r^2 = p^2$ . □

## Theorem

*If  $|G| = pq$ ,  $q < p$  primes, and  $G$  is nonabelian, then  $p \equiv 1 \pmod{q}$ .*

## Proof.

# More silly applications

## Theorem

*If  $|G| = p^2$ ,  $p$  prime, then  $G$  is abelian.*

## Proof.

Each degree  $d_j$  must be 1,  $p$  or  $p^2$ . The choices  $d_j = p$  or  $p^2$  are too big if  $1^2 + d_2^2 + \cdots + d_r^2 = p^2$ . □

## Theorem

*If  $|G| = pq$ ,  $q < p$  primes, and  $G$  is nonabelian, then  $p \equiv 1 \pmod{q}$ .*

## Proof.

Possible degrees are  $d_j \in \{1, q\}$ , and some  $d_j = q$ .

# More silly applications

## Theorem

*If  $|G| = p^2$ ,  $p$  prime, then  $G$  is abelian.*

## Proof.

Each degree  $d_j$  must be 1,  $p$  or  $p^2$ . The choices  $d_j = p$  or  $p^2$  are too big if  $1^2 + d_2^2 + \cdots + d_r^2 = p^2$ . □

## Theorem

*If  $|G| = pq$ ,  $q < p$  primes, and  $G$  is nonabelian, then  $p \equiv 1 \pmod{q}$ .*

## Proof.

Possible degrees are  $d_j \in \{1, q\}$ , and some  $d_j = q$ . Assume that there are  $m$  degrees satisfying  $d_j = q$ .

# More silly applications

## Theorem

*If  $|G| = p^2$ ,  $p$  prime, then  $G$  is abelian.*

## Proof.

Each degree  $d_j$  must be 1,  $p$  or  $p^2$ . The choices  $d_j = p$  or  $p^2$  are too big if  $1^2 + d_2^2 + \cdots + d_r^2 = p^2$ . □

## Theorem

*If  $|G| = pq$ ,  $q < p$  primes, and  $G$  is nonabelian, then  $p \equiv 1 \pmod{q}$ .*

## Proof.

Possible degrees are  $d_j \in \{1, q\}$ , and some  $d_j = q$ . Assume that there are  $m$  degrees satisfying  $d_j = q$ . The number of  $d_j$  equal to 1 is  $n := [G : G'] \in \{1, q, p\}$ .

# More silly applications

## Theorem

*If  $|G| = p^2$ ,  $p$  prime, then  $G$  is abelian.*

## Proof.

Each degree  $d_j$  must be 1,  $p$  or  $p^2$ . The choices  $d_j = p$  or  $p^2$  are too big if  $1^2 + d_2^2 + \cdots + d_r^2 = p^2$ . □

## Theorem

*If  $|G| = pq$ ,  $q < p$  primes, and  $G$  is nonabelian, then  $p \equiv 1 \pmod{q}$ .*

## Proof.

Possible degrees are  $d_j \in \{1, q\}$ , and some  $d_j = q$ . Assume that there are  $m$  degrees satisfying  $d_j = q$ . The number of  $d_j$  equal to 1 is  $n := [G : G'] \in \{1, q, p\}$ . We have  $pq = d_1^2 + \cdots + d_r^2 = mq^2 + n$ .

# More silly applications

## Theorem

*If  $|G| = p^2$ ,  $p$  prime, then  $G$  is abelian.*

## Proof.

Each degree  $d_j$  must be 1,  $p$  or  $p^2$ . The choices  $d_j = p$  or  $p^2$  are too big if  $1^2 + d_2^2 + \cdots + d_r^2 = p^2$ . □

## Theorem

*If  $|G| = pq$ ,  $q < p$  primes, and  $G$  is nonabelian, then  $p \equiv 1 \pmod{q}$ .*

## Proof.

Possible degrees are  $d_j \in \{1, q\}$ , and some  $d_j = q$ . Assume that there are  $m$  degrees satisfying  $d_j = q$ . The number of  $d_j$  equal to 1 is  $n := [G : G'] \in \{1, q, p\}$ . We have  $pq = d_1^2 + \cdots + d_r^2 = mq^2 + n$ . From this,  $q|n$ , so  $q = n$ .

# More silly applications

## Theorem

*If  $|G| = p^2$ ,  $p$  prime, then  $G$  is abelian.*

## Proof.

Each degree  $d_j$  must be 1,  $p$  or  $p^2$ . The choices  $d_j = p$  or  $p^2$  are too big if  $1^2 + d_2^2 + \cdots + d_r^2 = p^2$ . □

## Theorem

*If  $|G| = pq$ ,  $q < p$  primes, and  $G$  is nonabelian, then  $p \equiv 1 \pmod{q}$ .*

## Proof.

Possible degrees are  $d_j \in \{1, q\}$ , and some  $d_j = q$ . Assume that there are  $m$  degrees satisfying  $d_j = q$ . The number of  $d_j$  equal to 1 is  $n := [G : G'] \in \{1, q, p\}$ . We have  $pq = d_1^2 + \cdots + d_r^2 = mq^2 + n$ . From this,  $q|n$ , so  $q = n$ . Dividing  $pq = mq^2 + q$  by  $q$  we get  $p = mq + 1$ . □



## Theorem

*A finite nonabelian simple group  $G$  has no conjugacy class  $h^G$  of prime power size other than the class  $\{1\}$ .*

# Unsilly applications

## Theorem

*A finite nonabelian simple group  $G$  has no conjugacy class  $h^G$  of prime power size other than the class  $\{1\}$ .*

## Proof.

# Unsilly applications

## Theorem

*A finite nonabelian simple group  $G$  has no conjugacy class  $h^G$  of prime power size other than the class  $\{1\}$ .*

## Proof.

## Theorem

*A finite nonabelian simple group  $G$  has no conjugacy class  $h^G$  of prime power size other than the class  $\{1\}$ .*

## Proof.

If  $\chi \in \text{Irr}(G)$  is not the principal character, then  $K_\chi = Z_\chi = \{1\}$ , since these are proper normal subgroups.

# Unsilly applications

## Theorem

*A finite nonabelian simple group  $G$  has no conjugacy class  $h^G$  of prime power size other than the class  $\{1\}$ .*

## Proof.

If  $\chi \in \text{Irr}(G)$  is not the principal character, then  $K_\chi = Z_\chi = \{1\}$ , since these are proper normal subgroups. If  $p$  is the prime of the theorem, then  $p \nmid \chi(1)$  implies  $\chi(h) = 0$  for any such  $\chi$ .

# Unsilly applications

## Theorem

*A finite nonabelian simple group  $G$  has no conjugacy class  $h^G$  of prime power size other than the class  $\{1\}$ .*

## Proof.

If  $\chi \in \text{Irr}(G)$  is not the principal character, then  $K_\chi = Z_\chi = \{1\}$ , since these are proper normal subgroups. If  $p$  is the prime of the theorem, then  $p \nmid \chi(1)$  implies  $\chi(h) = 0$  for any such  $\chi$ . Column orthogonality, applied to the columns of  $1$  and  $h$ , yields  $1 + \chi_2(1)\chi_2(h) + \cdots + \chi_r(1)\chi_r(h) = 0$ ,

# Unsilly applications

## Theorem

*A finite nonabelian simple group  $G$  has no conjugacy class  $h^G$  of prime power size other than the class  $\{1\}$ .*

## Proof.

If  $\chi \in \text{Irr}(G)$  is not the principal character, then  $K_\chi = Z_\chi = \{1\}$ , since these are proper normal subgroups. If  $p$  is the prime of the theorem, then  $p \nmid \chi(1)$  implies  $\chi(h) = 0$  for any such  $\chi$ . Column orthogonality, applied to the columns of  $1$  and  $h$ , yields  $1 + \chi_2(1)\chi_2(h) + \cdots + \chi_r(1)\chi_r(h) = 0$ , hence

$$(\chi_2(1)\chi_2(h))/p + \cdots + (\chi_r(1)\chi_r(h))/p = -1/p.$$

# Unsilly applications

## Theorem

*A finite nonabelian simple group  $G$  has no conjugacy class  $h^G$  of prime power size other than the class  $\{1\}$ .*

## Proof.

If  $\chi \in \text{Irr}(G)$  is not the principal character, then  $K_\chi = Z_\chi = \{1\}$ , since these are proper normal subgroups. If  $p$  is the prime of the theorem, then  $p \nmid \chi(1)$  implies  $\chi(h) = 0$  for any such  $\chi$ . Column orthogonality, applied to the columns of  $1$  and  $h$ , yields  $1 + \chi_2(1)\chi_2(h) + \cdots + \chi_r(1)\chi_r(h) = 0$ , hence

$$(\chi_2(1)\chi_2(h))/p + \cdots + (\chi_r(1)\chi_r(h))/p = -1/p.$$

Each summand on the left is an algebraic integer, but  $-1/p$  is not. □



## Theorem

*If  $G$  is a finite nonabelian simple group and  $H$  is a subgroup of prime power index, then  $H$  is centerless.*

## Theorem

*If  $G$  is a finite nonabelian simple group and  $H$  is a subgroup of prime power index, then  $H$  is centerless.*

# Unsilly applications

## Theorem

*If  $G$  is a finite nonabelian simple group and  $H$  is a subgroup of prime power index, then  $H$  is centerless.*

(This is a restatement of the previous theorem in the language of subgroups instead of conjugacy classes.)

Let's see:

# Unsilly applications

## Theorem

*If  $G$  is a finite nonabelian simple group and  $H$  is a subgroup of prime power index, then  $H$  is centerless.*

(This is a restatement of the previous theorem in the language of subgroups instead of conjugacy classes.)

Let's see:

# Unsilly applications

## Theorem

*If  $G$  is a finite nonabelian simple group and  $H$  is a subgroup of prime power index, then  $H$  is centerless.*

(This is a restatement of the previous theorem in the language of subgroups instead of conjugacy classes.)

## Let's see:

*If  $H$  is a subgroup of  $G$  of  $p$ -power index with a nontrivial central element  $h$ , then  $h^G$  is a conjugacy class of  $G$  of  $p$ -power size.*

# Unsilly applications

## Theorem

*If  $G$  is a finite nonabelian simple group and  $H$  is a subgroup of prime power index, then  $H$  is centerless.*

(This is a restatement of the previous theorem in the language of subgroups instead of conjugacy classes.)

## Let's see:

*If  $H$  is a subgroup of  $G$  of  $p$ -power index with a nontrivial central element  $h$ , then  $h^G$  is a conjugacy class of  $G$  of  $p$ -power size. Conversely if  $h^G$  is a conjugacy class of  $G$  of  $p$ -power size, then  $C_G(h)$  is a subgroup of  $G$  of  $p$ -power index with a nontrivial central element  $h$ .*

# Burnside's $p^a q^b$ Theorem

## Theorem

*There is no nonabelian simple group  $G$  such that  $|G| = p^a q^b$ ,  $p, q$  primes.*

# Burnside's $p^a q^b$ Theorem

## Theorem

*There is no nonabelian simple group  $G$  such that  $|G| = p^a q^b$ ,  $p, q$  primes.*

## Proof.



# Burnside's $p^a q^b$ Theorem

## Theorem

*There is no nonabelian simple group  $G$  such that  $|G| = p^a q^b$ ,  $p, q$  primes.*

## Proof.

# Burnside's $p^a q^b$ Theorem

## Theorem

*There is no nonabelian simple group  $G$  such that  $|G| = p^a q^b$ ,  $p, q$  primes.*

## Proof.

Apply the preceding theorem to  $H \in \text{Syl}_p(G)$  to derive a contradiction. □

# Burnside's $p^a q^b$ Theorem

## Theorem

*There is no nonabelian simple group  $G$  such that  $|G| = p^a q^b$ ,  $p, q$  primes.*

## Proof.

Apply the preceding theorem to  $H \in \text{Syl}_p(G)$  to derive a contradiction. □

## Corollary

*If  $|G| = p^a q^b$ ,  $p, q$  primes, then  $G$  is solvable.*

# Burnside's $p^a q^b$ Theorem

## Theorem

*There is no nonabelian simple group  $G$  such that  $|G| = p^a q^b$ ,  $p, q$  primes.*

## Proof.

Apply the preceding theorem to  $H \in \text{Syl}_p(G)$  to derive a contradiction. □

## Corollary

*If  $|G| = p^a q^b$ ,  $p, q$  primes, then  $G$  is solvable.*

## Proof.

# Burnside's $p^a q^b$ Theorem

## Theorem

*There is no nonabelian simple group  $G$  such that  $|G| = p^a q^b$ ,  $p, q$  primes.*

## Proof.

Apply the preceding theorem to  $H \in \text{Syl}_p(G)$  to derive a contradiction. □

## Corollary

*If  $|G| = p^a q^b$ ,  $p, q$  primes, then  $G$  is solvable.*

## Proof.

# Burnside's $p^a q^b$ Theorem

## Theorem

*There is no nonabelian simple group  $G$  such that  $|G| = p^a q^b$ ,  $p, q$  primes.*

## Proof.

Apply the preceding theorem to  $H \in \text{Syl}_p(G)$  to derive a contradiction. □

## Corollary

*If  $|G| = p^a q^b$ ,  $p, q$  primes, then  $G$  is solvable.*

## Proof.

If  $|G| = p^a q^b$  and  $G$  is not solvable, then it must have a nonabelian simple section  $G'$  whose order divides  $p^a q^b$ .

# Burnside's $p^a q^b$ Theorem

## Theorem

*There is no nonabelian simple group  $G$  such that  $|G| = p^a q^b$ ,  $p, q$  primes.*

## Proof.

Apply the preceding theorem to  $H \in \text{Syl}_p(G)$  to derive a contradiction. ☐

## Corollary

*If  $|G| = p^a q^b$ ,  $p, q$  primes, then  $G$  is solvable.*

## Proof.

If  $|G| = p^a q^b$  and  $G$  is not solvable, then it must have a nonabelian simple section  $G'$  whose order divides  $p^a q^b$ . There is no such group. ☐

# A theorem about simple groups



# A theorem about simple groups

## Theorem

*If  $G$  is a nonabelian simple group and some irreducible character degree is a prime  $p$ , then the Sylow  $p$ -subgroups of  $G$  have order  $p$ .*

# A theorem about simple groups

## Theorem

*If  $G$  is a nonabelian simple group and some irreducible character degree is a prime  $p$ , then the Sylow  $p$ -subgroups of  $G$  have order  $p$ .*

# A theorem about simple groups

## Theorem

*If  $G$  is a nonabelian simple group and some irreducible character degree is a prime  $p$ , then the Sylow  $p$ -subgroups of  $G$  have order  $p$ .*

Recall, as an example:

# A theorem about simple groups

## Theorem

*If  $G$  is a nonabelian simple group and some irreducible character degree is a prime  $p$ , then the Sylow  $p$ -subgroups of  $G$  have order  $p$ .*

Recall, as an example:

	1	15	20	12	12
$A_5$	1	$(1\ 2)(3\ 4)$	$(1\ 2\ 3)$	$(1\ 2\ 3\ 4\ 5)$	$(1\ 2\ 3\ 5\ 4)$
$\chi_1$	1	1	1	1	1
$\chi_2$	3	-1	0	$\phi$	$\phi^{-1}$
$\chi_3$	3	-1	0	$\phi^{-1}$	$\phi$
$\chi_4$	4	0	1	-1	-1
$\chi_5$	5	1	-1	0	0

# A theorem about simple groups

## Theorem

*If  $G$  is a nonabelian simple group and some irreducible character degree is a prime  $p$ , then the Sylow  $p$ -subgroups of  $G$  have order  $p$ .*

Recall, as an example:

$A_5$	1	15	20	12	12
	1	(1 2)(3 4)	(1 2 3)	(1 2 3 4 5)	(1 2 3 5 4)
$\chi_1$	1	1	1	1	1
$\chi_2$	3	-1	0	$\phi$	$\phi^{-1}$
$\chi_3$	3	-1	0	$\phi^{-1}$	$\phi$
$\chi_4$	4	0	1	-1	-1
$\chi_5$	5	1	-1	0	0

# A theorem about simple groups

## Theorem

*If  $G$  is a nonabelian simple group and some irreducible character degree is a prime  $p$ , then the Sylow  $p$ -subgroups of  $G$  have order  $p$ .*

Recall, as an example:

$A_5$	1	15	20	12	12
	1	(1 2)(3 4)	(1 2 3)	(1 2 3 4 5)	(1 2 3 5 4)
$\chi_1$	1	1	1	1	1
$\chi_2$	3	-1	0	$\phi$	$\phi^{-1}$
$\chi_3$	3	-1	0	$\phi^{-1}$	$\phi$
$\chi_4$	4	0	1	-1	-1
$\chi_5$	5	1	-1	0	0

# A theorem about simple groups

## Theorem

*If  $G$  is a nonabelian simple group and some irreducible character degree is a prime  $p$ , then the Sylow  $p$ -subgroups of  $G$  have order  $p$ .*

Recall, as an example:

$A_5$	1	15	20	12	12
	1	(1 2)(3 4)	(1 2 3)	(1 2 3 4 5)	(1 2 3 5 4)
$\chi_1$	1	1	1	1	1
$\chi_2$	3	-1	0	$\phi$	$\phi^{-1}$
$\chi_3$	3	-1	0	$\phi^{-1}$	$\phi$
$\chi_4$	4	0	1	-1	-1
$\chi_5$	5	1	-1	0	0

# A theorem about simple groups

## Theorem

*If  $G$  is a nonabelian simple group and some irreducible character degree is a prime  $p$ , then the Sylow  $p$ -subgroups of  $G$  have order  $p$ .*

Recall, as an example:

$A_5$	1	15	20	12	12
	1	(1 2)(3 4)	(1 2 3)	(1 2 3 4 5)	(1 2 3 5 4)
$\chi_1$	1	1	1	1	1
$\chi_2$	3	-1	0	$\phi$	$\phi^{-1}$
$\chi_3$	3	-1	0	$\phi^{-1}$	$\phi$
$\chi_4$	4	0	1	-1	-1
$\chi_5$	5	1	-1	0	0

There are zeros at the centers of the colored crosses.





Let  $G$  be a nonabelian simple group with irreducible character  $\psi$  of degree  $p$ .

Let  $G$  be a nonabelian simple group with irreducible character  $\psi$  of degree  $p$ . If  $\rho: G \rightarrow \mathrm{GL}_p(\mathbb{C})$  affords  $\psi$ , then  $\rho$  is faithful.

Let  $G$  be a nonabelian simple group with irreducible character  $\psi$  of degree  $p$ . If  $\rho: G \rightarrow \mathrm{GL}_p(\mathbb{C})$  affords  $\psi$ , then  $\rho$  is faithful. In fact,  $K_\psi = Z_\psi = \{1\}$ .

Let  $G$  be a nonabelian simple group with irreducible character  $\psi$  of degree  $p$ . If  $\rho: G \rightarrow \mathrm{GL}_p(\mathbb{C})$  affords  $\psi$ , then  $\rho$  is faithful. In fact,  $K_\psi = Z_\psi = \{1\}$ .

Let  $P$  be a Sylow  $p$ -subgroup of  $G$ .

Let  $G$  be a nonabelian simple group with irreducible character  $\psi$  of degree  $p$ . If  $\rho: G \rightarrow \mathrm{GL}_p(\mathbb{C})$  affords  $\psi$ , then  $\rho$  is faithful. In fact,  $K_\psi = Z_\psi = \{1\}$ .

Let  $P$  be a Sylow  $p$ -subgroup of  $G$ . Choose any  $z \in Z(P) \setminus \{1\}$ .

Let  $G$  be a nonabelian simple group with irreducible character  $\psi$  of degree  $p$ . If  $\rho: G \rightarrow \mathrm{GL}_p(\mathbb{C})$  affords  $\psi$ , then  $\rho$  is faithful. In fact,  $K_\psi = Z_\psi = \{1\}$ .

Let  $P$  be a Sylow  $p$ -subgroup of  $G$ . Choose any  $z \in Z(P) \setminus \{1\}$ .  $[G : C_G(z)]$  divides  $[G : P]$ , hence  $z^G$  has size prime to  $p$ .

# Preliminaries

Let  $G$  be a nonabelian simple group with irreducible character  $\psi$  of degree  $p$ . If  $\rho: G \rightarrow \mathrm{GL}_p(\mathbb{C})$  affords  $\psi$ , then  $\rho$  is faithful. In fact,  $K_\psi = Z_\psi = \{1\}$ .

Let  $P$  be a Sylow  $p$ -subgroup of  $G$ . Choose any  $z \in Z(P) \setminus \{1\}$ .  $[G : C_G(z)]$  divides  $[G : P]$ , hence  $z^G$  has size prime to  $p$ . Thus  $\psi(z) = 0$  for any  $z \in Z(P) \setminus \{1\}$ .



# Preliminaries

Let  $G$  be a nonabelian simple group with irreducible character  $\psi$  of degree  $p$ . If  $\rho: G \rightarrow \mathrm{GL}_p(\mathbb{C})$  affords  $\psi$ , then  $\rho$  is faithful. In fact,  $K_\psi = Z_\psi = \{1\}$ .

Let  $P$  be a Sylow  $p$ -subgroup of  $G$ . Choose any  $z \in Z(P) \setminus \{1\}$ .  $[G : C_G(z)]$  divides  $[G : P]$ , hence  $z^G$  has size prime to  $p$ . Thus  $\psi(z) = 0$  for any  $z \in Z(P) \setminus \{1\}$ .

What we are seeing here is important:

# Preliminaries

Let  $G$  be a nonabelian simple group with irreducible character  $\psi$  of degree  $p$ . If  $\rho: G \rightarrow \mathrm{GL}_p(\mathbb{C})$  affords  $\psi$ , then  $\rho$  is faithful. In fact,  $K_\psi = Z_\psi = \{1\}$ .

Let  $P$  be a Sylow  $p$ -subgroup of  $G$ . Choose any  $z \in Z(P) \setminus \{1\}$ .  $[G : C_G(z)]$  divides  $[G : P]$ , hence  $z^G$  has size prime to  $p$ . Thus  $\psi(z) = 0$  for any  $z \in Z(P) \setminus \{1\}$ .

What we are seeing here is important: a character is vanishing on the nonidentity elements of a subgroup,  $Z(P)$ .

# Recognizing multiples of the regular representation

# Recognizing multiples of the regular representation

## Lemma

*Let  $\rho: H \rightarrow GL_n(\mathbb{C})$  be a (not necessarily irreducible) representation of a finite group, and let  $\psi$  be the character afforded by  $\rho$ . If  $\psi$  vanishes on  $H \setminus \{1\}$ , then  $|H|$  divides  $\psi(1)$ .*

# Recognizing multiples of the regular representation

## Lemma

*Let  $\rho: H \rightarrow GL_n(\mathbb{C})$  be a (not necessarily irreducible) representation of a finite group, and let  $\psi$  be the character afforded by  $\rho$ . If  $\psi$  vanishes on  $H \setminus \{1\}$ , then  $|H|$  divides  $\psi(1)$ .*

# Recognizing multiples of the regular representation

## Lemma

*Let  $\rho: H \rightarrow GL_n(\mathbb{C})$  be a (not necessarily irreducible) representation of a finite group, and let  $\psi$  be the character afforded by  $\rho$ . If  $\psi$  vanishes on  $H \setminus \{1\}$ , then  $|H|$  divides  $\psi(1)$ .*

*Thus,  $\psi$  is a multiple of the regular character,*

# Recognizing multiples of the regular representation

## Lemma

*Let  $\rho: H \rightarrow GL_n(\mathbb{C})$  be a (not necessarily irreducible) representation of a finite group, and let  $\psi$  be the character afforded by  $\rho$ . If  $\psi$  vanishes on  $H \setminus \{1\}$ , then  $|H|$  divides  $\psi(1)$ .*

*Thus,  $\psi$  is a multiple of the regular character, and so  $\rho$  is a multiple of the regular representation.*

# Recognizing multiples of the regular representation

## Lemma

*Let  $\rho: H \rightarrow GL_n(\mathbb{C})$  be a (not necessarily irreducible) representation of a finite group, and let  $\psi$  be the character afforded by  $\rho$ . If  $\psi$  vanishes on  $H \setminus \{1\}$ , then  $|H|$  divides  $\psi(1)$ .*

*Thus,  $\psi$  is a multiple of the regular character, and so  $\rho$  is a multiple of the regular representation.*

## Proof.



# Recognizing multiples of the regular representation

## Lemma

*Let  $\rho: H \rightarrow GL_n(\mathbb{C})$  be a (not necessarily irreducible) representation of a finite group, and let  $\psi$  be the character afforded by  $\rho$ . If  $\psi$  vanishes on  $H \setminus \{1\}$ , then  $|H|$  divides  $\psi(1)$ .*

*Thus,  $\psi$  is a multiple of the regular character, and so  $\rho$  is a multiple of the regular representation.*

## Proof.

# Recognizing multiples of the regular representation

## Lemma

Let  $\rho: H \rightarrow GL_n(\mathbb{C})$  be a (not necessarily irreducible) representation of a finite group, and let  $\psi$  be the character afforded by  $\rho$ . If  $\psi$  vanishes on  $H \setminus \{1\}$ , then  $|H|$  divides  $\psi(1)$ .

*Thus,  $\psi$  is a multiple of the regular character, and so  $\rho$  is a multiple of the regular representation.*

## Proof.

Compare  $\chi_1$  and  $\psi$ :

$H$	1	$k_2$	$\cdots$	$k_r$
	1	$a_2$	$\cdots$	$a_r$
$\chi_1$	1	1	$\cdots$	1
$\psi$	$\psi(1)$	0	$\cdots$	0

# Recognizing multiples of the regular representation

## Lemma

Let  $\rho: H \rightarrow GL_n(\mathbb{C})$  be a (not necessarily irreducible) representation of a finite group, and let  $\psi$  be the character afforded by  $\rho$ . If  $\psi$  vanishes on  $H \setminus \{1\}$ , then  $|H|$  divides  $\psi(1)$ .

*Thus,  $\psi$  is a multiple of the regular character, and so  $\rho$  is a multiple of the regular representation.*

## Proof.

Compare  $\chi_1$  and  $\psi$ :

	1	$k_2$	$\cdots$	$k_r$
$H$	1	$a_2$	$\cdots$	$a_r$
$\chi_1$	1	1	$\cdots$	1
$\psi$	$\psi(1)$	0	$\cdots$	0

$\langle \chi_1, \psi \rangle = \frac{\psi(1)}{|H|}$  must be an integer.

# Recognizing multiples of the regular representation

## Lemma

Let  $\rho: H \rightarrow GL_n(\mathbb{C})$  be a (not necessarily irreducible) representation of a finite group, and let  $\psi$  be the character afforded by  $\rho$ . If  $\psi$  vanishes on  $H \setminus \{1\}$ , then  $|H|$  divides  $\psi(1)$ .

*Thus,  $\psi$  is a multiple of the regular character, and so  $\rho$  is a multiple of the regular representation.*

## Proof.

Compare  $\chi_1$  and  $\psi$ :

	1	$k_2$	$\cdots$	$k_r$
$H$	1	$a_2$	$\cdots$	$a_r$
$\chi_1$	1	1	$\cdots$	1
$\psi$	$\psi(1)$	0	$\cdots$	0

$\langle \chi_1, \psi \rangle = \frac{\psi(1)}{|H|}$  must be an integer. Now use that representations are determined by their characters.



# Proof of theorem

# Proof of theorem

To be proved: If there is an irreducible representation  $\rho: G \rightarrow \mathrm{GL}_p(\mathbb{C})$  (affording  $\psi$ ) then a Sylow  $p$ -subgroup  $P \leq G$  has order  $p$ .

# Proof of theorem

To be proved: If there is an irreducible representation  $\rho: G \rightarrow \mathrm{GL}_p(\mathbb{C})$  (affording  $\psi$ ) then a Sylow  $p$ -subgroup  $P \leq G$  has order  $p$ .

Proof.

We have shown that  $\psi$  vanishes on the nonidentity elements of  $Z(P)$ , so  $|Z(P)| = p$ . Moreover,  $\rho_{Z(P)}$  is the regular representation of  $Z(P)$ ,

# Proof of theorem

To be proved: If there is an irreducible representation  $\rho: G \rightarrow \mathrm{GL}_p(\mathbb{C})$  (affording  $\psi$ ) then a Sylow  $p$ -subgroup  $P \leq G$  has order  $p$ .

Proof.

We have shown that  $\psi$  vanishes on the nonidentity elements of  $Z(P)$ , so  $|Z(P)| = p$ . Moreover,  $\rho_{Z(P)}$  is the regular representation of  $Z(P)$ ,



# Proof of theorem

To be proved: If there is an irreducible representation  $\rho: G \rightarrow \mathrm{GL}_p(\mathbb{C})$  (affording  $\psi$ ) then a Sylow  $p$ -subgroup  $P \leq G$  has order  $p$ .

Proof.

We have shown that  $\psi$  vanishes on the nonidentity elements of  $Z(P)$ , so  $|Z(P)| = p$ . Moreover,  $\rho_{Z(P)}$  is the regular representation of  $Z(P)$ , so if  $z \in Z(P)$  is a generator, then  $\rho(z)$  has distinct e-values.

# Proof of theorem

To be proved: If there is an irreducible representation  $\rho: G \rightarrow \mathrm{GL}_p(\mathbb{C})$  (affording  $\psi$ ) then a Sylow  $p$ -subgroup  $P \leq G$  has order  $p$ .

Proof.

We have shown that  $\psi$  vanishes on the nonidentity elements of  $Z(P)$ , so  $|Z(P)| = p$ . Moreover,  $\rho_{Z(P)}$  is the regular representation of  $Z(P)$ , so if  $z \in Z(P)$  is a generator, then  $\rho(z)$  has distinct e-values. It follows that  $C_{\mathrm{GL}_p(\mathbb{C})}(z)$  consists of diagonal matrices, hence is abelian.

# Proof of theorem

To be proved: If there is an irreducible representation  $\rho: G \rightarrow \mathrm{GL}_p(\mathbb{C})$  (affording  $\psi$ ) then a Sylow  $p$ -subgroup  $P \leq G$  has order  $p$ .

Proof.

We have shown that  $\psi$  vanishes on the nonidentity elements of  $Z(P)$ , so  $|Z(P)| = p$ . Moreover,  $\rho_{Z(P)}$  is the regular representation of  $Z(P)$ , so if  $z \in Z(P)$  is a generator, then  $\rho(z)$  has distinct  $e$ -values. It follows that  $C_{\mathrm{GL}_p(\mathbb{C})}(z)$  consists of diagonal matrices, hence is abelian. A fortiori,  $C_G(z)$  is abelian.

# Proof of theorem

To be proved: If there is an irreducible representation  $\rho: G \rightarrow \mathrm{GL}_p(\mathbb{C})$  (affording  $\psi$ ) then a Sylow  $p$ -subgroup  $P \leq G$  has order  $p$ .

## Proof.

We have shown that  $\psi$  vanishes on the nonidentity elements of  $Z(P)$ , so  $|Z(P)| = p$ . Moreover,  $\rho_{Z(P)}$  is the regular representation of  $Z(P)$ , so if  $z \in Z(P)$  is a generator, then  $\rho(z)$  has distinct  $e$ -values. It follows that  $C_{\mathrm{GL}_p(\mathbb{C})}(z)$  consists of diagonal matrices, hence is abelian. A fortiori,  $C_G(z)$  is abelian. Thus,  $P = Z(P)$ , which has order  $p$ . □

# Proof of theorem

To be proved: If there is an irreducible representation  $\rho: G \rightarrow \mathrm{GL}_p(\mathbb{C})$  (affording  $\psi$ ) then a Sylow  $p$ -subgroup  $P \leq G$  has order  $p$ .

## Proof.

We have shown that  $\psi$  vanishes on the nonidentity elements of  $Z(P)$ , so  $|Z(P)| = p$ . Moreover,  $\rho_{Z(P)}$  is the regular representation of  $Z(P)$ , so if  $z \in Z(P)$  is a generator, then  $\rho(z)$  has distinct e-values. It follows that  $C_{\mathrm{GL}_p(\mathbb{C})}(z)$  consists of diagonal matrices, hence is abelian. A fortiori,  $C_G(z)$  is abelian. Thus,  $P = Z(P)$ , which has order  $p$ . □

The converse is false.

# Proof of theorem

To be proved: If there is an irreducible representation  $\rho: G \rightarrow \mathrm{GL}_p(\mathbb{C})$  (affording  $\psi$ ) then a Sylow  $p$ -subgroup  $P \leq G$  has order  $p$ .

## Proof.

We have shown that  $\psi$  vanishes on the nonidentity elements of  $Z(P)$ , so  $|Z(P)| = p$ . Moreover,  $\rho_{Z(P)}$  is the regular representation of  $Z(P)$ , so if  $z \in Z(P)$  is a generator, then  $\rho(z)$  has distinct e-values. It follows that  $C_{\mathrm{GL}_p(\mathbb{C})}(z)$  consists of diagonal matrices, hence is abelian. A fortiori,  $C_G(z)$  is abelian. Thus,  $P = Z(P)$ , which has order  $p$ . □

The converse is false.  $\mathrm{PSL}_2(q)$  has Sylow 3-subgroups of size 3 for infinitely many prime powers  $q$ , but only  $\mathrm{PSL}_2(5)$  and  $\mathrm{PSL}_2(7)$  have irreps of degree 3.