

Solutions to HW 5.

1. (Exercise 2.7.14(a).) Point out how the hypothesis of Dirichlet's Test differs from that of Abel's Test in Exercise 2.7.13, but show that essentially the same strategy can be used to provide a proof.

The differences: Dirichlet's Test assumes that $\lim y_k = 0$, while Abel's Test does not. Abel's Test assumes that $\sum x_k$ converges, while Dirichlet's Test assumes only that the sequence of partial sums is bounded.

To prove Dirichlet's Test, we will use the summation by parts formula:

$$(2.7.12) \quad \sum_{k=1}^n x_k y_k = s_n y_{n+1} + \sum_{k=1}^n s_k (y_k - y_{k+1}).$$

Here $s_k = \sum_{n=1}^k x_n$ is the k th partial sum of the series $\sum_{n=1}^{\infty} x_n$. To show that the left hand side of (2.7.12) has a limit, it suffices to show that the right hand side has a limit. For this, it will suffice to show that the series on the right hand side is absolutely convergent. Let M be a bound such that $|s_k| \leq M$ for all k . Now

$$\sum_{k=1}^n |s_k (y_k - y_{k+1})| \leq \sum_{k=1}^n M (y_k - y_{k+1}) = M y_1 - M y_{n+1} \leq M y_1.$$

Since $\sum_{k=1}^{\infty} |s_k (y_k - y_{k+1})|$ has bounded partial sums, $\sum_{k=1}^{\infty} s_k (y_k - y_{k+1})$ is absolutely convergent, hence $\sum_{k=1}^{\infty} x_k y_k$ is convergent.

2. (a) Let θ be an arbitrary angle. Show that the partial sums of $\sum_{k=1}^{\infty} \sin(k\theta)$ are bounded. Hint: You may want to first prove and use the equality

$$2 \sin(k\theta) \sin(\theta/2) = \cos((k - 1/2)\theta) - \cos((k + 1/2)\theta).$$

The angle addition formula implies that

$$\begin{aligned} \cos((k - 1/2)\theta) &= \cos(k\theta) \cos(\theta/2) + \sin(k\theta) \sin(\theta/2) \\ \cos((k + 1/2)\theta) &= \cos(k\theta) \cos(\theta/2) - \sin(k\theta) \sin(\theta/2). \end{aligned}$$

Subtracting yields

$$\cos((k - 1/2)\theta) - \cos((k + 1/2)\theta) = 2 \sin(k\theta) \sin(\theta/2).$$

If $\sin(\theta/2) \neq 0$, then we can divide to obtain that

$$\sin(k\theta) = \frac{\cos((k - 1/2)\theta) - \cos((k + 1/2)\theta)}{2 \sin(\theta/2)}.$$

This means that the n th partial sum of $\sum \sin(k\theta)$ is

$$\begin{aligned} \sum_{k=1}^n \sin(k\theta) &= \sin(\theta) + \sin(2\theta) + \cdots + \sin(n\theta) \\ &= \frac{\cos((1/2)\theta) - \cos((3/2)\theta)}{2 \sin(\theta/2)} + \frac{\cos((3/2)\theta) - \cos((5/2)\theta)}{2 \sin(\theta/2)} + \cdots + \frac{\cos((n-1/2)\theta) - \cos((n+1/2)\theta)}{2 \sin(\theta/2)} \\ &= \frac{\cos((1/2)\theta) - \cos((n+1/2)\theta)}{2 \sin(\theta/2)} \\ &\leq \frac{2}{2 \sin(\theta/2)} = \frac{1}{\sin(\theta/2)}. \end{aligned}$$

This upper bound on partial sums is independent of n , so the partial sums are bounded.

The preceding argument required the assumption that $\sin(\theta/2) \neq 0$. If instead $\sin(\theta/2) = 0$, then $\sin(k\theta) = 0$ for all k . In this case the partial sums of $\sum \sin(k\theta)$ are still bounded, since they are all zero.

- (b) Show that $\sum_{k=1}^{\infty} \sin(k\theta)/k$ converges.

Here we just cite Dirichlet's Test with $y_k = 1/k$ and $x_k = \sin(k\theta)$.

3. (Exercise 3.2.3.) Decide whether the following sets are open, closed, or neither. If a set is not open, find a point in the set for which there is no ε -neighborhood contained in the set. If a set is not closed, find a limit point that is not contained in the set.

- (a) \mathbb{Q}

Neither open nor closed. $1 \in \mathbb{Q}$ is not an interior point, while $\sqrt{2} \notin \mathbb{Q}$ is a limit point.

- (b) \mathbb{N}

Closed. $1 \in \mathbb{N}$ is not an interior point.

- (c) $X := \{x \in \mathbb{R} \mid x \neq 0\}$.

Open. $0 \notin X$ is a limit point.

- (d) $Y := \{1 + 1/4 + \cdots + 1/n^2 \mid n \in \mathbb{N}^*\}$

Neither open nor closed. $1 \in Y$ is not an interior point, while $\sum_{k=1}^{\infty} 1/k^2 \notin Y$ is a limit point. The reason this value is not in Y is that the partial sums of $\sum_{k=1}^{\infty} 1/k^2$ are positive and strictly increasing, so the limit is greater than any partial sum.

- (e) $Z := \{1 + 1/2 + \cdots + 1/n \mid n \in \mathbb{N}^*\}$

Closed. $1 \in Z$ is not an interior point.