

Solutions to HW 4.

1. (Exercise 2.2.4.) Give an example of each or state that the request is impossible. For any that are impossible, give a compelling argument for why that is the case.

- (a) A sequence with an infinite number of ones that does not converge to one.

$$(0, 1, 0, 1, 0, 1, \dots)$$

- (b) A sequence with an infinite number of ones that converges to a limit not equal to one.

Such a sequence cannot exist.

Assume that $(a_i)_{i \in \mathbb{N}^*}$ has infinitely many 1's, and converges to $L \neq 1$. Let $\varepsilon = |1 - L|/2$. Choose N so that whenever $i > N$ it is the case that $|a_i - L| < \varepsilon$. Choose $i > N$ so that $a_i = 1$. Now $|a_i - L| = |1 - L| < \varepsilon = |1 - L|/2$. But $|1 - L| < |1 - L|/2$ leads to $|1 - L| < 0$, which is impossible.

- (c) A divergent sequence such that for every $n \in \mathbb{N}$ it is possible to find n consecutive ones somewhere in the sequence.

$$(a_1, a_2, a_3, a_4, a_5, \dots) = (0, 0, 1, 0, 1, 1, 1, 0, 1, 1, 1, 1, 1, 1, 1, 0, 1, \dots)$$

Here $a_n = 0$ if n is a power of 2, and $a_n = 1$ otherwise.

2. (Exercise 2.3.7 (a), (b), (c).) Give an example of each of the following, or state that such a request is impossible by referencing the proper theorem(s):

- (a) sequences (x_n) and (y_n) , which both diverge, but whose sum $(x_n + y_n)$ converges.

Let $x_n = n$ and $y_n = -n$. Then

$$(x_1) = (1, 2, 3, \dots) \quad \text{and} \quad (y_n) = (-1, -2, -3, \dots)$$

both diverge, but their sum $(x_n + y_n) = (0, 0, 0, \dots)$ converges.

- (b) sequences (x_n) and (y_n) , where (x_n) converges, (y_n) diverges, and $(x_n + y_n)$ converges.

Such sequences cannot exist.

Assume that (x_n) and $(x_n + y_n)$ converge. If $a_n = -x_n$ and $b_n = x_n + y_n$, then both (a_n) and (b_n) converge, by assumption and by the Algebraic Limit Theorem. Hence $(a_n + b_n) = (y_n)$ converges, by the Algebraic Limit Theorem.

- (c) a convergent sequence (b_n) with $b_n \neq 0$ for all n such that $(1/b_n)$ diverges.

$$(b_n) = \left(1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots\right)$$

3. (Exercise 2.4.4(a).) In Section 1.4 we used the Axiom of Completeness (AoC) to prove the Archimedean Property of \mathbb{R} (Theorem 1.4.2). Show that the Monotone Convergence Theorem can also be used to prove the Archimedean Property without making any use of AoC.

Proof. This is a proof by contradiction, so assume that \mathbb{R} is not Archimedean.

Since \mathbb{R} is not Archimedean, \mathbb{N} is bounded above, which implies that $(a_n) := (1, 2, 3, \dots)$ is a monotone increasing bounded sequence. By the Monotone Convergence Theorem, this sequence has a limit, $L \in \mathbb{R}$. There must exist some N such that for $i > N$ we have $a_i \in (L - 1, L + 1)$, or equivalently $L - 1 < a_i < L + 1$. In particular, this means that

$$L - 1 < a_{N+1} < a_{N+2} < a_{N+3} < \dots < L + 1.$$

But if $L - 1 < a_{N+1} < L + 1$, then (by adding 2 throughout) we have $L + 1 < a_{N+3} < L + 3$, leading to the contradiction that $a_{N+3} < L + 1 < a_{N+3}$.