


Homology and Cohomology



Enrico Betti



Why?

Why?

- ① Poincare introduced the idea of homology based on the concept of Betti numbers.

Why?

- ① Poincare introduced the idea of homology based on the concept of Betti numbers.

Why?

- ① Poincare introduced the idea of homology based on the concept of Betti numbers. ($b_d(X)$ = number of “ d -dimensional holes” of space X)

Why?

- ① Poincare introduced the idea of homology based on the concept of Betti numbers. ($b_d(X)$ = number of “ d -dimensional holes” of space X)
- ② Modern formulation is due to Emmy Noether, and focuses on homology and cohomology groups.

Why?

- ① Poincare introduced the idea of homology based on the concept of Betti numbers. ($b_d(X)$ = number of “ d -dimensional holes” of space X)
- ② Modern formulation is due to Emmy Noether, and focuses on homology and cohomology groups.

Why?

- ① Poincare introduced the idea of homology based on the concept of Betti numbers. ($b_d(X)$ = number of “ d -dimensional holes” of space X)
- ② Modern formulation is due to Emmy Noether, and focuses on homology and cohomology groups.
- ③ Homology is typically defined via the Tor functor, which is derived from the tensor functor.

Why?

- ① Poincare introduced the idea of homology based on the concept of Betti numbers. ($b_d(X)$ = number of “ d -dimensional holes” of space X)
- ② Modern formulation is due to Emmy Noether, and focuses on homology and cohomology groups.
- ③ Homology is typically defined via the Tor functor, which is derived from the tensor functor.

Why?

- ① Poincare introduced the idea of homology based on the concept of Betti numbers. ($b_d(X)$ = number of “ d -dimensional holes” of space X)
- ② Modern formulation is due to Emmy Noether, and focuses on homology and cohomology groups.
- ③ Homology is typically defined via the Tor functor, which is derived from the tensor functor.
- ④ Cohomology is typically defined via the Ext functor, which is derived from the Hom functor.

Why?

- ① Poincare introduced the idea of homology based on the concept of Betti numbers. ($b_d(X)$ = number of “ d -dimensional holes” of space X)
- ② Modern formulation is due to Emmy Noether, and focuses on homology and cohomology groups.
- ③ Homology is typically defined via the Tor functor, which is derived from the tensor functor.
- ④ Cohomology is typically defined via the Ext functor, which is derived from the Hom functor.

Why?

- 1 Poincare introduced the idea of homology based on the concept of Betti numbers. ($b_d(X)$ = number of “ d -dimensional holes” of space X)
- 2 Modern formulation is due to Emmy Noether, and focuses on homology and cohomology groups.
- 3 Homology is typically defined via the Tor functor, which is derived from the tensor functor.
- 4 Cohomology is typically defined via the Ext functor, which is derived from the Hom functor.
- 5 The machinery for computing these groups works outside of the topological setting.

Why?

- 1 Poincare introduced the idea of homology based on the concept of Betti numbers. ($b_d(X)$ = number of “ d -dimensional holes” of space X)
- 2 Modern formulation is due to Emmy Noether, and focuses on homology and cohomology groups.
- 3 Homology is typically defined via the Tor functor, which is derived from the tensor functor.
- 4 Cohomology is typically defined via the Ext functor, which is derived from the Hom functor.
- 5 The machinery for computing these groups works outside of the topological setting.

Why?

- 1 Poincare introduced the idea of homology based on the concept of Betti numbers. ($b_d(X)$ = number of “ d -dimensional holes” of space X)
- 2 Modern formulation is due to Emmy Noether, and focuses on homology and cohomology groups.
- 3 Homology is typically defined via the Tor functor, which is derived from the tensor functor.
- 4 Cohomology is typically defined via the Ext functor, which is derived from the Hom functor.
- 5 The machinery for computing these groups works outside of the topological setting.
- 6 Algebraically, the definition of these groups are chosen so that they “repair the loss of exactness” for Hom and tensor.

Why?

- 1 Poincare introduced the idea of homology based on the concept of Betti numbers. ($b_d(X)$ = number of “ d -dimensional holes” of space X)
- 2 Modern formulation is due to Emmy Noether, and focuses on homology and cohomology groups.
- 3 Homology is typically defined via the Tor functor, which is derived from the tensor functor.
- 4 Cohomology is typically defined via the Ext functor, which is derived from the Hom functor.
- 5 The machinery for computing these groups works outside of the topological setting.
- 6 Algebraically, the definition of these groups are chosen so that they “repair the loss of exactness” for Hom and tensor.

Why?

- 1 Poincare introduced the idea of homology based on the concept of Betti numbers. ($b_d(X)$ = number of “ d -dimensional holes” of space X)
- 2 Modern formulation is due to Emmy Noether, and focuses on homology and cohomology groups.
- 3 Homology is typically defined via the Tor functor, which is derived from the tensor functor.
- 4 Cohomology is typically defined via the Ext functor, which is derived from the Hom functor.
- 5 The machinery for computing these groups works outside of the topological setting.
- 6 Algebraically, the definition of these groups are chosen so that they “repair the loss of exactness” for Hom and tensor.
- 7 Nonvanishing of these groups in given dimension can be interpreted as “ d -dimensional obstructions to injectivity/projectivity/flatness”.

Why?

- 1 Poincare introduced the idea of homology based on the concept of Betti numbers. ($b_d(X)$ = number of “ d -dimensional holes” of space X)
- 2 Modern formulation is due to Emmy Noether, and focuses on homology and cohomology groups.
- 3 Homology is typically defined via the Tor functor, which is derived from the tensor functor.
- 4 Cohomology is typically defined via the Ext functor, which is derived from the Hom functor.
- 5 The machinery for computing these groups works outside of the topological setting.
- 6 Algebraically, the definition of these groups are chosen so that they “repair the loss of exactness” for Hom and tensor.
- 7 Nonvanishing of these groups in given dimension can be interpreted as “ d -dimensional obstructions to injectivity/projectivity/flatness”.

Why?

- 1 Poincare introduced the idea of homology based on the concept of Betti numbers. ($b_d(X)$ = number of “ d -dimensional holes” of space X)
- 2 Modern formulation is due to Emmy Noether, and focuses on homology and cohomology groups.
- 3 Homology is typically defined via the Tor functor, which is derived from the tensor functor.
- 4 Cohomology is typically defined via the Ext functor, which is derived from the Hom functor.
- 5 The machinery for computing these groups works outside of the topological setting.
- 6 Algebraically, the definition of these groups are chosen so that they “repair the loss of exactness” for Hom and tensor.
- 7 Nonvanishing of these groups in given dimension can be interpreted as “ d -dimensional obstructions to injectivity/projectivity/flatness”.
- 8 These groups are invariants, which can be used for making distinctions.

The idea behind the definitions

[Whiteboard!]

Computing $\mathrm{Tor}_n^R(A, B)$

Computing $\mathrm{Tor}_n^R(A, B)$

Start with a **projective resolution** of B :

$$\cdots \xrightarrow{\delta_3} P_2 \xrightarrow{\delta_2} P_1 \xrightarrow{\delta_1} P_0 \xrightarrow{\delta_0} B \rightarrow 0.$$

Computing $\text{Tor}_n^R(A, B)$

Start with a **projective resolution** of B :

$$\cdots \xrightarrow{\delta_3} P_2 \xrightarrow{\delta_2} P_1 \xrightarrow{\delta_1} P_0 \xrightarrow{\delta_0} B \rightarrow 0.$$

Apply the tensor functor $A \otimes_R _$:

$$\cdots \xrightarrow{1 \otimes \delta_3} A \otimes_R P_2 \xrightarrow{1 \otimes \delta_2} A \otimes_R P_1 \xrightarrow{1 \otimes \delta_1} A \otimes_R P_0 \xrightarrow{1 \otimes \delta_0} A \otimes_R B \rightarrow 0.$$

Computing $\text{Tor}_n^R(A, B)$

Start with a **projective resolution** of B :

$$\cdots \xrightarrow{\delta_3} P_2 \xrightarrow{\delta_2} P_1 \xrightarrow{\delta_1} P_0 \xrightarrow{\delta_0} B \rightarrow 0.$$

Apply the tensor functor $A \otimes_R _$:

$$\cdots \xrightarrow{1 \otimes \delta_3} A \otimes_R P_2 \xrightarrow{1 \otimes \delta_2} A \otimes_R P_1 \xrightarrow{1 \otimes \delta_1} A \otimes_R P_0 \xrightarrow{1 \otimes \delta_0} A \otimes_R B \rightarrow 0.$$

For $n > 0$, $\text{Tor}_n^R(A, B) = \text{homology at position } n = \ker(1 \otimes \delta_n) / \text{im}(1 \otimes \delta_{n+1})$.

$\text{Tor}_0^R(A, B) = (A \otimes_R P_0) / \text{im}(1 \otimes \delta_1) \cong A \otimes_R B$.

Computing $\text{Tor}_n^R(A, B)$

Start with a **projective resolution** of B :

$$\cdots \xrightarrow{\delta_3} P_2 \xrightarrow{\delta_2} P_1 \xrightarrow{\delta_1} P_0 \xrightarrow{\delta_0} B \rightarrow 0.$$

Apply the tensor functor $A \otimes_R _$:

$$\cdots \xrightarrow{1 \otimes \delta_3} A \otimes_R P_2 \xrightarrow{1 \otimes \delta_2} A \otimes_R P_1 \xrightarrow{1 \otimes \delta_1} A \otimes_R P_0 \xrightarrow{1 \otimes \delta_0} A \otimes_R B \rightarrow 0.$$

For $n > 0$, $\text{Tor}_n^R(A, B) = \text{homology at position } n = \ker(1 \otimes \delta_n) / \text{im}(1 \otimes \delta_{n+1})$.

$\text{Tor}_0^R(A, B) = (A \otimes_R P_0) / \text{im}(1 \otimes \delta_1) \cong A \otimes_R B$.

Applying $A \otimes_R _$ to $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$, the loss of exactness is repaired by Tor in the sense that there exist maps yielding the “LES in homology”:

Computing $\text{Tor}_n^R(A, B)$

Start with a **projective resolution** of B :

$$\cdots \xrightarrow{\delta_3} P_2 \xrightarrow{\delta_2} P_1 \xrightarrow{\delta_1} P_0 \xrightarrow{\delta_0} B \rightarrow 0.$$

Apply the tensor functor $A \otimes_R _$:

$$\cdots \xrightarrow{1 \otimes \delta_3} A \otimes_R P_2 \xrightarrow{1 \otimes \delta_2} A \otimes_R P_1 \xrightarrow{1 \otimes \delta_1} A \otimes_R P_0 \xrightarrow{1 \otimes \delta_0} A \otimes_R B \rightarrow 0.$$

For $n > 0$, $\text{Tor}_n^R(A, B) = \text{homology at position } n = \ker(1 \otimes \delta_n) / \text{im}(1 \otimes \delta_{n+1})$.

$\text{Tor}_0^R(A, B) = (A \otimes_R P_0) / \text{im}(1 \otimes \delta_1) \cong A \otimes_R B$.

Applying $A \otimes_R _$ to $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$, the loss of exactness is repaired by Tor in the sense that there exist maps yielding the “LES in homology”:

$$\begin{array}{ccccccc} \cdots \rightarrow & \text{Tor}_2^R(A, N) \rightarrow & \text{Tor}_1^R(A, L) \rightarrow & \text{Tor}_1^R(A, M) \rightarrow & & & \\ & \rightarrow \text{Tor}_1^R(A, N) & \rightarrow A \otimes_R L \rightarrow & A \otimes_R M \rightarrow & A \otimes_R N \rightarrow & 0. \end{array}$$

$\mathrm{Tor}_1^{\mathbb{Z}}$ computes torsion

$\mathrm{Tor}_1^{\mathbb{Z}}$ computes torsion

$$\mathrm{Tor}_1^{\mathbb{Z}}(A, \mathbb{Z}/d\mathbb{Z})$$

$\mathrm{Tor}_1^{\mathbb{Z}}$ computes torsion

$$\mathrm{Tor}_1^{\mathbb{Z}}(A, \mathbb{Z}/d\mathbb{Z}) = \{a \in A \mid da = 0\}$$

$\mathrm{Tor}_1^{\mathbb{Z}}$ computes torsion

$$\mathrm{Tor}_1^{\mathbb{Z}}(A, \mathbb{Z}/d\mathbb{Z}) = \{a \in A \mid da = 0\} = {}_dA$$

$\mathrm{Tor}_1^{\mathbb{Z}}$ computes torsion

$$\mathrm{Tor}_1^{\mathbb{Z}}(A, \mathbb{Z}/d\mathbb{Z}) = \{a \in A \mid da = 0\} = {}_dA = \ker(d \times _ : A \rightarrow A).$$

$\mathrm{Tor}_1^{\mathbb{Z}}$ computes torsion

$$\mathrm{Tor}_1^{\mathbb{Z}}(A, \mathbb{Z}/d\mathbb{Z}) = \{a \in A \mid da = 0\} = {}_dA = \ker(d \times _ : A \rightarrow A).$$

The sequence $\cdots \rightarrow 0 \rightarrow 0 \rightarrow \mathbb{Z} \xrightarrow{d \times} \mathbb{Z} \rightarrow \mathbb{Z}/d\mathbb{Z} \rightarrow 0$ is a projective/free resolution of $\mathbb{Z}/d\mathbb{Z}$.

$\text{Tor}_1^{\mathbb{Z}}$ computes torsion

$$\text{Tor}_1^{\mathbb{Z}}(A, \mathbb{Z}/d\mathbb{Z}) = \{a \in A \mid da = 0\} = {}_dA = \ker(d \times _ : A \rightarrow A).$$

The sequence $\cdots \rightarrow 0 \rightarrow 0 \rightarrow \mathbb{Z} \xrightarrow{d \times} \mathbb{Z} \rightarrow \mathbb{Z}/d\mathbb{Z} \rightarrow 0$ is a projective/free resolution of $\mathbb{Z}/d\mathbb{Z}$.

Tensoring with A yields the complex

$$\cdots \rightarrow 0 \rightarrow 0 \rightarrow \underline{A} \xrightarrow{d \times} A \rightarrow A/dA \rightarrow 0,$$

$\mathrm{Tor}_1^{\mathbb{Z}}$ computes torsion

$$\mathrm{Tor}_1^{\mathbb{Z}}(A, \mathbb{Z}/d\mathbb{Z}) = \{a \in A \mid da = 0\} = {}_dA = \ker(d \times _ : A \rightarrow A).$$

The sequence $\cdots \rightarrow 0 \rightarrow 0 \rightarrow \mathbb{Z} \xrightarrow{d \times} \mathbb{Z} \rightarrow \mathbb{Z}/d\mathbb{Z} \rightarrow 0$ is a projective/free resolution of $\mathbb{Z}/d\mathbb{Z}$.

Tensoring with A yields the complex

$$\cdots \rightarrow 0 \rightarrow 0 \rightarrow \underline{A} \xrightarrow{d \times} A \rightarrow A/dA \rightarrow 0,$$

which can only fail to be exact at the underlined term.

$\mathrm{Tor}_1^{\mathbb{Z}}$ computes torsion

$$\mathrm{Tor}_1^{\mathbb{Z}}(A, \mathbb{Z}/d\mathbb{Z}) = \{a \in A \mid da = 0\} = {}_dA = \ker(d \times _ : A \rightarrow A).$$

The sequence $\cdots \rightarrow 0 \rightarrow 0 \rightarrow \mathbb{Z} \xrightarrow{d \times} \mathbb{Z} \rightarrow \mathbb{Z}/d\mathbb{Z} \rightarrow 0$ is a projective/free resolution of $\mathbb{Z}/d\mathbb{Z}$.

Tensoring with A yields the complex

$$\cdots \rightarrow 0 \rightarrow 0 \rightarrow \underline{A} \xrightarrow{d \times} A \rightarrow A/dA \rightarrow 0,$$

which can only fail to be exact at the underlined term. Homology at that point is $\ker(d \times _)$

$\mathrm{Tor}_1^{\mathbb{Z}}$ computes torsion

$$\mathrm{Tor}_1^{\mathbb{Z}}(A, \mathbb{Z}/d\mathbb{Z}) = \{a \in A \mid da = 0\} = {}_dA = \ker(d \times _ : A \rightarrow A).$$

The sequence $\cdots \rightarrow 0 \rightarrow 0 \rightarrow \mathbb{Z} \xrightarrow{d \times} \mathbb{Z} \rightarrow \mathbb{Z}/d\mathbb{Z} \rightarrow 0$ is a projective/free resolution of $\mathbb{Z}/d\mathbb{Z}$.

Tensoring with A yields the complex

$$\cdots \rightarrow 0 \rightarrow 0 \rightarrow \underline{A} \xrightarrow{d \times} A \rightarrow A/dA \rightarrow 0,$$

which can only fail to be exact at the underlined term. Homology at that point is $\ker(d \times _) = \mathrm{Tor}_1^{\mathbb{Z}}(A, \mathbb{Z}/d\mathbb{Z})$.

$\mathrm{Tor}_1^{\mathbb{Z}}$ computes torsion

$$\mathrm{Tor}_1^{\mathbb{Z}}(A, \mathbb{Z}/d\mathbb{Z}) = \{a \in A \mid da = 0\} = {}_dA = \ker(d \times _ : A \rightarrow A).$$

The sequence $\cdots \rightarrow 0 \rightarrow 0 \rightarrow \mathbb{Z} \xrightarrow{d \times} \mathbb{Z} \rightarrow \mathbb{Z}/d\mathbb{Z} \rightarrow 0$ is a projective/free resolution of $\mathbb{Z}/d\mathbb{Z}$.

Tensoring with A yields the complex

$$\cdots \rightarrow 0 \rightarrow 0 \rightarrow \underline{A} \xrightarrow{d \times} A \rightarrow A/dA \rightarrow 0,$$

which can only fail to be exact at the underlined term. Homology at that point is $\ker(d \times _) = \mathrm{Tor}_1^{\mathbb{Z}}(A, \mathbb{Z}/d\mathbb{Z})$. We also see that $\mathrm{Tor}_n^{\mathbb{Z}}(A, \mathbb{Z}/d\mathbb{Z}) = 0$ for $n > 1$.

$\mathrm{Tor}_1^{\mathbb{Z}}$ computes torsion

$$\mathrm{Tor}_1^{\mathbb{Z}}(A, \mathbb{Z}/d\mathbb{Z}) = \{a \in A \mid da = 0\} = {}_dA = \ker(d \times _ : A \rightarrow A).$$

The sequence $\cdots \rightarrow 0 \rightarrow 0 \rightarrow \mathbb{Z} \xrightarrow{d \times} \mathbb{Z} \rightarrow \mathbb{Z}/d\mathbb{Z} \rightarrow 0$ is a projective/free resolution of $\mathbb{Z}/d\mathbb{Z}$.

Tensoring with A yields the complex

$$\cdots \rightarrow 0 \rightarrow 0 \rightarrow \underline{A} \xrightarrow{d \times} A \rightarrow A/dA \rightarrow 0,$$

which can only fail to be exact at the underlined term. Homology at that point is $\ker(d \times _) = \mathrm{Tor}_1^{\mathbb{Z}}(A, \mathbb{Z}/d\mathbb{Z})$. We also see that $\mathrm{Tor}_n^{\mathbb{Z}}(A, \mathbb{Z}/d\mathbb{Z}) = 0$ for $n > 1$.

Generalizations

Everything from the previous slide works in exactly the same way if we replace \mathbb{Z} by some other commutative ring R , provided $d \in R$ is not a zero divisor.

Generalizations

Everything from the previous slide works in exactly the same way if we replace \mathbb{Z} by some other commutative ring R , provided $d \in R$ is not a zero divisor. We then get $\mathrm{Tor}_1^R(A, R/(d)) = {}_dA$ and $\mathrm{Tor}_n^R(A, R/(d)) = 0$ for $n > 1$.

Generalizations

Everything from the previous slide works in exactly the same way if we replace \mathbb{Z} by some other commutative ring R , provided $d \in R$ is not a zero divisor. We then get $\mathrm{Tor}_1^R(A, R/(d)) = {}_dA$ and $\mathrm{Tor}_n^R(A, R/(d)) = 0$ for $n > 1$.

With a little work, one can show that $\mathrm{Tor}_1^{\mathbb{Z}}(A, \mathbb{Q}/\mathbb{Z})$ is isomorphic to the subgroup of all torsion elements of A .

Everything from the previous slide works in exactly the same way if we replace \mathbb{Z} by some other commutative ring R , provided $d \in R$ is not a zero divisor. We then get $\mathrm{Tor}_1^R(A, R/(d)) = {}_dA$ and $\mathrm{Tor}_n^R(A, R/(d)) = 0$ for $n > 1$.

With a little work, one can show that $\mathrm{Tor}_1^{\mathbb{Z}}(A, \mathbb{Q}/\mathbb{Z})$ is isomorphic to the subgroup of all torsion elements of A .

Auslander showed in the 60's that $\mathrm{Tor}(M, N)$ is related to torsion in another way.

Everything from the previous slide works in exactly the same way if we replace \mathbb{Z} by some other commutative ring R , provided $d \in R$ is not a zero divisor. We then get $\mathrm{Tor}_1^R(A, R/(d)) = {}_dA$ and $\mathrm{Tor}_n^R(A, R/(d)) = 0$ for $n > 1$.

With a little work, one can show that $\mathrm{Tor}_1^{\mathbb{Z}}(A, \mathbb{Q}/\mathbb{Z})$ is isomorphic to the subgroup of all torsion elements of A .

Auslander showed in the 60's that $\mathrm{Tor}(M, N)$ is related to torsion in another way.

Theorem. Let R be a regular local ring that is not a field, and let M and N be finitely generated R -modules. TFAE:

- ① $M \otimes_R N$ is torsion-free.

Everything from the previous slide works in exactly the same way if we replace \mathbb{Z} by some other commutative ring R , provided $d \in R$ is not a zero divisor. We then get $\mathrm{Tor}_1^R(A, R/(d)) = {}_dA$ and $\mathrm{Tor}_n^R(A, R/(d)) = 0$ for $n > 1$.

With a little work, one can show that $\mathrm{Tor}_1^{\mathbb{Z}}(A, \mathbb{Q}/\mathbb{Z})$ is isomorphic to the subgroup of all torsion elements of A .

Auslander showed in the 60's that $\mathrm{Tor}(M, N)$ is related to torsion in another way.

Theorem. Let R be a regular local ring that is not a field, and let M and N be finitely generated R -modules. TFAE:

- ① $M \otimes_R N$ is torsion-free.

Everything from the previous slide works in exactly the same way if we replace \mathbb{Z} by some other commutative ring R , provided $d \in R$ is not a zero divisor. We then get $\mathrm{Tor}_1^R(A, R/(d)) = {}_dA$ and $\mathrm{Tor}_n^R(A, R/(d)) = 0$ for $n > 1$.

With a little work, one can show that $\mathrm{Tor}_1^{\mathbb{Z}}(A, \mathbb{Q}/\mathbb{Z})$ is isomorphic to the subgroup of all torsion elements of A .

Auslander showed in the 60's that $\mathrm{Tor}(M, N)$ is related to torsion in another way.

Theorem. Let R be a regular local ring that is not a field, and let M and N be finitely generated R -modules. TFAE:

- ① $M \otimes_R N$ is torsion-free.
- ② ① M and N are torsion-free.

Everything from the previous slide works in exactly the same way if we replace \mathbb{Z} by some other commutative ring R , provided $d \in R$ is not a zero divisor. We then get $\mathrm{Tor}_1^R(A, R/(d)) = {}_dA$ and $\mathrm{Tor}_n^R(A, R/(d)) = 0$ for $n > 1$.

With a little work, one can show that $\mathrm{Tor}_1^{\mathbb{Z}}(A, \mathbb{Q}/\mathbb{Z})$ is isomorphic to the subgroup of all torsion elements of A .

Auslander showed in the 60's that $\mathrm{Tor}(M, N)$ is related to torsion in another way.

Theorem. Let R be a regular local ring that is not a field, and let M and N be finitely generated R -modules. TFAE:

- ① $M \otimes_R N$ is torsion-free.
- ②
 - ① M and N are torsion-free.
 - ② $\mathrm{Tor}_n^R(M, N) = 0$ for all $n > 0$.

Everything from the previous slide works in exactly the same way if we replace \mathbb{Z} by some other commutative ring R , provided $d \in R$ is not a zero divisor. We then get $\mathrm{Tor}_1^R(A, R/(d)) = {}_dA$ and $\mathrm{Tor}_n^R(A, R/(d)) = 0$ for $n > 1$.

With a little work, one can show that $\mathrm{Tor}_1^{\mathbb{Z}}(A, \mathbb{Q}/\mathbb{Z})$ is isomorphic to the subgroup of all torsion elements of A .

Auslander showed in the 60's that $\mathrm{Tor}(M, N)$ is related to torsion in another way.

Theorem. Let R be a regular local ring that is not a field, and let M and N be finitely generated R -modules. TFAE:

- ① $M \otimes_R N$ is torsion-free.
- ②
 - ① M and N are torsion-free.
 - ② $\mathrm{Tor}_n^R(M, N) = 0$ for all $n > 0$.
 - ③ The projective dimensions of M and N sum to the projective dimension of $M \otimes_R N$, and this is less than the Krull dimension of R .

Computing $\text{Ext}_R^n(A, B)$

Computing $\text{Ext}_R^n(A, B)$

Start with a **projective resolution** of A :

$$\cdots \xrightarrow{\delta_3} P_2 \xrightarrow{\delta_2} P_1 \xrightarrow{\delta_1} P_0 \xrightarrow{\delta_0} A \rightarrow 0.$$

Computing $Ext_R^n(A, B)$

Start with a **projective resolution** of A :

$$\cdots \xrightarrow{\delta_3} P_2 \xrightarrow{\delta_2} P_1 \xrightarrow{\delta_1} P_0 \xrightarrow{\delta_0} A \rightarrow 0.$$

Apply contravariant the hom functor $\text{Hom}_R(_, B)$:

$$0 \rightarrow \text{Hom}_R(A, B) \xrightarrow{\circ \delta_0} \text{Hom}_R(P_0, B) \xrightarrow{\circ \delta_1} \text{Hom}_R(P_1, B) \xrightarrow{\circ \delta_2} \cdots$$

Computing $\text{Ext}_R^n(A, B)$

Start with a **projective resolution** of A :

$$\cdots \xrightarrow{\delta_3} P_2 \xrightarrow{\delta_2} P_1 \xrightarrow{\delta_1} P_0 \xrightarrow{\delta_0} A \rightarrow 0.$$

Apply contravariant the hom functor $\text{Hom}_R(_, B)$:

$$0 \rightarrow \text{Hom}_R(A, B) \xrightarrow{\circ \delta_0} \text{Hom}_R(P_0, B) \xrightarrow{\circ \delta_1} \text{Hom}_R(P_1, B) \xrightarrow{\circ \delta_2} \cdots$$

For $n > 0$, $\text{Ext}_R^n(A, B) = \text{homology at } n = \ker(_ \circ \delta_{n+1}) / \text{im}(_ \circ \delta_n).$

Computing $\text{Ext}_R^n(A, B)$

Start with a **projective resolution** of A :

$$\cdots \xrightarrow{\delta_3} P_2 \xrightarrow{\delta_2} P_1 \xrightarrow{\delta_1} P_0 \xrightarrow{\delta_0} A \rightarrow 0.$$

Apply contravariant the hom functor $\text{Hom}_R(_, B)$:

$$0 \rightarrow \text{Hom}_R(A, B) \xrightarrow{\circ \delta_0} \text{Hom}_R(P_0, B) \xrightarrow{\circ \delta_1} \text{Hom}_R(P_1, B) \xrightarrow{\circ \delta_2} \cdots$$

For $n > 0$, $\text{Ext}_R^n(A, B) = \text{homology at } n = \ker(_ \circ \delta_{n+1}) / \text{im}(_ \circ \delta_n)$.

$\text{Ext}_R^0(A, B) = \ker(_ \circ \delta_1) \cong \text{Hom}_R(A, B)$.

Computing $\text{Ext}_R^n(A, B)$

Start with a **projective resolution** of A :

$$\cdots \xrightarrow{\delta_3} P_2 \xrightarrow{\delta_2} P_1 \xrightarrow{\delta_1} P_0 \xrightarrow{\delta_0} A \rightarrow 0.$$

Apply contravariant the hom functor $\text{Hom}_R(_, B)$:

$$0 \rightarrow \text{Hom}_R(A, B) \xrightarrow{\circ \delta_0} \text{Hom}_R(P_0, B) \xrightarrow{\circ \delta_1} \text{Hom}_R(P_1, B) \xrightarrow{\circ \delta_2} \cdots$$

For $n > 0$, $\text{Ext}_R^n(A, B) = \text{homology at } n = \ker(_ \circ \delta_{n+1}) / \text{im}(_ \circ \delta_n)$.

$\text{Ext}_R^0(A, B) = \ker(_ \circ \delta_1) \cong \text{Hom}_R(A, B)$.

Ext can be computed using covariant hom instead, $\text{Hom}_R(A, _)$,

Computing $\text{Ext}_R^n(A, B)$

Start with a **projective resolution** of A :

$$\cdots \xrightarrow{\delta_3} P_2 \xrightarrow{\delta_2} P_1 \xrightarrow{\delta_1} P_0 \xrightarrow{\delta_0} A \rightarrow 0.$$

Apply contravariant the hom functor $\text{Hom}_R(_, B)$:

$$0 \rightarrow \text{Hom}_R(A, B) \xrightarrow{\circ \delta_0} \text{Hom}_R(P_0, B) \xrightarrow{\circ \delta_1} \text{Hom}_R(P_1, B) \xrightarrow{\circ \delta_2} \cdots$$

For $n > 0$, $\text{Ext}_R^n(A, B) = \text{homology at } n = \ker(_ \circ \delta_{n+1}) / \text{im}(_ \circ \delta_n)$.

$\text{Ext}_R^0(A, B) = \ker(_ \circ \delta_1) \cong \text{Hom}_R(A, B)$.

Ext can be computed using covariant hom instead, $\text{Hom}_R(A, _)$, but one should apply covariant hom to an injective resolution of B instead of a projective resolution of B (or A).

An Example: Ext computes extension groups

An Example: Ext computes extension groups

$\text{Ext}_R^1(A, B)$ computes the group of inequivalent extensions

$$0 \rightarrow B \rightarrow E \rightarrow A \rightarrow 0.$$

An Example: Ext computes extension groups

$\text{Ext}_R^1(A, B)$ computes the group of inequivalent extensions

$0 \rightarrow B \rightarrow E \rightarrow A \rightarrow 0$. Concrete example: $0 \rightarrow \mathbb{Z}_2 \rightarrow E \rightarrow \mathbb{Z}_4 \rightarrow 0$.

An Example: Ext computes extension groups

$\text{Ext}_R^1(A, B)$ computes the group of inequivalent extensions

$0 \rightarrow B \rightarrow E \rightarrow A \rightarrow 0$. Concrete example: $0 \rightarrow \mathbb{Z}_2 \rightarrow E \rightarrow \mathbb{Z}_4 \rightarrow 0$.

Exercise.

An Example: Ext computes extension groups

$\text{Ext}_R^1(A, B)$ computes the group of inequivalent extensions

$0 \rightarrow B \rightarrow E \rightarrow A \rightarrow 0$. Concrete example: $0 \rightarrow \mathbb{Z}_2 \rightarrow E \rightarrow \mathbb{Z}_4 \rightarrow 0$.

Exercise. Compute $\text{Ext}_{\mathbb{Z}}^1(\mathbb{Z}_4, \mathbb{Z}_2)$

An Example: Ext computes extension groups

$\text{Ext}_R^1(A, B)$ computes the group of inequivalent extensions

$0 \rightarrow B \rightarrow E \rightarrow A \rightarrow 0$. Concrete example: $0 \rightarrow \mathbb{Z}_2 \rightarrow E \rightarrow \mathbb{Z}_4 \rightarrow 0$.

Exercise. Compute $\text{Ext}_{\mathbb{Z}}^1(\mathbb{Z}_4, \mathbb{Z}_2)$

The sequence $\cdots \rightarrow 0 \rightarrow 0 \rightarrow \mathbb{Z} \xrightarrow{4\times} \mathbb{Z} \rightarrow \mathbb{Z}_4 \rightarrow 0$ is a projective/free resolution of \mathbb{Z}_4 .

An Example: Ext computes extension groups

$\text{Ext}_R^1(A, B)$ computes the group of inequivalent extensions

$0 \rightarrow B \rightarrow E \rightarrow A \rightarrow 0$. Concrete example: $0 \rightarrow \mathbb{Z}_2 \rightarrow E \rightarrow \mathbb{Z}_4 \rightarrow 0$.

Exercise. Compute $\text{Ext}_{\mathbb{Z}}^1(\mathbb{Z}_4, \mathbb{Z}_2)$

The sequence $\cdots \rightarrow 0 \rightarrow 0 \rightarrow \mathbb{Z} \xrightarrow{4\times} \mathbb{Z} \rightarrow \mathbb{Z}_4 \rightarrow 0$ is a projective/free resolution of \mathbb{Z}_4 . Apply $\text{Hom}_R(_, \mathbb{Z}_2)$:

An Example: Ext computes extension groups

$\text{Ext}_R^1(A, B)$ computes the group of inequivalent extensions

$0 \rightarrow B \rightarrow E \rightarrow A \rightarrow 0$. Concrete example: $0 \rightarrow \mathbb{Z}_2 \rightarrow E \rightarrow \mathbb{Z}_4 \rightarrow 0$.

Exercise. Compute $\text{Ext}_{\mathbb{Z}}^1(\mathbb{Z}_4, \mathbb{Z}_2)$

The sequence $\cdots \rightarrow 0 \rightarrow 0 \rightarrow \mathbb{Z} \xrightarrow{4\times} \mathbb{Z} \rightarrow \mathbb{Z}_4 \rightarrow 0$ is a projective/free resolution of \mathbb{Z}_4 . Apply $\text{Hom}_R(_, \mathbb{Z}_2)$:

$$0 \rightarrow \text{Hom}_R(\mathbb{Z}_4, \mathbb{Z}_2) \rightarrow \text{Hom}_R(\mathbb{Z}, \mathbb{Z}_2) \xrightarrow{4\times} \underline{\text{Hom}_R(\mathbb{Z}, \mathbb{Z}_2)} \rightarrow 0 \rightarrow \cdots$$

An Example: Ext computes extension groups

$\text{Ext}_R^1(A, B)$ computes the group of inequivalent extensions

$0 \rightarrow B \rightarrow E \rightarrow A \rightarrow 0$. Concrete example: $0 \rightarrow \mathbb{Z}_2 \rightarrow E \rightarrow \mathbb{Z}_4 \rightarrow 0$.

Exercise. Compute $\text{Ext}_{\mathbb{Z}}^1(\mathbb{Z}_4, \mathbb{Z}_2)$

The sequence $\cdots \rightarrow 0 \rightarrow 0 \rightarrow \mathbb{Z} \xrightarrow{4\times} \mathbb{Z} \rightarrow \mathbb{Z}_4 \rightarrow 0$ is a projective/free resolution of \mathbb{Z}_4 . Apply $\text{Hom}_R(, \mathbb{Z}_2)$:

$$0 \rightarrow \text{Hom}_R(\mathbb{Z}_4, \mathbb{Z}_2) \rightarrow \text{Hom}_R(\mathbb{Z}, \mathbb{Z}_2) \xrightarrow{4\times} \underline{\text{Hom}_R(\mathbb{Z}, \mathbb{Z}_2)} \rightarrow 0 \rightarrow \cdots$$

$$\text{Ext}_{\mathbb{Z}}^1(\mathbb{Z}_4, \mathbb{Z}_2) = \mathbb{Z}_2/0 \cong \mathbb{Z}_2.$$

An Example: Ext computes extension groups

$\text{Ext}_R^1(A, B)$ computes the group of inequivalent extensions

$0 \rightarrow B \rightarrow E \rightarrow A \rightarrow 0$. Concrete example: $0 \rightarrow \mathbb{Z}_2 \rightarrow E \rightarrow \mathbb{Z}_4 \rightarrow 0$.

Exercise. Compute $\text{Ext}_{\mathbb{Z}}^1(\mathbb{Z}_4, \mathbb{Z}_2)$

The sequence $\cdots \rightarrow 0 \rightarrow 0 \rightarrow \mathbb{Z} \xrightarrow{4\times} \mathbb{Z} \rightarrow \mathbb{Z}_4 \rightarrow 0$ is a projective/free resolution of \mathbb{Z}_4 . Apply $\text{Hom}_R(_, \mathbb{Z}_2)$:

$$0 \rightarrow \text{Hom}_R(\mathbb{Z}_4, \mathbb{Z}_2) \rightarrow \text{Hom}_R(\mathbb{Z}, \mathbb{Z}_2) \xrightarrow{4\times} \underline{\text{Hom}_R(\mathbb{Z}, \mathbb{Z}_2)} \rightarrow 0 \rightarrow \cdots$$

$$\text{Ext}_{\mathbb{Z}}^1(\mathbb{Z}_4, \mathbb{Z}_2) = \mathbb{Z}_2/0 \cong \mathbb{Z}_2.$$

$\text{Ext}_{\mathbb{Z}}^n(A, B)$ can be interpreted as the groups of inequivalent n -extensions:

$$0 \rightarrow B \rightarrow E_n \rightarrow \cdots \rightarrow E_1 \rightarrow A \rightarrow 0.$$