

Localization

The problem.

The problem. Given a ring A and elements $u, v \in R$, find the most general extension of A in which u and v are units.

The problem. Given a ring A and elements $u, v \in R$, find the most general extension of A in which u and v are units.

The obvious solution.

The problem. Given a ring A and elements $u, v \in R$, find the most general extension of A in which u and v are units.

The obvious solution. If $A = \langle G \mid R \rangle$, then the solution is

$$\langle G \cup \{x, y\} \mid R \cup \{ux = 1 = vy\} \rangle.$$

The problem. Given a ring A and elements $u, v \in R$, find the most general extension of A in which u and v are units.

The obvious solution. If $A = \langle G \mid R \rangle$, then the solution is

$$\langle G \cup \{x, y\} \mid R \cup \{ux = 1 = vy\} \rangle.$$

$$(\langle G \mid R \rangle = A.$$

The problem. Given a ring A and elements $u, v \in R$, find the most general extension of A in which u and v are units.

The obvious solution. If $A = \langle G \mid R \rangle$, then the solution is

$$\langle G \cup \{x, y\} \mid R \cup \{ux = 1 = vy\} \rangle.$$

$$(\langle G \mid R \rangle = A. \langle G \cup \{x, y\} \mid R \rangle \cong A \otimes_{\mathbb{Z}} \mathbb{Z}[x, y])$$

The problem. Given a ring A and elements $u, v \in R$, find the most general extension of A in which u and v are units.

The obvious solution. If $A = \langle G \mid R \rangle$, then the solution is

$$\langle G \cup \{x, y\} \mid R \cup \{ux = 1 = vy\} \rangle.$$

$$(\langle G \mid R \rangle = A. \langle G \cup \{x, y\} \mid R \rangle \cong A \otimes_{\mathbb{Z}} \mathbb{Z}[x, y] \cong A[x, y].$$

The problem. Given a ring A and elements $u, v \in R$, find the most general extension of A in which u and v are units.

The obvious solution. If $A = \langle G \mid R \rangle$, then the solution is

$$\langle G \cup \{x, y\} \mid R \cup \{ux = 1 = vy\} \rangle.$$

$$\begin{aligned} (\langle G \mid R \rangle = A. \langle G \cup \{x, y\} \mid R \rangle &\cong A \otimes_{\mathbb{Z}} \mathbb{Z}[x, y] \cong A[x, y]. \\ \langle G \cup \{x, y\} \mid R \cup \{ux = 1 = vy\} \rangle &\cong A[x, y]/(ux - 1, vy - 1).) \end{aligned}$$

The problem. Given a ring A and elements $u, v \in R$, find the most general extension of A in which u and v are units.

The obvious solution. If $A = \langle G \mid R \rangle$, then the solution is

$$\langle G \cup \{x, y\} \mid R \cup \{ux = 1 = vy\} \rangle.$$

$$\begin{aligned} (\langle G \mid R \rangle = A. \langle G \cup \{x, y\} \mid R \rangle &\cong A \otimes_{\mathbb{Z}} \mathbb{Z}[x, y] \cong A[x, y]. \\ \langle G \cup \{x, y\} \mid R \cup \{ux = 1 = vy\} \rangle &\cong A[x, y]/(ux - 1, vy - 1).) \end{aligned}$$

Elements of $A[x, y]/(ux - 1, vy - 1)$

The problem. Given a ring A and elements $u, v \in R$, find the most general extension of A in which u and v are units.

The obvious solution. If $A = \langle G \mid R \rangle$, then the solution is

$$\langle G \cup \{x, y\} \mid R \cup \{ux = 1 = vy\} \rangle.$$

$$\begin{aligned} (\langle G \mid R \rangle = A. \langle G \cup \{x, y\} \mid R \rangle &\cong A \otimes_{\mathbb{Z}} \mathbb{Z}[x, y] \cong A[x, y]. \\ \langle G \cup \{x, y\} \mid R \cup \{ux = 1 = vy\} \rangle &\cong A[x, y]/(ux - 1, vy - 1).) \end{aligned}$$

Elements of $A[x, y]/(ux - 1, vy - 1) = A[u^{-1}, v^{-1}]$

The problem. Given a ring A and elements $u, v \in R$, find the most general extension of A in which u and v are units.

The obvious solution. If $A = \langle G \mid R \rangle$, then the solution is

$$\langle G \cup \{x, y\} \mid R \cup \{ux = 1 = vy\} \rangle.$$

$$(\langle G \mid R \rangle = A. \langle G \cup \{x, y\} \mid R \rangle \cong A \otimes_{\mathbb{Z}} \mathbb{Z}[x, y] \cong A[x, y].$$

$$\langle G \cup \{x, y\} \mid R \cup \{ux = 1 = vy\} \rangle \cong A[x, y]/(ux - 1, vy - 1).)$$

Elements of $A[x, y]/(ux - 1, vy - 1) = A[u^{-1}, v^{-1}]$ simplify to the form $a(u^{-1})^m(v^{-1})^n$.

The problem. Given a ring A and elements $u, v \in R$, find the most general extension of A in which u and v are units.

The obvious solution. If $A = \langle G \mid R \rangle$, then the solution is

$$\langle G \cup \{x, y\} \mid R \cup \{ux = 1 = vy\} \rangle.$$

$$\begin{aligned} (\langle G \mid R \rangle = A. \langle G \cup \{x, y\} \mid R \rangle &\cong A \otimes_{\mathbb{Z}} \mathbb{Z}[x, y] \cong A[x, y]. \\ \langle G \cup \{x, y\} \mid R \cup \{ux = 1 = vy\} \rangle &\cong A[x, y]/(ux - 1, vy - 1).) \end{aligned}$$

Elements of $A[x, y]/(ux - 1, vy - 1) = A[u^{-1}, v^{-1}]$ simplify to the form $a(u^{-1})^m(v^{-1})^n$.

Simpler, more direct (and standard) approach.

The problem. Given a ring A and elements $u, v \in R$, find the most general extension of A in which u and v are units.

The obvious solution. If $A = \langle G \mid R \rangle$, then the solution is

$$\langle G \cup \{x, y\} \mid R \cup \{ux = 1 = vy\} \rangle.$$

$$\begin{aligned} (\langle G \mid R \rangle = A. \langle G \cup \{x, y\} \mid R \rangle &\cong A \otimes_{\mathbb{Z}} \mathbb{Z}[x, y] \cong A[x, y]. \\ \langle G \cup \{x, y\} \mid R \cup \{ux = 1 = vy\} \rangle &\cong A[x, y]/(ux - 1, vy - 1).) \end{aligned}$$

Elements of $A[x, y]/(ux - 1, vy - 1) = A[u^{-1}, v^{-1}]$ simplify to the form $a(u^{-1})^m(v^{-1})^n$.

Simpler, more direct (and standard) approach.

- 1 Start instead with $S = \{u, v\}^* = \{1, u, v, u^2, uv, \dots\}$, an MC set.

The problem. Given a ring A and elements $u, v \in R$, find the most general extension of A in which u and v are units.

The obvious solution. If $A = \langle G \mid R \rangle$, then the solution is

$$\langle G \cup \{x, y\} \mid R \cup \{ux = 1 = vy\} \rangle.$$

$$\begin{aligned} \langle \langle G \mid R \rangle = A. \langle G \cup \{x, y\} \mid R \rangle &\cong A \otimes_{\mathbb{Z}} \mathbb{Z}[x, y] \cong A[x, y]. \\ \langle G \cup \{x, y\} \mid R \cup \{ux = 1 = vy\} \rangle &\cong A[x, y]/(ux - 1, vy - 1). \end{aligned}$$

Elements of $A[x, y]/(ux - 1, vy - 1) = A[u^{-1}, v^{-1}]$ simplify to the form $a(u^{-1})^m(v^{-1})^n$.

Simpler, more direct (and standard) approach.

- 1 Start instead with $S = \{u, v\}^* = \{1, u, v, u^2, uv, \dots\}$, an MC set.
- 2 Define $S^{-1}A$ to consist of ordered pairs $a/s := (a, s)$ modulo equiv. reln.
 $a/s \sim b/t$ iff $(\exists u \in S)(u(sb - ta) = 0)$.

The problem. Given a ring A and elements $u, v \in R$, find the most general extension of A in which u and v are units.

The obvious solution. If $A = \langle G \mid R \rangle$, then the solution is

$$\langle G \cup \{x, y\} \mid R \cup \{ux = 1 = vy\} \rangle.$$

$$\begin{aligned} \langle \langle G \mid R \rangle = A. \langle G \cup \{x, y\} \mid R \rangle &\cong A \otimes_{\mathbb{Z}} \mathbb{Z}[x, y] \cong A[x, y]. \\ \langle G \cup \{x, y\} \mid R \cup \{ux = 1 = vy\} \rangle &\cong A[x, y]/(ux - 1, vy - 1). \end{aligned}$$

Elements of $A[x, y]/(ux - 1, vy - 1) = A[u^{-1}, v^{-1}]$ simplify to the form $a(u^{-1})^m(v^{-1})^n$.

Simpler, more direct (and standard) approach.

- 1 Start instead with $S = \{u, v\}^* = \{1, u, v, u^2, uv, \dots\}$, an MC set.
- 2 Define $S^{-1}A$ to consist of ordered pairs $a/s := (a, s)$ modulo equiv. reln.
 $a/s \sim b/t$ iff $(\exists u \in S)(u(sb - ta) = 0)$.

The problem. Given a ring A and elements $u, v \in R$, find the most general extension of A in which u and v are units.

The obvious solution. If $A = \langle G \mid R \rangle$, then the solution is

$$\langle G \cup \{x, y\} \mid R \cup \{ux = 1 = vy\} \rangle.$$

$$\begin{aligned} \langle \langle G \mid R \rangle = A. \langle G \cup \{x, y\} \mid R \rangle &\cong A \otimes_{\mathbb{Z}} \mathbb{Z}[x, y] \cong A[x, y]. \\ \langle G \cup \{x, y\} \mid R \cup \{ux = 1 = vy\} \rangle &\cong A[x, y]/(ux - 1, vy - 1). \end{aligned}$$

Elements of $A[x, y]/(ux - 1, vy - 1) = A[u^{-1}, v^{-1}]$ simplify to the form $a(u^{-1})^m(v^{-1})^n$.

Simpler, more direct (and standard) approach.

- 1 Start instead with $S = \{u, v\}^* = \{1, u, v, u^2, uv, \dots\}$, an MC set.
- 2 Define $S^{-1}A$ to consist of ordered pairs $a/s := (a, s)$ modulo equiv. reln.
 $a/s \sim b/t$ iff $(\exists u \in S)(u(sb - ta) = 0)$. (u = “premultiplier”)

Check some things

Check some things

\sim is transitive.

Check some things

\sim is transitive.

If $a/r \sim b/s \sim c/t$,

Check some things

\sim is transitive.

If $a/r \sim b/s \sim c/t$, $\exists u, v \in S$ such that $u(rb - sa) = 0 = v(sc - tb)$.

Check some things

\sim is transitive.

If $a/r \sim b/s \sim c/t$, $\exists u, v \in S$ such that $u(rb - sa) = 0 = v(sc - tb)$. Then $w = uvs \in S$ and

$$w(rc - ta) = uvsr - uvsta = vt(u(rb - sa)) + ur(v(sc - tb)) = 0,$$

Check some things

\sim is transitive.

If $a/r \sim b/s \sim c/t$, $\exists u, v \in S$ such that $u(rb - sa) = 0 = v(sc - tb)$. Then $w = uvs \in S$ and

$$w(rc - ta) = uvsr - uvsta = vt(u(rb - sa)) + ur(v(sc - tb)) = 0,$$

so $a/r \sim c/t$.

Check some things

\sim is transitive.

If $a/r \sim b/s \sim c/t$, $\exists u, v \in S$ such that $u(rb - sa) = 0 = v(sc - tb)$. Then $w = uvs \in S$ and

$$w(rc - ta) = uvsr - uvsta = vt(u(rb - sa)) + ur(v(sc - tb)) = 0,$$

so $a/r \sim c/t$.

premultiplier $u \in S$ is necessary for transitivity of \sim .

Check some things

\sim is transitive.

If $a/r \sim b/s \sim c/t$, $\exists u, v \in S$ such that $u(rb - sa) = 0 = v(sc - tb)$. Then $w = uvs \in S$ and

$$w(rc - ta) = uvsr - uvsta = vt(u(rb - sa)) + ur(v(sc - tb)) = 0,$$

so $a/r \sim c/t$.

premultiplier $u \in S$ is necessary for transitivity of \sim .

E.g., when $0 \in S$,

Check some things

\sim is transitive.

If $a/r \sim b/s \sim c/t$, $\exists u, v \in S$ such that $u(rb - sa) = 0 = v(sc - tb)$. Then $w = uvs \in S$ and

$$w(rc - ta) = uvsr - uvsta = vt(u(rb - sa)) + ur(v(sc - tb)) = 0,$$

so $a/r \sim c/t$.

premultiplier $u \in S$ is necessary for transitivity of \sim .

E.g., when $0 \in S$,

$$0/1 \sim 0/0 \sim 1/0,$$

Check some things

\sim is transitive.

If $a/r \sim b/s \sim c/t$, $\exists u, v \in S$ such that $u(rb - sa) = 0 = v(sc - tb)$. Then $w = uvs \in S$ and

$$w(rc - ta) = uvsr - uvsta = vt(u(rb - sa)) + ur(v(sc - tb)) = 0,$$

so $a/r \sim c/t$.

premultiplier $u \in S$ is necessary for transitivity of \sim .

E.g., when $0 \in S$,

$0/1 \sim 0/0 \sim 1/0$, yet $0/1 \not\sim 1/0$ without the premultiplier.

The universal property

The universal property

Since $S^{-1}A$ is meant to equal $\langle A \cup S^{-1} \mid \{ss^{-1} = 1 \mid s^{-1} \in S^{-1}\} \rangle$,

The universal property

Since $S^{-1}A$ is meant to equal $\langle A \cup S^{-1} \mid \{ss^{-1} = 1 \mid s^{-1} \in S^{-1}\} \rangle$, and since the inverse of a unit is unique,

The universal property

Since $S^{-1}A$ is meant to equal $\langle A \cup S^{-1} \mid \{ss^{-1} = 1 \mid s^{-1} \in S^{-1}\} \rangle$, and since the inverse of a unit is unique, $S^{-1}A$ has the following universal property:

The universal property

Since $S^{-1}A$ is meant to equal $\langle A \cup S^{-1} \mid \{ss^{-1} = 1 \mid s^{-1} \in S^{-1}\} \rangle$, and since the inverse of a unit is unique, $S^{-1}A$ has the following universal property:

Thm. If $\alpha : A \rightarrow B$ is a homomorphism and $\alpha(S) \subseteq B^\times$, then α has a unique extension to $\bar{\alpha} : S^{-1}A \rightarrow B$.

Some notation and terminology

Some notation and terminology

If the elements of S are to become units, then so must any divisor of an element of S .

Some notation and terminology

If the elements of S are to become units, then so must any divisor of an element of S . Call an MC set **saturated** if it is closed under divisors.

Some notation and terminology

If the elements of S are to become units, then so must any divisor of an element of S . Call an MC set **saturated** if it is closed under divisors.

$(ab \in S \Rightarrow a, b \in S.)$

Some notation and terminology

If the elements of S are to become units, then so must any divisor of an element of S . Call an MC set **saturated** if it is closed under divisors.

$(ab \in S \Rightarrow a, b \in S.)$

Thm.

Some notation and terminology

If the elements of S are to become units, then so must any divisor of an element of S . Call an MC set **saturated** if it is closed under divisors.

$(ab \in S \Rightarrow a, b \in S.)$

Thm. A subset $S \subseteq A$ is a saturated MC set iff $A - S$ is a union of prime ideals.

Some notation and terminology

If the elements of S are to become units, then so must any divisor of an element of S . Call an MC set **saturated** if it is closed under divisors.

$(ab \in S \Rightarrow a, b \in S.)$

Thm. A subset $S \subseteq A$ is a saturated MC set iff $A - S$ is a union of prime ideals. (If this union contains one prime \mathfrak{p} , we write $A_{\mathfrak{p}}$ for $S^{-1}A$ and call it the localization of A at \mathfrak{p} .)

Some notation and terminology

If the elements of S are to become units, then so must any divisor of an element of S . Call an MC set **saturated** if it is closed under divisors.

$(ab \in S \Rightarrow a, b \in S.)$

Thm. A subset $S \subseteq A$ is a saturated MC set iff $A - S$ is a union of prime ideals. (If this union contains one prime \mathfrak{p} , we write $A_{\mathfrak{p}}$ for $S^{-1}A$ and call it the localization of A at \mathfrak{p} .)

Proof.

Some notation and terminology

If the elements of S are to become units, then so must any divisor of an element of S . Call an MC set **saturated** if it is closed under divisors.

$(ab \in S \Rightarrow a, b \in S.)$

Thm. A subset $S \subseteq A$ is a saturated MC set iff $A - S$ is a union of prime ideals. (If this union contains one prime \mathfrak{p} , we write $A_{\mathfrak{p}}$ for $S^{-1}A$ and call it the localization of A at \mathfrak{p} .)

Proof.

[If:]

Some notation and terminology

If the elements of S are to become units, then so must any divisor of an element of S . Call an MC set **saturated** if it is closed under divisors.

$(ab \in S \Rightarrow a, b \in S.)$

Thm. A subset $S \subseteq A$ is a saturated MC set iff $A - S$ is a union of prime ideals. (If this union contains one prime \mathfrak{p} , we write $A_{\mathfrak{p}}$ for $S^{-1}A$ and call it the localization of A at \mathfrak{p} .)

Proof.

[If:] The complement of any ideal is saturated.

Some notation and terminology

If the elements of S are to become units, then so must any divisor of an element of S . Call an MC set **saturated** if it is closed under divisors.

$(ab \in S \Rightarrow a, b \in S.)$

Thm. A subset $S \subseteq A$ is a saturated MC set iff $A - S$ is a union of prime ideals. (If this union contains one prime \mathfrak{p} , we write $A_{\mathfrak{p}}$ for $S^{-1}A$ and call it the localization of A at \mathfrak{p} .)

Proof.

[If:] The complement of any ideal is saturated. The complement of a prime ideal is also MC.

Some notation and terminology

If the elements of S are to become units, then so must any divisor of an element of S . Call an MC set **saturated** if it is closed under divisors.

$(ab \in S \Rightarrow a, b \in S.)$

Thm. A subset $S \subseteq A$ is a saturated MC set iff $A - S$ is a union of prime ideals. (If this union contains one prime \mathfrak{p} , we write $A_{\mathfrak{p}}$ for $S^{-1}A$ and call it the localization of A at \mathfrak{p} .)

Proof.

[If:] The complement of any ideal is saturated. The complement of a prime ideal is also MC. The class of SAT+MC subsets of A is closed under arbitrary intersection.

Some notation and terminology

If the elements of S are to become units, then so must any divisor of an element of S . Call an MC set **saturated** if it is closed under divisors.

$(ab \in S \Rightarrow a, b \in S.)$

Thm. A subset $S \subseteq A$ is a saturated MC set iff $A - S$ is a union of prime ideals. (If this union contains one prime \mathfrak{p} , we write $A_{\mathfrak{p}}$ for $S^{-1}A$ and call it the localization of A at \mathfrak{p} .)

Proof.

[If:] The complement of any ideal is saturated. The complement of a prime ideal is also MC. The class of SAT+MC subsets of A is closed under arbitrary intersection. Hence the complement of a union of primes is SAT+MC.

Some notation and terminology

If the elements of S are to become units, then so must any divisor of an element of S . Call an MC set **saturated** if it is closed under divisors.

$(ab \in S \Rightarrow a, b \in S.)$

Thm. A subset $S \subseteq A$ is a saturated MC set iff $A - S$ is a union of prime ideals. (If this union contains one prime \mathfrak{p} , we write $A_{\mathfrak{p}}$ for $S^{-1}A$ and call it the localization of A at \mathfrak{p} .)

Proof.

[If:] The complement of any ideal is saturated. The complement of a prime ideal is also MC. The class of SAT+MC subsets of A is closed under arbitrary intersection. Hence the complement of a union of primes is SAT+MC. \square

Some notation and terminology

If the elements of S are to become units, then so must any divisor of an element of S . Call an MC set **saturated** if it is closed under divisors.

$(ab \in S \Rightarrow a, b \in S.)$

Thm. A subset $S \subseteq A$ is a saturated MC set iff $A - S$ is a union of prime ideals. (If this union contains one prime \mathfrak{p} , we write $A_{\mathfrak{p}}$ for $S^{-1}A$ and call it the localization of A at \mathfrak{p} .)

Proof.

[If:] The complement of any ideal is saturated. The complement of a prime ideal is also MC. The class of SAT+MC subsets of A is closed under arbitrary intersection. Hence the complement of a union of primes is SAT+MC. \square

[Only if:]

Some notation and terminology

If the elements of S are to become units, then so must any divisor of an element of S . Call an MC set **saturated** if it is closed under divisors.

$(ab \in S \Rightarrow a, b \in S.)$

Thm. A subset $S \subseteq A$ is a saturated MC set iff $A - S$ is a union of prime ideals. (If this union contains one prime \mathfrak{p} , we write $A_{\mathfrak{p}}$ for $S^{-1}A$ and call it the localization of A at \mathfrak{p} .)

Proof.

[If:] The complement of any ideal is saturated. The complement of a prime ideal is also MC. The class of SAT+MC subsets of A is closed under arbitrary intersection. Hence the complement of a union of primes is SAT+MC. \square

[Only if:] If S is SAT, then $A - S = \bigcup_{t \notin S} (t)$ is a union of ideals.

Some notation and terminology

If the elements of S are to become units, then so must any divisor of an element of S . Call an MC set **saturated** if it is closed under divisors.

$(ab \in S \Rightarrow a, b \in S.)$

Thm. A subset $S \subseteq A$ is a saturated MC set iff $A - S$ is a union of prime ideals. (If this union contains one prime \mathfrak{p} , we write $A_{\mathfrak{p}}$ for $S^{-1}A$ and call it the localization of A at \mathfrak{p} .)

Proof.

[If:] The complement of any ideal is saturated. The complement of a prime ideal is also MC. The class of SAT+MC subsets of A is closed under arbitrary intersection. Hence the complement of a union of primes is SAT+MC. \square

[Only if:] If S is SAT, then $A - S = \bigcup_{t \notin S} (t)$ is a union of ideals. Since S is MC, any (t) disjoint from S is extendible to a prime \mathfrak{p}_t disjoint from S .

Some notation and terminology

If the elements of S are to become units, then so must any divisor of an element of S . Call an MC set **saturated** if it is closed under divisors.

$(ab \in S \Rightarrow a, b \in S.)$

Thm. A subset $S \subseteq A$ is a saturated MC set iff $A - S$ is a union of prime ideals. (If this union contains one prime \mathfrak{p} , we write $A_{\mathfrak{p}}$ for $S^{-1}A$ and call it the localization of A at \mathfrak{p} .)

Proof.

[If:] The complement of any ideal is saturated. The complement of a prime ideal is also MC. The class of SAT+MC subsets of A is closed under arbitrary intersection. Hence the complement of a union of primes is SAT+MC. \square

[Only if:] If S is SAT, then $A - S = \bigcup_{t \notin S} (t)$ is a union of ideals. Since S is MC, any (t) disjoint from S is extendible to a prime \mathfrak{p}_t disjoint from S .

$A - S = \bigcup_{t \notin S} \mathfrak{p}_t.$

Some notation and terminology

If the elements of S are to become units, then so must any divisor of an element of S . Call an MC set **saturated** if it is closed under divisors.

$(ab \in S \Rightarrow a, b \in S.)$

Thm. A subset $S \subseteq A$ is a saturated MC set iff $A - S$ is a union of prime ideals. (If this union contains one prime \mathfrak{p} , we write $A_{\mathfrak{p}}$ for $S^{-1}A$ and call it the localization of A at \mathfrak{p} .)

Proof.

[If:] The complement of any ideal is saturated. The complement of a prime ideal is also MC. The class of SAT+MC subsets of A is closed under arbitrary intersection. Hence the complement of a union of primes is SAT+MC. \square

[Only if:] If S is SAT, then $A - S = \bigcup_{t \notin S} (t)$ is a union of ideals. Since S is MC, any (t) disjoint from S is extendible to a prime \mathfrak{p}_t disjoint from S .

$A - S = \bigcup_{t \notin S} \mathfrak{p}_t.$ \square

The effect of localization upon ideals

The effect of localization upon ideals

The homomorphism $\alpha : A \rightarrow S^{-1}A : a \mapsto a/1$ allows us to extend and contract ideals from A to $S^{-1}A$:

The effect of localization upon ideals

The homomorphism $\alpha : A \rightarrow S^{-1}A : a \mapsto a/1$ allows us to extend and contract ideals from A to $S^{-1}A$: $I^e = S^{-1}I = \{i/s \mid i \in I, s \in S\}$,

The effect of localization upon ideals

The homomorphism $\alpha : A \rightarrow S^{-1}A : a \mapsto a/1$ allows us to extend and contract ideals from A to $S^{-1}A$: $I^e = S^{-1}I = \{i/s \mid i \in I, s \in S\}$, $J^c = \alpha^{-1}(J)$.

The effect of localization upon ideals

The homomorphism $\alpha : A \rightarrow S^{-1}A : a \mapsto a/1$ allows us to extend and contract ideals from A to $S^{-1}A$: $I^e = S^{-1}I = \{i/s \mid i \in I, s \in S\}$, $J^c = \alpha^{-1}(J)$.

Thm.

The effect of localization upon ideals

The homomorphism $\alpha : A \rightarrow S^{-1}A : a \mapsto a/1$ allows us to extend and contract ideals from A to $S^{-1}A$: $I^e = S^{-1}I = \{i/s \mid i \in I, s \in S\}$, $J^c = \alpha^{-1}(J)$.

Thm. (See AM Prop. 3.11)

The effect of localization upon ideals

The homomorphism $\alpha : A \rightarrow S^{-1}A : a \mapsto a/1$ allows us to extend and contract ideals from A to $S^{-1}A$: $I^e = S^{-1}I = \{i/s \mid i \in I, s \in S\}$, $J^c = \alpha^{-1}(J)$.

Thm. (See AM Prop. 3.11)

- 1 Every ideal of $S^{-1}A$ is an extended ideal.

The effect of localization upon ideals

The homomorphism $\alpha : A \rightarrow S^{-1}A : a \mapsto a/1$ allows us to extend and contract ideals from A to $S^{-1}A$: $I^e = S^{-1}I = \{i/s \mid i \in I, s \in S\}$, $J^c = \alpha^{-1}(J)$.

Thm. (See AM Prop. 3.11)

- 1 Every ideal of $S^{-1}A$ is an extended ideal.

The effect of localization upon ideals

The homomorphism $\alpha : A \rightarrow S^{-1}A : a \mapsto a/1$ allows us to extend and contract ideals from A to $S^{-1}A$: $I^e = S^{-1}I = \{i/s \mid i \in I, s \in S\}$, $J^c = \alpha^{-1}(J)$.

Thm. (See AM Prop. 3.11)

- ① Every ideal of $S^{-1}A$ is an extended ideal. ($J = J^{ce}$.)
- ② $I^{ec} = \bigcup_{s \in S} (I : s)$

The effect of localization upon ideals

The homomorphism $\alpha : A \rightarrow S^{-1}A : a \mapsto a/1$ allows us to extend and contract ideals from A to $S^{-1}A$: $I^e = S^{-1}I = \{i/s \mid i \in I, s \in S\}$, $J^c = \alpha^{-1}(J)$.

Thm. (See AM Prop. 3.11)

- ① Every ideal of $S^{-1}A$ is an extended ideal. ($J = J^{ce}$.)
- ② $I^{ec} = \bigcup_{s \in S} (I : s)$

The effect of localization upon ideals

The homomorphism $\alpha : A \rightarrow S^{-1}A : a \mapsto a/1$ allows us to extend and contract ideals from A to $S^{-1}A$: $I^e = S^{-1}I = \{i/s \mid i \in I, s \in S\}$, $J^c = \alpha^{-1}(J)$.

Thm. (See AM Prop. 3.11)

- ① Every ideal of $S^{-1}A$ is an extended ideal. ($J = J^{ce}$.)
- ② $I^{ec} = \bigcup_{s \in S} (I : s) = \text{sat}(I)$.

The effect of localization upon ideals

The homomorphism $\alpha : A \rightarrow S^{-1}A : a \mapsto a/1$ allows us to extend and contract ideals from A to $S^{-1}A$: $I^e = S^{-1}I = \{i/s \mid i \in I, s \in S\}$, $J^c = \alpha^{-1}(J)$.

Thm. (See AM Prop. 3.11)

- ① Every ideal of $S^{-1}A$ is an extended ideal. ($J = J^{ce}$.)
- ② $I^{ec} = \bigcup_{s \in S} (I : s) = \text{sat}(I)$. ($I = \text{sat}(I)$ iff $\lambda_s : A/I \rightarrow A/I$ is 1-1 $\forall s$.)

The effect of localization upon ideals

The homomorphism $\alpha : A \rightarrow S^{-1}A : a \mapsto a/1$ allows us to extend and contract ideals from A to $S^{-1}A$: $I^e = S^{-1}I = \{i/s \mid i \in I, s \in S\}$, $J^c = \alpha^{-1}(J)$.

Thm. (See AM Prop. 3.11)

- ① Every ideal of $S^{-1}A$ is an extended ideal. ($J = J^{ce}$.)
- ② $I^{ec} = \bigcup_{s \in S} (I : s) = \text{sat}(I)$. ($I = \text{sat}(I)$ iff $\lambda_s : A/I \rightarrow A/I$ is 1-1 $\forall s$.)
- ③ Extension and contraction restrict to inverse bijections between the primes of $S^{-1}A$ and the primes of A disjoint from S .

The effect of localization upon ideals

The homomorphism $\alpha : A \rightarrow S^{-1}A : a \mapsto a/1$ allows us to extend and contract ideals from A to $S^{-1}A$: $I^e = S^{-1}I = \{i/s \mid i \in I, s \in S\}$, $J^c = \alpha^{-1}(J)$.

Thm. (See AM Prop. 3.11)

- ① Every ideal of $S^{-1}A$ is an extended ideal. ($J = J^{ce}$.)
- ② $I^{ec} = \bigcup_{s \in S} (I : s) = \text{sat}(I)$. ($I = \text{sat}(I)$ iff $\lambda_s : A/I \rightarrow A/I$ is 1-1 $\forall s$.)
- ③ Extension and contraction restrict to inverse bijections between the primes of $S^{-1}A$ and the primes of A disjoint from S .

The effect of localization upon ideals

The homomorphism $\alpha : A \rightarrow S^{-1}A : a \mapsto a/1$ allows us to extend and contract ideals from A to $S^{-1}A$: $I^e = S^{-1}I = \{i/s \mid i \in I, s \in S\}$, $J^c = \alpha^{-1}(J)$.

Thm. (See AM Prop. 3.11)

- ① Every ideal of $S^{-1}A$ is an extended ideal. ($J = J^{ce}$.)
- ② $I^{ec} = \bigcup_{s \in S} (I : s) = \text{sat}(I)$. ($I = \text{sat}(I)$ iff $\lambda_s : A/I \rightarrow A/I$ is 1-1 $\forall s$.)
- ③ Extension and contraction restrict to inverse bijections between the primes of $S^{-1}A$ and the primes of A disjoint from S . ($\text{sat}(\mathfrak{p}) = \mathfrak{p}$ or A .)

The effect of localization upon ideals

The homomorphism $\alpha : A \rightarrow S^{-1}A : a \mapsto a/1$ allows us to extend and contract ideals from A to $S^{-1}A$: $I^e = S^{-1}I = \{i/s \mid i \in I, s \in S\}$, $J^c = \alpha^{-1}(J)$.

Thm. (See AM Prop. 3.11)

- ① Every ideal of $S^{-1}A$ is an extended ideal. ($J = J^{ce}$.)
- ② $I^{ec} = \bigcup_{s \in S} (I : s) = \text{sat}(I)$. ($I = \text{sat}(I)$ iff $\lambda_s : A/I \rightarrow A/I$ is 1-1 $\forall s$.)
- ③ Extension and contraction restrict to inverse bijections between the primes of $S^{-1}A$ and the primes of A disjoint from S . ($\text{sat}(\mathfrak{p}) = \mathfrak{p}$ or A .)
- ④ $S^{-1}(\sum I_j) = \sum S^{-1}I_j$, $S^{-1}(I \cap J) = S^{-1}I \cap S^{-1}J$
 $S^{-1}(IJ) = (S^{-1}I)(S^{-1}J)$, $S^{-1}\sqrt{I} = \sqrt{S^{-1}I}$.

The effect of localization upon ideals

The homomorphism $\alpha : A \rightarrow S^{-1}A : a \mapsto a/1$ allows us to extend and contract ideals from A to $S^{-1}A$: $I^e = S^{-1}I = \{i/s \mid i \in I, s \in S\}$, $J^c = \alpha^{-1}(J)$.

Thm. (See AM Prop. 3.11)

- ① Every ideal of $S^{-1}A$ is an extended ideal. ($J = J^{ce}$.)
- ② $I^{ec} = \bigcup_{s \in S} (I : s) = \text{sat}(I)$. ($I = \text{sat}(I)$ iff $\lambda_s : A/I \rightarrow A/I$ is 1-1 $\forall s$.)
- ③ Extension and contraction restrict to inverse bijections between the primes of $S^{-1}A$ and the primes of A disjoint from S . ($\text{sat}(\mathfrak{p}) = \mathfrak{p}$ or A .)
- ④ $S^{-1}(\sum I_j) = \sum S^{-1}I_j$, $S^{-1}(I \cap J) = S^{-1}I \cap S^{-1}J$
 $S^{-1}(IJ) = (S^{-1}I)(S^{-1}J)$, $S^{-1}\sqrt{I} = \sqrt{S^{-1}I}$.

The effect of localization upon ideals

The homomorphism $\alpha : A \rightarrow S^{-1}A : a \mapsto a/1$ allows us to extend and contract ideals from A to $S^{-1}A$: $I^e = S^{-1}I = \{i/s \mid i \in I, s \in S\}$, $J^c = \alpha^{-1}(J)$.

Thm. (See AM Prop. 3.11)

- ① Every ideal of $S^{-1}A$ is an extended ideal. ($J = J^{ce}$.)
- ② $I^{ec} = \bigcup_{s \in S} (I : s) = \text{sat}(I)$. ($I = \text{sat}(I)$ iff $\lambda_s : A/I \rightarrow A/I$ is 1-1 $\forall s$.)
- ③ Extension and contraction restrict to inverse bijections between the primes of $S^{-1}A$ and the primes of A disjoint from S . ($\text{sat}(\mathfrak{p}) = \mathfrak{p}$ or A .)
- ④ $S^{-1}(\sum I_j) = \sum S^{-1}I_j$, $S^{-1}(I \cap J) = S^{-1}I \cap S^{-1}J$
 $S^{-1}(IJ) = (S^{-1}I)(S^{-1}J)$, $S^{-1}\sqrt{I} = \sqrt{S^{-1}I}$.
- ⑤ $S^{-1}J(I)$ need not equal $J(S^{-1}I)$.

The effect of localization upon ideals

The homomorphism $\alpha : A \rightarrow S^{-1}A : a \mapsto a/1$ allows us to extend and contract ideals from A to $S^{-1}A$: $I^e = S^{-1}I = \{i/s \mid i \in I, s \in S\}$, $J^c = \alpha^{-1}(J)$.

Thm. (See AM Prop. 3.11)

- ① Every ideal of $S^{-1}A$ is an extended ideal. ($J = J^{ce}$.)
- ② $I^{ec} = \bigcup_{s \in S} (I : s) = \text{sat}(I)$. ($I = \text{sat}(I)$ iff $\lambda_s : A/I \rightarrow A/I$ is 1-1 $\forall s$.)
- ③ Extension and contraction restrict to inverse bijections between the primes of $S^{-1}A$ and the primes of A disjoint from S . ($\text{sat}(\mathfrak{p}) = \mathfrak{p}$ or A .)
- ④ $S^{-1}(\sum I_j) = \sum S^{-1}I_j$, $S^{-1}(I \cap J) = S^{-1}I \cap S^{-1}J$
 $S^{-1}(IJ) = (S^{-1}I)(S^{-1}J)$, $S^{-1}\sqrt{I} = \sqrt{S^{-1}I}$.
- ⑤ $S^{-1}J(I)$ need not equal $J(S^{-1}I)$.

The effect of localization upon ideals

The homomorphism $\alpha : A \rightarrow S^{-1}A : a \mapsto a/1$ allows us to extend and contract ideals from A to $S^{-1}A$: $I^e = S^{-1}I = \{i/s \mid i \in I, s \in S\}$, $J^c = \alpha^{-1}(J)$.

Thm. (See AM Prop. 3.11)

- 1 Every ideal of $S^{-1}A$ is an extended ideal. ($J = J^{ce}$.)
- 2 $I^{ec} = \bigcup_{s \in S} (I : s) = \text{sat}(I)$. ($I = \text{sat}(I)$ iff $\lambda_s : A/I \rightarrow A/I$ is 1-1 $\forall s$.)
- 3 Extension and contraction restrict to inverse bijections between the primes of $S^{-1}A$ and the primes of A disjoint from S . ($\text{sat}(\mathfrak{p}) = \mathfrak{p}$ or A .)
- 4 $S^{-1}(\sum I_j) = \sum S^{-1}I_j$, $S^{-1}(I \cap J) = S^{-1}I \cap S^{-1}J$
 $S^{-1}(IJ) = (S^{-1}I)(S^{-1}J)$, $S^{-1}\sqrt{I} = \sqrt{S^{-1}I}$.
- 5 $S^{-1}J(I)$ need not equal $J(S^{-1}I)$. $A = I = \mathbb{Z}$, $S = \{3, 5, 7, \dots\}^*$.

The effect of localization upon ideals

The homomorphism $\alpha : A \rightarrow S^{-1}A : a \mapsto a/1$ allows us to extend and contract ideals from A to $S^{-1}A$: $I^e = S^{-1}I = \{i/s \mid i \in I, s \in S\}$, $J^c = \alpha^{-1}(J)$.

Thm. (See AM Prop. 3.11)

- ① Every ideal of $S^{-1}A$ is an extended ideal. ($J = J^{ce}$.)
- ② $I^{ec} = \bigcup_{s \in S} (I : s) = \text{sat}(I)$. ($I = \text{sat}(I)$ iff $\lambda_s : A/I \rightarrow A/I$ is 1-1 $\forall s$.)
- ③ Extension and contraction restrict to inverse bijections between the primes of $S^{-1}A$ and the primes of A disjoint from S . ($\text{sat}(\mathfrak{p}) = \mathfrak{p}$ or A .)
- ④ $S^{-1}(\sum I_j) = \sum S^{-1}I_j$, $S^{-1}(I \cap J) = S^{-1}I \cap S^{-1}J$
 $S^{-1}(IJ) = (S^{-1}I)(S^{-1}J)$, $S^{-1}\sqrt{I} = \sqrt{S^{-1}I}$.
- ⑤ $S^{-1}J(I)$ need not equal $J(S^{-1}I)$. $A = I = \mathbb{Z}$, $S = \{3, 5, 7, \dots\}^*$.

(2)

The effect of localization upon ideals

The homomorphism $\alpha : A \rightarrow S^{-1}A : a \mapsto a/1$ allows us to extend and contract ideals from A to $S^{-1}A$: $I^e = S^{-1}I = \{i/s \mid i \in I, s \in S\}$, $J^c = \alpha^{-1}(J)$.

Thm. (See AM Prop. 3.11)

- ① Every ideal of $S^{-1}A$ is an extended ideal. ($J = J^{ce}$.)
- ② $I^{ec} = \bigcup_{s \in S} (I : s) = \text{sat}(I)$. ($I = \text{sat}(I)$ iff $\lambda_s : A/I \rightarrow A/I$ is 1-1 $\forall s$.)
- ③ Extension and contraction restrict to inverse bijections between the primes of $S^{-1}A$ and the primes of A disjoint from S . ($\text{sat}(\mathfrak{p}) = \mathfrak{p}$ or A .)
- ④ $S^{-1}(\sum I_j) = \sum S^{-1}I_j$, $S^{-1}(I \cap J) = S^{-1}I \cap S^{-1}J$
 $S^{-1}(IJ) = (S^{-1}I)(S^{-1}J)$, $S^{-1}\sqrt{I} = \sqrt{S^{-1}I}$.
- ⑤ $S^{-1}J(I)$ need not equal $J(S^{-1}I)$. $A = I = \mathbb{Z}$, $S = \{3, 5, 7, \dots\}^*$.

(2) $x \in I^{ec}$

The effect of localization upon ideals

The homomorphism $\alpha : A \rightarrow S^{-1}A : a \mapsto a/1$ allows us to extend and contract ideals from A to $S^{-1}A$: $I^e = S^{-1}I = \{i/s \mid i \in I, s \in S\}$, $J^c = \alpha^{-1}(J)$.

Thm. (See AM Prop. 3.11)

- ① Every ideal of $S^{-1}A$ is an extended ideal. ($J = J^{ce}$.)
- ② $I^{ec} = \bigcup_{s \in S} (I : s) = \text{sat}(I)$. ($I = \text{sat}(I)$ iff $\lambda_s : A/I \rightarrow A/I$ is 1-1 $\forall s$.)
- ③ Extension and contraction restrict to inverse bijections between the primes of $S^{-1}A$ and the primes of A disjoint from S . ($\text{sat}(\mathfrak{p}) = \mathfrak{p}$ or A .)
- ④ $S^{-1}(\sum I_j) = \sum S^{-1}I_j$, $S^{-1}(I \cap J) = S^{-1}I \cap S^{-1}J$
 $S^{-1}(IJ) = (S^{-1}I)(S^{-1}J)$, $S^{-1}\sqrt{I} = \sqrt{S^{-1}I}$.
- ⑤ $S^{-1}J(I)$ need not equal $J(S^{-1}I)$. $A = I = \mathbb{Z}$, $S = \{3, 5, 7, \dots\}^*$.

(2) $x \in I^{ec}$ iff $x/1 \sim i/t \in I^e$

The effect of localization upon ideals

The homomorphism $\alpha : A \rightarrow S^{-1}A : a \mapsto a/1$ allows us to extend and contract ideals from A to $S^{-1}A$: $I^e = S^{-1}I = \{i/s \mid i \in I, s \in S\}$, $J^c = \alpha^{-1}(J)$.

Thm. (See AM Prop. 3.11)

- ① Every ideal of $S^{-1}A$ is an extended ideal. ($J = J^{ce}$.)
- ② $I^{ec} = \bigcup_{s \in S} (I : s) = \text{sat}(I)$. ($I = \text{sat}(I)$ iff $\lambda_s : A/I \rightarrow A/I$ is 1-1 $\forall s$.)
- ③ Extension and contraction restrict to inverse bijections between the primes of $S^{-1}A$ and the primes of A disjoint from S . ($\text{sat}(\mathfrak{p}) = \mathfrak{p}$ or A .)
- ④ $S^{-1}(\sum I_j) = \sum S^{-1}I_j$, $S^{-1}(I \cap J) = S^{-1}I \cap S^{-1}J$
 $S^{-1}(IJ) = (S^{-1}I)(S^{-1}J)$, $S^{-1}\sqrt{I} = \sqrt{S^{-1}I}$.
- ⑤ $S^{-1}J(I)$ need not equal $J(S^{-1}I)$. $A = I = \mathbb{Z}$, $S = \{3, 5, 7, \dots\}^*$.

(2) $x \in I^{ec}$ iff $x/1 \sim i/t \in I^e$ iff $(\exists u)(u(i - tx) = 0)$

The effect of localization upon ideals

The homomorphism $\alpha : A \rightarrow S^{-1}A : a \mapsto a/1$ allows us to extend and contract ideals from A to $S^{-1}A$: $I^e = S^{-1}I = \{i/s \mid i \in I, s \in S\}$, $J^c = \alpha^{-1}(J)$.

Thm. (See AM Prop. 3.11)

- ① Every ideal of $S^{-1}A$ is an extended ideal. ($J = J^{ce}$.)
- ② $I^{ec} = \bigcup_{s \in S} (I : s) = \text{sat}(I)$. ($I = \text{sat}(I)$ iff $\lambda_s : A/I \rightarrow A/I$ is 1-1 $\forall s$.)
- ③ Extension and contraction restrict to inverse bijections between the primes of $S^{-1}A$ and the primes of A disjoint from S . ($\text{sat}(\mathfrak{p}) = \mathfrak{p}$ or A .)
- ④ $S^{-1}(\sum I_j) = \sum S^{-1}I_j$, $S^{-1}(I \cap J) = S^{-1}I \cap S^{-1}J$
 $S^{-1}(IJ) = (S^{-1}I)(S^{-1}J)$, $S^{-1}\sqrt{I} = \sqrt{S^{-1}I}$.
- ⑤ $S^{-1}J(I)$ need not equal $J(S^{-1}I)$. $A = I = \mathbb{Z}$, $S = \{3, 5, 7, \dots\}^*$.

(2) $x \in I^{ec}$ iff $x/1 \sim i/t \in I^e$ iff $(\exists u)(u(i - tx) = 0)$ iff $(\exists u)(ui = utx)$

The effect of localization upon ideals

The homomorphism $\alpha : A \rightarrow S^{-1}A : a \mapsto a/1$ allows us to extend and contract ideals from A to $S^{-1}A$: $I^e = S^{-1}I = \{i/s \mid i \in I, s \in S\}$, $J^c = \alpha^{-1}(J)$.

Thm. (See AM Prop. 3.11)

- ① Every ideal of $S^{-1}A$ is an extended ideal. ($J = J^{ce}$.)
- ② $I^{ec} = \bigcup_{s \in S} (I : s) = \text{sat}(I)$. ($I = \text{sat}(I)$ iff $\lambda_s : A/I \rightarrow A/I$ is 1-1 $\forall s$.)
- ③ Extension and contraction restrict to inverse bijections between the primes of $S^{-1}A$ and the primes of A disjoint from S . ($\text{sat}(\mathfrak{p}) = \mathfrak{p}$ or A .)
- ④ $S^{-1}(\sum I_j) = \sum S^{-1}I_j$, $S^{-1}(I \cap J) = S^{-1}I \cap S^{-1}J$
 $S^{-1}(IJ) = (S^{-1}I)(S^{-1}J)$, $S^{-1}\sqrt{I} = \sqrt{S^{-1}I}$.
- ⑤ $S^{-1}J(I)$ need not equal $J(S^{-1}I)$. $A = I = \mathbb{Z}$, $S = \{3, 5, 7, \dots\}^*$.

(2) $x \in I^{ec}$ iff $x/1 \sim i/t \in I^e$ iff $(\exists u)(u(i - tx) = 0)$ iff $(\exists u)(ui = utx)$ implies $x \in \bigcup_{s \in S} (I : s)$

The effect of localization upon ideals

The homomorphism $\alpha : A \rightarrow S^{-1}A : a \mapsto a/1$ allows us to extend and contract ideals from A to $S^{-1}A$: $I^e = S^{-1}I = \{i/s \mid i \in I, s \in S\}$, $J^c = \alpha^{-1}(J)$.

Thm. (See AM Prop. 3.11)

- ① Every ideal of $S^{-1}A$ is an extended ideal. ($J = J^{ce}$.)
- ② $I^{ec} = \bigcup_{s \in S} (I : s) = \text{sat}(I)$. ($I = \text{sat}(I)$ iff $\lambda_s : A/I \rightarrow A/I$ is 1-1 $\forall s$.)
- ③ Extension and contraction restrict to inverse bijections between the primes of $S^{-1}A$ and the primes of A disjoint from S . ($\text{sat}(\mathfrak{p}) = \mathfrak{p}$ or A .)
- ④ $S^{-1}(\sum I_j) = \sum S^{-1}I_j$, $S^{-1}(I \cap J) = S^{-1}I \cap S^{-1}J$
 $S^{-1}(IJ) = (S^{-1}I)(S^{-1}J)$, $S^{-1}\sqrt{I} = \sqrt{S^{-1}I}$.
- ⑤ $S^{-1}J(I)$ need not equal $J(S^{-1}I)$. $A = I = \mathbb{Z}$, $S = \{3, 5, 7, \dots\}^*$.

(2) $x \in I^{ec}$ iff $x/1 \sim i/t \in I^e$ iff $(\exists u)(u(i - tx) = 0)$ iff $(\exists u)(ui = utx)$ implies $x \in \bigcup_{s \in S} (I : s)$ implies $(\exists s)(xs = j \in I)$

The effect of localization upon ideals

The homomorphism $\alpha : A \rightarrow S^{-1}A : a \mapsto a/1$ allows us to extend and contract ideals from A to $S^{-1}A$: $I^e = S^{-1}I = \{i/s \mid i \in I, s \in S\}$, $J^c = \alpha^{-1}(J)$.

Thm. (See AM Prop. 3.11)

- ① Every ideal of $S^{-1}A$ is an extended ideal. ($J = J^{ce}$.)
- ② $I^{ec} = \bigcup_{s \in S} (I : s) = \text{sat}(I)$. ($I = \text{sat}(I)$ iff $\lambda_s : A/I \rightarrow A/I$ is 1-1 $\forall s$.)
- ③ Extension and contraction restrict to inverse bijections between the primes of $S^{-1}A$ and the primes of A disjoint from S . ($\text{sat}(\mathfrak{p}) = \mathfrak{p}$ or A .)
- ④ $S^{-1}(\sum I_j) = \sum S^{-1}I_j$, $S^{-1}(I \cap J) = S^{-1}I \cap S^{-1}J$
 $S^{-1}(IJ) = (S^{-1}I)(S^{-1}J)$, $S^{-1}\sqrt{I} = \sqrt{S^{-1}I}$.
- ⑤ $S^{-1}J(I)$ need not equal $J(S^{-1}I)$. $A = I = \mathbb{Z}$, $S = \{3, 5, 7, \dots\}^*$.

(2) $x \in I^{ec}$ iff $x/1 \sim i/t \in I^e$ iff $(\exists u)(u(i - tx) = 0)$ iff $(\exists u)(ui = utx)$ implies $x \in \bigcup_{s \in S} (I : s)$ implies $(\exists s)(xs = j \in I)$ implies $x/1 \sim j/s \in I^e$.

The effect of localization upon ideals

The homomorphism $\alpha : A \rightarrow S^{-1}A : a \mapsto a/1$ allows us to extend and contract ideals from A to $S^{-1}A$: $I^e = S^{-1}I = \{i/s \mid i \in I, s \in S\}$, $J^c = \alpha^{-1}(J)$.

Thm. (See AM Prop. 3.11)

- ① Every ideal of $S^{-1}A$ is an extended ideal. ($J = J^{ce}$.)
- ② $I^{ec} = \bigcup_{s \in S} (I : s) = \text{sat}(I)$. ($I = \text{sat}(I)$ iff $\lambda_s : A/I \rightarrow A/I$ is 1-1 $\forall s$.)
- ③ Extension and contraction restrict to inverse bijections between the primes of $S^{-1}A$ and the primes of A disjoint from S . ($\text{sat}(\mathfrak{p}) = \mathfrak{p}$ or A .)
- ④ $S^{-1}(\sum I_j) = \sum S^{-1}I_j$, $S^{-1}(I \cap J) = S^{-1}I \cap S^{-1}J$
 $S^{-1}(IJ) = (S^{-1}I)(S^{-1}J)$, $S^{-1}\sqrt{I} = \sqrt{S^{-1}I}$.
- ⑤ $S^{-1}J(I)$ need not equal $J(S^{-1}I)$. $A = I = \mathbb{Z}$, $S = \{3, 5, 7, \dots\}^*$.

(2) $x \in I^{ec}$ iff $x/1 \sim i/t \in I^e$ iff $(\exists u)(u(i - tx) = 0)$ iff $(\exists u)(ui = utx)$ implies $x \in \bigcup_{s \in S} (I : s)$ implies $(\exists s)(xs = j \in I)$ implies $x/1 \sim j/s \in I^e$. \square

Restriction and extension of scalars

Restriction and extension of scalars

The homomorphism $A \rightarrow S^{-1}A : a \mapsto a/1$ induces functors of restriction and extension of scalars between $A\text{-Mod}$ and $S^{-1}A\text{-Mod}$.

Restriction and extension of scalars

The homomorphism $A \rightarrow S^{-1}A : a \mapsto a/1$ induces functors of restriction and extension of scalars between $A\text{-Mod}$ and $S^{-1}A\text{-Mod}$. The functor of extension of scalars takes M to $S^{-1}A \otimes_A M \cong S^{-1}M$, where $S^{-1}M$ is constructed in the expected way.

Restriction and extension of scalars

The homomorphism $A \rightarrow S^{-1}A : a \mapsto a/1$ induces functors of restriction and extension of scalars between $A\text{-Mod}$ and $S^{-1}A\text{-Mod}$. The functor of extension of scalars takes M to $S^{-1}A \otimes_A M \cong S^{-1}M$, where $S^{-1}M$ is constructed in the expected way.

($S^{-1}M$ consists of fractions m/s modulo \sim .)

Restriction and extension of scalars

The homomorphism $A \rightarrow S^{-1}A : a \mapsto a/1$ induces functors of restriction and extension of scalars between $A\text{-Mod}$ and $S^{-1}A\text{-Mod}$. The functor of extension of scalars takes M to $S^{-1}A \otimes_A M \cong S^{-1}M$, where $S^{-1}M$ is constructed in the expected way.

($S^{-1}M$ consists of fractions m/s modulo \sim .)

(Isomorphism $S^{-1}A \otimes_A M \rightarrow S^{-1}M$ is induced by $(a/s) \otimes m \mapsto am/s$.)

Localization is an exact additive functor

Localization is an exact additive functor

Thm. If $L \xrightarrow{\alpha} M \xrightarrow{\beta} N$ is exact, then

$$S^{-1}L \xrightarrow{S^{-1}\alpha} S^{-1}M \xrightarrow{S^{-1}\beta} S^{-1}N \quad (\dagger)$$

is exact for any MC S .

Localization is an exact additive functor

Thm. If $L \xrightarrow{\alpha} M \xrightarrow{\beta} N$ is exact, then

$$S^{-1}L \xrightarrow{S^{-1}\alpha} S^{-1}M \xrightarrow{S^{-1}\beta} S^{-1}N \quad (\dagger)$$

is exact for any MC S .

Proof.

Localization is an exact additive functor

Thm. If $L \xrightarrow{\alpha} M \xrightarrow{\beta} N$ is exact, then

$$S^{-1}L \xrightarrow{S^{-1}\alpha} S^{-1}M \xrightarrow{S^{-1}\beta} S^{-1}N \quad (\dagger)$$

is exact for any MC S .

Proof.

Localization is an additive functor, so (\dagger) is a chain complex.

Localization is an exact additive functor

Thm. If $L \xrightarrow{\alpha} M \xrightarrow{\beta} N$ is exact, then

$$S^{-1}L \xrightarrow{S^{-1}\alpha} S^{-1}M \xrightarrow{S^{-1}\beta} S^{-1}N \quad (\dagger)$$

is exact for any MC S .

Proof.

Localization is an additive functor, so (\dagger) is a chain complex. Hence only need to prove $\ker(S^{-1}\beta) \subseteq \operatorname{im}(S^{-1}\alpha)$.

Localization is an exact additive functor

Thm. If $L \xrightarrow{\alpha} M \xrightarrow{\beta} N$ is exact, then

$$S^{-1}L \xrightarrow{S^{-1}\alpha} S^{-1}M \xrightarrow{S^{-1}\beta} S^{-1}N \quad (\dagger)$$

is exact for any MC S .

Proof.

Localization is an additive functor, so (\dagger) is a chain complex. Hence only need to prove $\ker(S^{-1}\beta) \subseteq \operatorname{im}(S^{-1}\alpha)$.

If $m/s \in \ker(S^{-1}\beta)$, then $\beta(m)/s = 0/1$ in $S^{-1}N$.

Localization is an exact additive functor

Thm. If $L \xrightarrow{\alpha} M \xrightarrow{\beta} N$ is exact, then

$$S^{-1}L \xrightarrow{S^{-1}\alpha} S^{-1}M \xrightarrow{S^{-1}\beta} S^{-1}N \quad (\dagger)$$

is exact for any MC S .

Proof.

Localization is an additive functor, so (\dagger) is a chain complex. Hence only need to prove $\ker(S^{-1}\beta) \subseteq \operatorname{im}(S^{-1}\alpha)$.

If $m/s \in \ker(S^{-1}\beta)$, then $\beta(m)/s = 0/1$ in $S^{-1}N$. Hence $(\exists u)(u(\beta(m) - 0) = 0)$,

Localization is an exact additive functor

Thm. If $L \xrightarrow{\alpha} M \xrightarrow{\beta} N$ is exact, then

$$S^{-1}L \xrightarrow{S^{-1}\alpha} S^{-1}M \xrightarrow{S^{-1}\beta} S^{-1}N \quad (\dagger)$$

is exact for any MC S .

Proof.

Localization is an additive functor, so (\dagger) is a chain complex. Hence only need to prove $\ker(S^{-1}\beta) \subseteq \operatorname{im}(S^{-1}\alpha)$.

If $m/s \in \ker(S^{-1}\beta)$, then $\beta(m)/s = 0/1$ in $S^{-1}N$. Hence $(\exists u)(u(\beta(m) - 0) = 0)$, or $um \in \ker(\beta) = \operatorname{im}(\alpha)$.

Localization is an exact additive functor

Thm. If $L \xrightarrow{\alpha} M \xrightarrow{\beta} N$ is exact, then

$$S^{-1}L \xrightarrow{S^{-1}\alpha} S^{-1}M \xrightarrow{S^{-1}\beta} S^{-1}N \quad (\dagger)$$

is exact for any MC S .

Proof.

Localization is an additive functor, so (\dagger) is a chain complex. Hence only need to prove $\ker(S^{-1}\beta) \subseteq \operatorname{im}(S^{-1}\alpha)$.

If $m/s \in \ker(S^{-1}\beta)$, then $\beta(m)/s = 0/1$ in $S^{-1}N$. Hence

$(\exists u)(u(\beta(m) - 0) = 0)$, or $um \in \ker(\beta) = \operatorname{im}(\alpha)$. Hence $\alpha(\ell) = um$ for some $\ell \in L$.

Localization is an exact additive functor

Thm. If $L \xrightarrow{\alpha} M \xrightarrow{\beta} N$ is exact, then

$$S^{-1}L \xrightarrow{S^{-1}\alpha} S^{-1}M \xrightarrow{S^{-1}\beta} S^{-1}N \quad (\dagger)$$

is exact for any MC S .

Proof.

Localization is an additive functor, so (\dagger) is a chain complex. Hence only need to prove $\ker(S^{-1}\beta) \subseteq \operatorname{im}(S^{-1}\alpha)$.

If $m/s \in \ker(S^{-1}\beta)$, then $\beta(m)/s = 0/1$ in $S^{-1}N$. Hence $(\exists u)(u(\beta(m) - 0) = 0)$, or $um \in \ker(\beta) = \operatorname{im}(\alpha)$. Hence $\alpha(\ell) = um$ for some $\ell \in L$. Hence $m/s = \alpha(\ell/(us)) \in \operatorname{im}(S^{-1}\alpha)$.

Localization is an exact additive functor

Thm. If $L \xrightarrow{\alpha} M \xrightarrow{\beta} N$ is exact, then

$$S^{-1}L \xrightarrow{S^{-1}\alpha} S^{-1}M \xrightarrow{S^{-1}\beta} S^{-1}N \quad (\dagger)$$

is exact for any MC S .

Proof.

Localization is an additive functor, so (\dagger) is a chain complex. Hence only need to prove $\ker(S^{-1}\beta) \subseteq \operatorname{im}(S^{-1}\alpha)$.

If $m/s \in \ker(S^{-1}\beta)$, then $\beta(m)/s = 0/1$ in $S^{-1}N$. Hence $(\exists u)(u(\beta(m) - 0) = 0)$, or $um \in \ker(\beta) = \operatorname{im}(\alpha)$. Hence $\alpha(\ell) = um$ for some $\ell \in L$. Hence $m/s = \alpha(\ell/(us)) \in \operatorname{im}(S^{-1}\alpha)$. \square

Local properties

A property \mathcal{P} of a ring or module is local if “ A (or M) has \mathcal{P} ” iff “ $A_{\mathfrak{p}}$ (or $M_{\mathfrak{p}}$) has \mathcal{P} for all prime ideals \mathfrak{p} ”.

Local properties

A property \mathcal{P} of a ring or module is local if “ A (or M) has \mathcal{P} ” iff “ $A_{\mathfrak{p}}$ (or $M_{\mathfrak{p}}$) has \mathcal{P} for all prime ideals \mathfrak{p} ”. (\mathcal{P} is preserved and reflected by localization to all primes.)

Local properties

A property \mathcal{P} of a ring or module is local if “ A (or M) has \mathcal{P} ” iff “ $A_{\mathfrak{p}}$ (or $M_{\mathfrak{p}}$) has \mathcal{P} for all prime ideals \mathfrak{p} ”. (\mathcal{P} is preserved and reflected by localization to all primes.)

Examples.

Local properties

A property \mathcal{P} of a ring or module is local if “ A (or M) has \mathcal{P} ” iff “ $A_{\mathfrak{p}}$ (or $M_{\mathfrak{p}}$) has \mathcal{P} for all prime ideals \mathfrak{p} ”. (\mathcal{P} is preserved and reflected by localization to all primes.)

Examples.

- 1 $M = 0$.

Local properties

A property \mathcal{P} of a ring or module is local if “ A (or M) has \mathcal{P} ” iff “ $A_{\mathfrak{p}}$ (or $M_{\mathfrak{p}}$) has \mathcal{P} for all prime ideals \mathfrak{p} ”. (\mathcal{P} is preserved and reflected by localization to all primes.)

Examples.

- 1 $M = 0$.

Local properties

A property \mathcal{P} of a ring or module is local if “ A (or M) has \mathcal{P} ” iff “ $A_{\mathfrak{p}}$ (or $M_{\mathfrak{p}}$) has \mathcal{P} for all prime ideals \mathfrak{p} ”. (\mathcal{P} is preserved and reflected by localization to all primes.)

Examples.

- 1 $M = 0$.
- 2 $L \rightarrow M \rightarrow N$ is exact at M .

Local properties

A property \mathcal{P} of a ring or module is local if “ A (or M) has \mathcal{P} ” iff “ $A_{\mathfrak{p}}$ (or $M_{\mathfrak{p}}$) has \mathcal{P} for all prime ideals \mathfrak{p} ”. (\mathcal{P} is preserved and reflected by localization to all primes.)

Examples.

- 1 $M = 0$.
- 2 $L \rightarrow M \rightarrow N$ is exact at M .

Local properties

A property \mathcal{P} of a ring or module is local if “ A (or M) has \mathcal{P} ” iff “ $A_{\mathfrak{p}}$ (or $M_{\mathfrak{p}}$) has \mathcal{P} for all prime ideals \mathfrak{p} ”. (\mathcal{P} is preserved and reflected by localization to all primes.)

Examples.

- ① $M = 0$.
- ② $L \rightarrow M \rightarrow N$ is exact at M .

(1) [$M = 0$ locally implies $M = 0$; contradiction]

Local properties

A property \mathcal{P} of a ring or module is local if “ A (or M) has \mathcal{P} ” iff “ $A_{\mathfrak{p}}$ (or $M_{\mathfrak{p}}$) has \mathcal{P} for all prime ideals \mathfrak{p} ”. (\mathcal{P} is preserved and reflected by localization to all primes.)

Examples.

- ① $M = 0$.
- ② $L \rightarrow M \rightarrow N$ is exact at M .

(1) [$M = 0$ locally implies $M = 0$; contradiction]

Assume that $x \in M - \{0\}$.

Local properties

A property \mathcal{P} of a ring or module is local if “ A (or M) has \mathcal{P} ” iff “ $A_{\mathfrak{p}}$ (or $M_{\mathfrak{p}}$) has \mathcal{P} for all prime ideals \mathfrak{p} ”. (\mathcal{P} is preserved and reflected by localization to all primes.)

Examples.

- ① $M = 0$.
- ② $L \rightarrow M \rightarrow N$ is exact at M .

(1) [$M = 0$ locally implies $M = 0$; contradiction]

Assume that $x \in M - \{0\}$. Let \mathfrak{p} be a maximal ideal containing $\text{Ann}(x)$.

Local properties

A property \mathcal{P} of a ring or module is local if “ A (or M) has \mathcal{P} ” iff “ $A_{\mathfrak{p}}$ (or $M_{\mathfrak{p}}$) has \mathcal{P} for all prime ideals \mathfrak{p} ”. (\mathcal{P} is preserved and reflected by localization to all primes.)

Examples.

- ① $M = 0$.
- ② $L \rightarrow M \rightarrow N$ is exact at M .

(1) [$M = 0$ locally implies $M = 0$; contradiction]

Assume that $x \in M - \{0\}$. Let \mathfrak{p} be a maximal ideal containing $\text{Ann}(x)$.

Claim: $x/1 \in M_{\mathfrak{p}} - 0$.

Local properties

A property \mathcal{P} of a ring or module is local if “ A (or M) has \mathcal{P} ” iff “ $A_{\mathfrak{p}}$ (or $M_{\mathfrak{p}}$) has \mathcal{P} for all prime ideals \mathfrak{p} ”. (\mathcal{P} is preserved and reflected by localization to all primes.)

Examples.

- ① $M = 0$.
- ② $L \rightarrow M \rightarrow N$ is exact at M .

(1) [$M = 0$ locally implies $M = 0$; contradiction]

Assume that $x \in M - \{0\}$. Let \mathfrak{p} be a maximal ideal containing $\text{Ann}(x)$.

Claim: $x/1 \in M_{\mathfrak{p}} - 0$. Else $x/1 \sim 0/s$,

Local properties

A property \mathcal{P} of a ring or module is local if “ A (or M) has \mathcal{P} ” iff “ $A_{\mathfrak{p}}$ (or $M_{\mathfrak{p}}$) has \mathcal{P} for all prime ideals \mathfrak{p} ”. (\mathcal{P} is preserved and reflected by localization to all primes.)

Examples.

- ① $M = 0$.
- ② $L \rightarrow M \rightarrow N$ is exact at M .

(1) [$M = 0$ locally implies $M = 0$; contradiction]

Assume that $x \in M - \{0\}$. Let \mathfrak{p} be a maximal ideal containing $\text{Ann}(x)$.

Claim: $x/1 \in M_{\mathfrak{p}} - 0$. Else $x/1 \sim 0/s$, leading to $(\exists u)(usx = 0)$. But $us \in S = A - \mathfrak{p}$,

Local properties

A property \mathcal{P} of a ring or module is local if “ A (or M) has \mathcal{P} ” iff “ $A_{\mathfrak{p}}$ (or $M_{\mathfrak{p}}$) has \mathcal{P} for all prime ideals \mathfrak{p} ”. (\mathcal{P} is preserved and reflected by localization to all primes.)

Examples.

- ① $M = 0$.
- ② $L \rightarrow M \rightarrow N$ is exact at M .

(1) [$M = 0$ locally implies $M = 0$; contradiction]

Assume that $x \in M - \{0\}$. Let \mathfrak{p} be a maximal ideal containing $\text{Ann}(x)$.

Claim: $x/1 \in M_{\mathfrak{p}} - 0$. Else $x/1 \sim 0/s$, leading to $(\exists u)(usx = 0)$. But $us \in S = A - \mathfrak{p}$, so $us \notin \text{Ann}(x)$. \square

Local properties

A property \mathcal{P} of a ring or module is local if “ A (or M) has \mathcal{P} ” iff “ $A_{\mathfrak{p}}$ (or $M_{\mathfrak{p}}$) has \mathcal{P} for all prime ideals \mathfrak{p} ”. (\mathcal{P} is preserved and reflected by localization to all primes.)

Examples.

- ① $M = 0$.
- ② $L \rightarrow M \rightarrow N$ is exact at M .

(1) [$M = 0$ locally implies $M = 0$; contradiction]

Assume that $x \in M - \{0\}$. Let \mathfrak{p} be a maximal ideal containing $\text{Ann}(x)$.

Claim: $x/1 \in M_{\mathfrak{p}} - 0$. Else $x/1 \sim 0/s$, leading to $(\exists u)(usx = 0)$. But $us \in S = A - \mathfrak{p}$, so $us \notin \text{Ann}(x)$. \square

(2) [Exactness of complex is local; contradiction]

Local properties

A property \mathcal{P} of a ring or module is local if “ A (or M) has \mathcal{P} ” iff “ $A_{\mathfrak{p}}$ (or $M_{\mathfrak{p}}$) has \mathcal{P} for all prime ideals \mathfrak{p} ”. (\mathcal{P} is preserved and reflected by localization to all primes.)

Examples.

- ① $M = 0$.
- ② $L \rightarrow M \rightarrow N$ is exact at M .

(1) [$M = 0$ locally implies $M = 0$; contradiction]

Assume that $x \in M - \{0\}$. Let \mathfrak{p} be a maximal ideal containing $\text{Ann}(x)$.

Claim: $x/1 \in M_{\mathfrak{p}} - 0$. Else $x/1 \sim 0/s$, leading to $(\exists u)(usx = 0)$. But $us \in S = A - \mathfrak{p}$, so $us \notin \text{Ann}(x)$. \square

(2) [Exactness of complex is local; contradiction]

If $L \xrightarrow{\alpha} M \xrightarrow{\beta} N$ inexact at M , and $m \in \ker(\beta) - \text{im}(\alpha)$, then there is a maximal \mathfrak{p} containing $(\text{im}(\alpha) : m)$.

Local properties

A property \mathcal{P} of a ring or module is local if “ A (or M) has \mathcal{P} ” iff “ $A_{\mathfrak{p}}$ (or $M_{\mathfrak{p}}$) has \mathcal{P} for all prime ideals \mathfrak{p} ”. (\mathcal{P} is preserved and reflected by localization to all primes.)

Examples.

- ① $M = 0$.
- ② $L \rightarrow M \rightarrow N$ is exact at M .

(1) [$M = 0$ locally implies $M = 0$; contradiction]

Assume that $x \in M - \{0\}$. Let \mathfrak{p} be a maximal ideal containing $\text{Ann}(x)$.

Claim: $x/1 \in M_{\mathfrak{p}} - 0$. Else $x/1 \sim 0/s$, leading to $(\exists u)(usx = 0)$. But $us \in S = A - \mathfrak{p}$, so $us \notin \text{Ann}(x)$. \square

(2) [Exactness of complex is local; contradiction]

If $L \xrightarrow{\alpha} M \xrightarrow{\beta} N$ inexact at M , and $m \in \ker(\beta) - \text{im}(\alpha)$, then there is a maximal \mathfrak{p} containing $(\text{im}(\alpha) : m)$. Then $L_{\mathfrak{p}} \rightarrow M_{\mathfrak{p}} \rightarrow N_{\mathfrak{p}}$ inexact at $M_{\mathfrak{p}}$.

Local properties

A property \mathcal{P} of a ring or module is local if “ A (or M) has \mathcal{P} ” iff “ $A_{\mathfrak{p}}$ (or $M_{\mathfrak{p}}$) has \mathcal{P} for all prime ideals \mathfrak{p} ”. (\mathcal{P} is preserved and reflected by localization to all primes.)

Examples.

- ① $M = 0$.
- ② $L \rightarrow M \rightarrow N$ is exact at M .

(1) [$M = 0$ locally implies $M = 0$; contradiction]

Assume that $x \in M - \{0\}$. Let \mathfrak{p} be a maximal ideal containing $\text{Ann}(x)$.

Claim: $x/1 \in M_{\mathfrak{p}} - 0$. Else $x/1 \sim 0/s$, leading to $(\exists u)(usx = 0)$. But $us \in S = A - \mathfrak{p}$, so $us \notin \text{Ann}(x)$. \square

(2) [Exactness of complex is local; contradiction]

If $L \xrightarrow{\alpha} M \xrightarrow{\beta} N$ inexact at M , and $m \in \ker(\beta) - \text{im}(\alpha)$, then there is a maximal \mathfrak{p} containing $(\text{im}(\alpha) : m)$. Then $L_{\mathfrak{p}} \rightarrow M_{\mathfrak{p}} \rightarrow N_{\mathfrak{p}}$ inexact at $M_{\mathfrak{p}}$. \square

Check last claim

Assume that $L \xrightarrow{\alpha} M \xrightarrow{\beta} N$ is inexact at M , and that $m \in \ker(\beta) - \operatorname{im}(\alpha)$.

Check last claim

Assume that $L \xrightarrow{\alpha} M \xrightarrow{\beta} N$ is inexact at M , and that $m \in \ker(\beta) - \operatorname{im}(\alpha)$.
Assume that \mathfrak{p} is a maximal ideal containing $(\operatorname{im}(\alpha) : m)$.

Check last claim

Assume that $L \xrightarrow{\alpha} M \xrightarrow{\beta} N$ is inexact at M , and that $m \in \ker(\beta) - \operatorname{im}(\alpha)$.

Assume that \mathfrak{p} is a maximal ideal containing $(\operatorname{im}(\alpha) : m)$. Assume that

$L_{\mathfrak{p}} \xrightarrow{\alpha_{\mathfrak{p}}} M_{\mathfrak{p}} \xrightarrow{\beta_{\mathfrak{p}}} N_{\mathfrak{p}}$ is exact at $M_{\mathfrak{p}}$.

Check last claim

Assume that $L \xrightarrow{\alpha} M \xrightarrow{\beta} N$ is inexact at M , and that $m \in \ker(\beta) - \operatorname{im}(\alpha)$.

Assume that \mathfrak{p} is a maximal ideal containing $(\operatorname{im}(\alpha) : m)$. Assume that

$L_{\mathfrak{p}} \xrightarrow{\alpha_{\mathfrak{p}}} M_{\mathfrak{p}} \xrightarrow{\beta_{\mathfrak{p}}} N_{\mathfrak{p}}$ is exact at $M_{\mathfrak{p}}$.

Since $m \in \ker(\beta)$, $\beta_{\mathfrak{p}}(m/1) = \beta(m)/1 = 0$,

Check last claim

Assume that $L \xrightarrow{\alpha} M \xrightarrow{\beta} N$ is inexact at M , and that $m \in \ker(\beta) - \operatorname{im}(\alpha)$.

Assume that \mathfrak{p} is a maximal ideal containing $(\operatorname{im}(\alpha) : m)$. Assume that

$L_{\mathfrak{p}} \xrightarrow{\alpha_{\mathfrak{p}}} M_{\mathfrak{p}} \xrightarrow{\beta_{\mathfrak{p}}} N_{\mathfrak{p}}$ is exact at $M_{\mathfrak{p}}$.

Since $m \in \ker(\beta)$, $\beta_{\mathfrak{p}}(m/1) = \beta(m)/1 = 0$, so $m/1 \in \ker(\beta_{\mathfrak{p}})$.

Check last claim

Assume that $L \xrightarrow{\alpha} M \xrightarrow{\beta} N$ is inexact at M , and that $m \in \ker(\beta) - \operatorname{im}(\alpha)$. Assume that \mathfrak{p} is a maximal ideal containing $(\operatorname{im}(\alpha) : m)$. Assume that $L_{\mathfrak{p}} \xrightarrow{\alpha_{\mathfrak{p}}} M_{\mathfrak{p}} \xrightarrow{\beta_{\mathfrak{p}}} N_{\mathfrak{p}}$ is exact at $M_{\mathfrak{p}}$.

Since $m \in \ker(\beta)$, $\beta_{\mathfrak{p}}(m/1) = \beta(m)/1 = 0$, so $m/1 \in \ker(\beta_{\mathfrak{p}})$. Hence $m/1 = \alpha_{\mathfrak{p}}(\ell/s) = \alpha(\ell)/s$.

Check last claim

Assume that $L \xrightarrow{\alpha} M \xrightarrow{\beta} N$ is inexact at M , and that $m \in \ker(\beta) - \operatorname{im}(\alpha)$. Assume that \mathfrak{p} is a maximal ideal containing $(\operatorname{im}(\alpha) : m)$. Assume that $L_{\mathfrak{p}} \xrightarrow{\alpha_{\mathfrak{p}}} M_{\mathfrak{p}} \xrightarrow{\beta_{\mathfrak{p}}} N_{\mathfrak{p}}$ is exact at $M_{\mathfrak{p}}$.

Since $m \in \ker(\beta)$, $\beta_{\mathfrak{p}}(m/1) = \beta(m)/1 = 0$, so $m/1 \in \ker(\beta_{\mathfrak{p}})$. Hence $m/1 = \alpha_{\mathfrak{p}}(\ell/s) = \alpha(\ell)/s$. Hence $(\exists u)(u(\alpha(\ell) - sm) = 0)$,

Check last claim

Assume that $L \xrightarrow{\alpha} M \xrightarrow{\beta} N$ is inexact at M , and that $m \in \ker(\beta) - \operatorname{im}(\alpha)$.

Assume that \mathfrak{p} is a maximal ideal containing $(\operatorname{im}(\alpha) : m)$. Assume that

$L_{\mathfrak{p}} \xrightarrow{\alpha_{\mathfrak{p}}} M_{\mathfrak{p}} \xrightarrow{\beta_{\mathfrak{p}}} N_{\mathfrak{p}}$ is exact at $M_{\mathfrak{p}}$.

Since $m \in \ker(\beta)$, $\beta_{\mathfrak{p}}(m/1) = \beta(m)/1 = 0$, so $m/1 \in \ker(\beta_{\mathfrak{p}})$. Hence $m/1 = \alpha_{\mathfrak{p}}(\ell/s) = \alpha(\ell)/s$. Hence $(\exists u)(u(\alpha(\ell) - sm) = 0)$, or $\alpha(u\ell) = usm$.

Check last claim

Assume that $L \xrightarrow{\alpha} M \xrightarrow{\beta} N$ is inexact at M , and that $m \in \ker(\beta) - \operatorname{im}(\alpha)$.

Assume that \mathfrak{p} is a maximal ideal containing $(\operatorname{im}(\alpha) : m)$. Assume that

$L_{\mathfrak{p}} \xrightarrow{\alpha_{\mathfrak{p}}} M_{\mathfrak{p}} \xrightarrow{\beta_{\mathfrak{p}}} N_{\mathfrak{p}}$ is exact at $M_{\mathfrak{p}}$.

Since $m \in \ker(\beta)$, $\beta_{\mathfrak{p}}(m/1) = \beta(m)/1 = 0$, so $m/1 \in \ker(\beta_{\mathfrak{p}})$. Hence $m/1 = \alpha_{\mathfrak{p}}(\ell/s) = \alpha(\ell)/s$. Hence $(\exists u)(u(\alpha(\ell) - sm) = 0)$, or $\alpha(u\ell) = usm$. Hence $us \in (\operatorname{im}(\alpha) : m) \subseteq \mathfrak{p}$,

Check last claim

Assume that $L \xrightarrow{\alpha} M \xrightarrow{\beta} N$ is inexact at M , and that $m \in \ker(\beta) - \operatorname{im}(\alpha)$.

Assume that \mathfrak{p} is a maximal ideal containing $(\operatorname{im}(\alpha) : m)$. Assume that

$L_{\mathfrak{p}} \xrightarrow{\alpha_{\mathfrak{p}}} M_{\mathfrak{p}} \xrightarrow{\beta_{\mathfrak{p}}} N_{\mathfrak{p}}$ is exact at $M_{\mathfrak{p}}$.

Since $m \in \ker(\beta)$, $\beta_{\mathfrak{p}}(m/1) = \beta(m)/1 = 0$, so $m/1 \in \ker(\beta_{\mathfrak{p}})$. Hence $m/1 = \alpha_{\mathfrak{p}}(\ell/s) = \alpha(\ell)/s$. Hence $(\exists u)(u(\alpha(\ell) - sm) = 0)$, or $\alpha(u\ell) = usm$. Hence $us \in (\operatorname{im}(\alpha) : m) \subseteq \mathfrak{p}$, contradicting $u, s \in S = A - \mathfrak{p}$.

Check last claim

Assume that $L \xrightarrow{\alpha} M \xrightarrow{\beta} N$ is inexact at M , and that $m \in \ker(\beta) - \operatorname{im}(\alpha)$.

Assume that \mathfrak{p} is a maximal ideal containing $(\operatorname{im}(\alpha) : m)$. Assume that

$L_{\mathfrak{p}} \xrightarrow{\alpha_{\mathfrak{p}}} M_{\mathfrak{p}} \xrightarrow{\beta_{\mathfrak{p}}} N_{\mathfrak{p}}$ is exact at $M_{\mathfrak{p}}$.

Since $m \in \ker(\beta)$, $\beta_{\mathfrak{p}}(m/1) = \beta(m)/1 = 0$, so $m/1 \in \ker(\beta_{\mathfrak{p}})$. Hence $m/1 = \alpha_{\mathfrak{p}}(\ell/s) = \alpha(\ell)/s$. Hence $(\exists u)(u(\alpha(\ell) - sm) = 0)$, or $\alpha(u\ell) = usm$. Hence $us \in (\operatorname{im}(\alpha) : m) \subseteq \mathfrak{p}$, contradicting $u, s \in S = A - \mathfrak{p}$. \square

The effect of localization on Spec

The effect of localization on Spec

We view $\text{Spec}(A)$ as a topological space, and each $f \in A$ as a “continuous” or “polynomial” function defined on $\text{Spec}(A)$. The “value” of the function f at the point \mathfrak{p} is $f/\mathfrak{p} \in A/\mathfrak{p}$.

The effect of localization on Spec

We view $\text{Spec}(A)$ as a topological space, and each $f \in A$ as a “continuous” or “polynomial” function defined on $\text{Spec}(A)$. The “value” of the function f at the point \mathfrak{p} is $f/\mathfrak{p} \in A/\mathfrak{p}$. The function f “vanishes” at \mathfrak{p} if $f/\mathfrak{p} = 0/\mathfrak{p}$, i.e. if $f \in \mathfrak{p}$.

The effect of localization on Spec

We view $\text{Spec}(A)$ as a topological space, and each $f \in A$ as a “continuous” or “polynomial” function defined on $\text{Spec}(A)$. The “value” of the function f at the point \mathfrak{p} is $f/\mathfrak{p} \in A/\mathfrak{p}$. The function f “vanishes” at \mathfrak{p} if $f/\mathfrak{p} = 0/\mathfrak{p}$, i.e. if $f \in \mathfrak{p}$.

If $S = \{f\}^* = \{1, f, f^2, \dots\}$, then we often write A_f for $S^{-1}A$. These are viewed as rational functions of the form a/f^k defined on $\text{Spec}(A_f) = \text{Spec}(A) - V(f)$

The effect of localization on Spec

We view $\text{Spec}(A)$ as a topological space, and each $f \in A$ as a “continuous” or “polynomial” function defined on $\text{Spec}(A)$. The “value” of the function f at the point \mathfrak{p} is $f/\mathfrak{p} \in A/\mathfrak{p}$. The function f “vanishes” at \mathfrak{p} if $f/\mathfrak{p} = 0/\mathfrak{p}$, i.e. if $f \in \mathfrak{p}$.

If $S = \{f\}^* = \{1, f, f^2, \dots\}$, then we often write A_f for $S^{-1}A$. These are viewed as rational functions of the form a/f^k defined on $\text{Spec}(A_f) = \text{Spec}(A) - V(f) = D(f)$

The effect of localization on Spec

We view $\text{Spec}(A)$ as a topological space, and each $f \in A$ as a “continuous” or “polynomial” function defined on $\text{Spec}(A)$. The “value” of the function f at the point \mathfrak{p} is $f/\mathfrak{p} \in A/\mathfrak{p}$. The function f “vanishes” at \mathfrak{p} if $f/\mathfrak{p} = 0/\mathfrak{p}$, i.e. if $f \in \mathfrak{p}$.

If $S = \{f\}^* = \{1, f, f^2, \dots\}$, then we often write A_f for $S^{-1}A$. These are viewed as rational functions of the form a/f^k defined on $\text{Spec}(A_f) = \text{Spec}(A) - V(f) = D(f) = \text{supp}(f)$.

The effect of localization on Spec

We view $\text{Spec}(A)$ as a topological space, and each $f \in A$ as a “continuous” or “polynomial” function defined on $\text{Spec}(A)$. The “value” of the function f at the point \mathfrak{p} is $f/\mathfrak{p} \in A/\mathfrak{p}$. The function f “vanishes” at \mathfrak{p} if $f/\mathfrak{p} = 0/\mathfrak{p}$, i.e. if $f \in \mathfrak{p}$.

If $S = \{f\}^* = \{1, f, f^2, \dots\}$, then we often write A_f for $S^{-1}A$. These are viewed as rational functions of the form a/f^k defined on $\text{Spec}(A_f) = \text{Spec}(A) - V(f) = D(f) = \text{supp}(f)$.

More generally, for any MC S , the ring $S^{-1}A$ is viewed as the ring of rational functions of the form a/s defined on $\text{Spec}(S^{-1}A) = \text{Spec}(A) - \bigcup_{s \in S} V(s)$

The effect of localization on Spec

We view $\text{Spec}(A)$ as a topological space, and each $f \in A$ as a “continuous” or “polynomial” function defined on $\text{Spec}(A)$. The “value” of the function f at the point \mathfrak{p} is $f/\mathfrak{p} \in A/\mathfrak{p}$. The function f “vanishes” at \mathfrak{p} if $f/\mathfrak{p} = 0/\mathfrak{p}$, i.e. if $f \in \mathfrak{p}$.

If $S = \{f\}^* = \{1, f, f^2, \dots\}$, then we often write A_f for $S^{-1}A$. These are viewed as rational functions of the form a/f^k defined on $\text{Spec}(A_f) = \text{Spec}(A) - V(f) = D(f) = \text{supp}(f)$.

More generally, for any MC S , the ring $S^{-1}A$ is viewed as the ring of rational functions of the form a/s defined on $\text{Spec}(S^{-1}A) = \text{Spec}(A) - \bigcup_{s \in S} V(s) = \bigcap_{s \in S} D(s)$

The effect of localization on Spec

We view $\text{Spec}(A)$ as a topological space, and each $f \in A$ as a “continuous” or “polynomial” function defined on $\text{Spec}(A)$. The “value” of the function f at the point \mathfrak{p} is $f/\mathfrak{p} \in A/\mathfrak{p}$. The function f “vanishes” at \mathfrak{p} if $f/\mathfrak{p} = 0/\mathfrak{p}$, i.e. if $f \in \mathfrak{p}$.

If $S = \{f\}^* = \{1, f, f^2, \dots\}$, then we often write A_f for $S^{-1}A$. These are viewed as rational functions of the form a/f^k defined on $\text{Spec}(A_f) = \text{Spec}(A) - V(f) = D(f) = \text{supp}(f)$.

More generally, for any MC S , the ring $S^{-1}A$ is viewed as the ring of rational functions of the form a/s defined on $\text{Spec}(S^{-1}A) = \text{Spec}(A) - \bigcup_{s \in S} V(s) = \bigcap_{s \in S} D(s) = \bigcap_{s \in S} \text{supp}(s)$

The effect of localization on Spec

We view $\text{Spec}(A)$ as a topological space, and each $f \in A$ as a “continuous” or “polynomial” function defined on $\text{Spec}(A)$. The “value” of the function f at the point \mathfrak{p} is $f/\mathfrak{p} \in A/\mathfrak{p}$. The function f “vanishes” at \mathfrak{p} if $f/\mathfrak{p} = 0/\mathfrak{p}$, i.e. if $f \in \mathfrak{p}$.

If $S = \{f\}^* = \{1, f, f^2, \dots\}$, then we often write A_f for $S^{-1}A$. These are viewed as rational functions of the form a/f^k defined on $\text{Spec}(A_f) = \text{Spec}(A) - V(f) = D(f) = \text{supp}(f)$.

More generally, for any MC S , the ring $S^{-1}A$ is viewed as the ring of rational functions of the form a/s defined on $\text{Spec}(S^{-1}A) = \text{Spec}(A) - \bigcup_{s \in S} V(s) = \bigcap_{s \in S} D(s) = \bigcap_{s \in S} \text{supp}(s) = \text{supp}(S)$.

The effect of localization on Spec

We view $\text{Spec}(A)$ as a topological space, and each $f \in A$ as a “continuous” or “polynomial” function defined on $\text{Spec}(A)$. The “value” of the function f at the point \mathfrak{p} is $f/\mathfrak{p} \in A/\mathfrak{p}$. The function f “vanishes” at \mathfrak{p} if $f/\mathfrak{p} = 0/\mathfrak{p}$, i.e. if $f \in \mathfrak{p}$.

If $S = \{f\}^* = \{1, f, f^2, \dots\}$, then we often write A_f for $S^{-1}A$. These are viewed as rational functions of the form a/f^k defined on $\text{Spec}(A_f) = \text{Spec}(A) - V(f) = D(f) = \text{supp}(f)$.

More generally, for any MC S , the ring $S^{-1}A$ is viewed as the ring of rational functions of the form a/s defined on $\text{Spec}(S^{-1}A) = \text{Spec}(A) - \bigcup_{s \in S} V(s) = \bigcap_{s \in S} D(s) = \bigcap_{s \in S} \text{supp}(s) = \text{supp}(S)$.

An interesting special case is when $S = A - \mathfrak{p}$, where $\text{Spec}(A_{\mathfrak{p}})$ equals the intersection of all distinguished open sets containing the point \mathfrak{p} .