


# Chain conditions







ACC = every strictly ascending chain  $a_0 < a_1 < \dots$  is finite.



ACC = every strictly ascending chain  $a_0 < a_1 < \dots$  is finite.

This is the class of posets which omit the substructure  $\langle \omega; < \rangle$ .



ACC = every strictly ascending chain  $a_0 < a_1 < \dots$  is finite.  
This is the class of posets which omit the substructure  $\langle \omega; < \rangle$ .  
(DCC = dual.)



ACC = every strictly ascending chain  $a_0 < a_1 < \dots$  is finite.  
This is the class of posets which omit the substructure  $\langle \omega; < \rangle$ .  
(DCC = dual.)

**Easy observations.**



ACC = every strictly ascending chain  $a_0 < a_1 < \dots$  is finite.  
This is the class of posets which omit the substructure  $\langle \omega; < \rangle$ .  
(DCC = dual.)

## Easy observations.

- 1 A nonempty poset  $P$  has the DCC iff it is well-founded iff every nonempty subset has a minimal element.



ACC = every strictly ascending chain  $a_0 < a_1 < \dots$  is finite.  
This is the class of posets which omit the substructure  $\langle \omega; < \rangle$ .  
(DCC = dual.)

## Easy observations.

- 1 A nonempty poset  $P$  has the DCC iff it is well-founded iff every nonempty subset has a minimal element.





ACC = every strictly ascending chain  $a_0 < a_1 < \dots$  is finite.  
This is the class of posets which omit the substructure  $\langle \omega; < \rangle$ .  
(DCC = dual.)

## Easy observations.

- 1 A nonempty poset  $P$  has the DCC iff it is well-founded iff every nonempty subset has a minimal element. (ACC = dual)



ACC = every strictly ascending chain  $a_0 < a_1 < \dots$  is finite.  
This is the class of posets which omit the substructure  $\langle \omega; < \rangle$ .  
(DCC = dual.)

## Easy observations.

- 1 A nonempty poset  $P$  has the DCC iff it is well-founded iff every nonempty subset has a minimal element. (ACC = dual)
- 2 A nonempty poset  $P$  has the ACC+DCC iff every chain in  $P$  is finite.





TFAE and they define modularity for lattices.



TFAE and they define modularity for lattices.



TFAE and they define modularity for lattices.

- 1  $L$  satisfies  $(x \vee y) \wedge (y \vee z) = y \vee ((x \vee y) \wedge z)$ .



TFAE and they define modularity for lattices.

- 1  $L$  satisfies  $(x \vee y) \wedge (y \vee z) = y \vee ((x \vee y) \wedge z)$ .



TFAE and they define modularity for lattices.

- 1  $L$  satisfies  $(x \vee y) \wedge (y \vee z) = y \vee ((x \vee y) \wedge z)$ .
- 2  $L$  omits the pentagon as a sublattice.





TFAE and they define modularity for lattices.

- 1  $L$  satisfies  $(x \vee y) \wedge (y \vee z) = y \vee ((x \vee y) \wedge z)$ .
- 2  $L$  omits the pentagon as a sublattice.



TFAE and they define modularity for lattices.

- ①  $L$  satisfies  $(x \vee y) \wedge (y \vee z) = y \vee ((x \vee y) \wedge z)$ .
- ②  $L$  omits the pentagon as a sublattice.
- ③ Perspective intervals  $[a \wedge b, a]$  and  $[b, a \vee b]$  are isomorphic via  $x \mapsto b \vee x$  and  $y \mapsto a \wedge y$ .



TFAE and they define modularity for lattices.

- ①  $L$  satisfies  $(x \vee y) \wedge (y \vee z) = y \vee ((x \vee y) \wedge z)$ .
- ②  $L$  omits the pentagon as a sublattice.
- ③ Perspective intervals  $[a \wedge b, a]$  and  $[b, a \vee b]$  are isomorphic via  $x \mapsto b \vee x$  and  $y \mapsto a \wedge y$ .



TFAE and they define modularity for lattices.

- ①  $L$  satisfies  $(x \vee y) \wedge (y \vee z) = y \vee ((x \vee y) \wedge z)$ .
- ②  $L$  omits the pentagon as a sublattice.
- ③ Perspective intervals  $[a \wedge b, a]$  and  $[b, a \vee b]$  are isomorphic via  $x \mapsto b \vee x$  and  $y \mapsto a \wedge y$ .

Modularity is a self-dual property.



TFAE and they define modularity for lattices.

- ①  $L$  satisfies  $(x \vee y) \wedge (y \vee z) = y \vee ((x \vee y) \wedge z)$ .
- ②  $L$  omits the pentagon as a sublattice.
- ③ Perspective intervals  $[a \wedge b, a]$  and  $[b, a \vee b]$  are isomorphic via  $x \mapsto b \vee x$  and  $y \mapsto a \wedge y$ .

Modularity is a self-dual property.

The normal subgroup lattice of a group is modular.



TFAE and they define modularity for lattices.

- ①  $L$  satisfies  $(x \vee y) \wedge (y \vee z) = y \vee ((x \vee y) \wedge z)$ .
- ②  $L$  omits the pentagon as a sublattice.
- ③ Perspective intervals  $[a \wedge b, a]$  and  $[b, a \vee b]$  are isomorphic via  $x \mapsto b \vee x$  and  $y \mapsto a \wedge y$ .

Modularity is a self-dual property.

The normal subgroup lattice of a group is modular. The submodule lattice of a module is modular.



TFAE and they define modularity for lattices.

- ①  $L$  satisfies  $(x \vee y) \wedge (y \vee z) = y \vee ((x \vee y) \wedge z)$ .
- ②  $L$  omits the pentagon as a sublattice.
- ③ Perspective intervals  $[a \wedge b, a]$  and  $[b, a \vee b]$  are isomorphic via  $x \mapsto b \vee x$  and  $y \mapsto a \wedge y$ .


Modularity is a self-dual property.

The normal subgroup lattice of a group is modular. The submodule lattice of a module is modular. The left, right, or 2-sided ideal lattices of a ring are modular.

# Chain conditions for modular lattices



# Chain conditions for modular lattices



# Chain conditions for complemented modular lattices

# Chain conditions for complemented modular lattices

As one might expect, a complemented modular lattice has ACC iff it has DCC.

# Chain conditions for complemented modular lattices

As one might expect, a complemented modular lattice has ACC iff it has DCC. One might expect that whenever  $a_0 < a_1 < a_2 < \cdots$ , and  $a_i^\perp$  is a complement of  $a_i$  for each  $i$ , then  $a_0^\perp > a_1^\perp > a_2^\perp > \cdots$ .

# Chain conditions for complemented modular lattices

As one might expect, a complemented modular lattice has ACC iff it has DCC. One might expect that whenever  $a_0 < a_1 < a_2 < \cdots$ , and  $a_i^\perp$  is a complement of  $a_i$  for each  $i$ , then  $a_0^\perp > a_1^\perp > a_2^\perp > \cdots$ . This turns out to be false, due to the fact that complements need not be unique.

# Chain conditions for complemented modular lattices

As one might expect, a complemented modular lattice has ACC iff it has DCC. One might expect that whenever  $a_0 < a_1 < a_2 < \cdots$ , and  $a_i^\perp$  is a complement of  $a_i$  for each  $i$ , then  $a_0^\perp > a_1^\perp > a_2^\perp > \cdots$ . This turns out to be false, due to the fact that complements need not be unique. Instead:

# Chain conditions for complemented modular lattices

As one might expect, a complemented modular lattice has ACC iff it has DCC. One might expect that whenever  $a_0 < a_1 < a_2 < \cdots$ , and  $a_i^\perp$  is a complement of  $a_i$  for each  $i$ , then  $a_0^\perp > a_1^\perp > a_2^\perp > \cdots$ . This turns out to be false, due to the fact that complements need not be unique. Instead:

**Lm.** If  $L$  is a modular lattice and  $a < b$  are both complemented, then for any complement  $a^\perp$  of  $a$  there exists a complement  $b^\perp$  of  $b$ , such that  $a^\perp > b^\perp$ .

# Chain conditions for complemented modular lattices

As one might expect, a complemented modular lattice has ACC iff it has DCC. One might expect that whenever  $a_0 < a_1 < a_2 < \dots$ , and  $a_i^\perp$  is a complement of  $a_i$  for each  $i$ , then  $a_0^\perp > a_1^\perp > a_2^\perp > \dots$ . This turns out to be false, due to the fact that complements need not be unique. Instead:

**Lm.** If  $L$  is a modular lattice and  $a < b$  are both complemented, then for any complement  $a^\perp$  of  $a$  there exists a complement  $b^\perp$  of  $b$ , such that  $a^\perp > b^\perp$ .

*Proof.*



# Chain conditions for complemented modular lattices

As one might expect, a complemented modular lattice has ACC iff it has DCC. One might expect that whenever  $a_0 < a_1 < a_2 < \dots$ , and  $a_i^\perp$  is a complement of  $a_i$  for each  $i$ , then  $a_0^\perp > a_1^\perp > a_2^\perp > \dots$ . This turns out to be false, due to the fact that complements need not be unique. Instead:

**Lm.** If  $L$  is a modular lattice and  $a < b$  are both complemented, then for any complement  $a^\perp$  of  $a$  there exists a complement  $b^\perp$  of  $b$ , such that  $a^\perp > b^\perp$ .

*Proof.*

Given a complement  $a^\perp$  of  $a$ ,

# Chain conditions for complemented modular lattices

As one might expect, a complemented modular lattice has ACC iff it has DCC. One might expect that whenever  $a_0 < a_1 < a_2 < \dots$ , and  $a_i^\perp$  is a complement of  $a_i$  for each  $i$ , then  $a_0^\perp > a_1^\perp > a_2^\perp > \dots$ . This turns out to be false, due to the fact that complements need not be unique. Instead:

**Lm.** If  $L$  is a modular lattice and  $a < b$  are both complemented, then for any complement  $a^\perp$  of  $a$  there exists a complement  $b^\perp$  of  $b$ , such that  $a^\perp > b^\perp$ .

*Proof.*

Given a complement  $a^\perp$  of  $a$ , choose any complement  $b'$  of  $b$ .

# Chain conditions for complemented modular lattices

As one might expect, a complemented modular lattice has ACC iff it has DCC. One might expect that whenever  $a_0 < a_1 < a_2 < \dots$ , and  $a_i^\perp$  is a complement of  $a_i$  for each  $i$ , then  $a_0^\perp > a_1^\perp > a_2^\perp > \dots$ . This turns out to be false, due to the fact that complements need not be unique. Instead:

**Lm.** If  $L$  is a modular lattice and  $a < b$  are both complemented, then for any complement  $a^\perp$  of  $a$  there exists a complement  $b^\perp$  of  $b$ , such that  $a^\perp > b^\perp$ .

*Proof.*

Given a complement  $a^\perp$  of  $a$ , choose any complement  $b'$  of  $b$ . Define  $b^\perp := a^\perp \wedge (a \vee b')$ .

# Chain conditions for complemented modular lattices

As one might expect, a complemented modular lattice has ACC iff it has DCC. One might expect that whenever  $a_0 < a_1 < a_2 < \dots$ , and  $a_i^\perp$  is a complement of  $a_i$  for each  $i$ , then  $a_0^\perp > a_1^\perp > a_2^\perp > \dots$ . This turns out to be false, due to the fact that complements need not be unique. Instead:

**Lm.** If  $L$  is a modular lattice and  $a < b$  are both complemented, then for any complement  $a^\perp$  of  $a$  there exists a complement  $b^\perp$  of  $b$ , such that  $a^\perp > b^\perp$ .

*Proof.*

Given a complement  $a^\perp$  of  $a$ , choose any complement  $b'$  of  $b$ . Define  $b^\perp := a^\perp \wedge (a \vee b')$ . Check that  $b^\perp$  is a complement of  $b$  and  $a^\perp > b^\perp$ .

# Chain conditions for complemented modular lattices

As one might expect, a complemented modular lattice has ACC iff it has DCC. One might expect that whenever  $a_0 < a_1 < a_2 < \dots$ , and  $a_i^\perp$  is a complement of  $a_i$  for each  $i$ , then  $a_0^\perp > a_1^\perp > a_2^\perp > \dots$ . This turns out to be false, due to the fact that complements need not be unique. Instead:

**Lm.** If  $L$  is a modular lattice and  $a < b$  are both complemented, then for any complement  $a^\perp$  of  $a$  there exists a complement  $b^\perp$  of  $b$ , such that  $a^\perp > b^\perp$ .

*Proof.*

Given a complement  $a^\perp$  of  $a$ , choose any complement  $b'$  of  $b$ . Define  $b^\perp := a^\perp \wedge (a \vee b')$ . Check that  $b^\perp$  is a complement of  $b$  and  $a^\perp > b^\perp$ .  $\square$

# Chain conditions for complemented modular lattices

As one might expect, a complemented modular lattice has ACC iff it has DCC. One might expect that whenever  $a_0 < a_1 < a_2 < \cdots$ , and  $a_i^\perp$  is a complement of  $a_i$  for each  $i$ , then  $a_0^\perp > a_1^\perp > a_2^\perp > \cdots$ . This turns out to be false, due to the fact that complements need not be unique. Instead:

**Lm.** If  $L$  is a modular lattice and  $a < b$  are both complemented, then for any complement  $a^\perp$  of  $a$  there exists a complement  $b^\perp$  of  $b$ , such that  $a^\perp > b^\perp$ .

*Proof.*

Given a complement  $a^\perp$  of  $a$ , choose any complement  $b'$  of  $b$ . Define  $b^\perp := a^\perp \wedge (a \vee b')$ . Check that  $b^\perp$  is a complement of  $b$  and  $a^\perp > b^\perp$ .  $\square$

**Examples.** Subspace lattices of  $k$ -vector spaces are complemented modular lattices.

# Chain conditions for complemented modular lattices

As one might expect, a complemented modular lattice has ACC iff it has DCC. One might expect that whenever  $a_0 < a_1 < a_2 < \cdots$ , and  $a_i^\perp$  is a complement of  $a_i$  for each  $i$ , then  $a_0^\perp > a_1^\perp > a_2^\perp > \cdots$ . This turns out to be false, due to the fact that complements need not be unique. Instead:

**Lm.** If  $L$  is a modular lattice and  $a < b$  are both complemented, then for any complement  $a^\perp$  of  $a$  there exists a complement  $b^\perp$  of  $b$ , such that  $a^\perp > b^\perp$ .

*Proof.*

Given a complement  $a^\perp$  of  $a$ , choose any complement  $b'$  of  $b$ . Define  $b^\perp := a^\perp \wedge (a \vee b')$ . Check that  $b^\perp$  is a complement of  $b$  and  $a^\perp > b^\perp$ .  $\square$

**Examples.** Subspace lattices of  $k$ -vector spaces are complemented modular lattices. If  $A = k_1 \times \cdots \times k_n$  is a finite product of fields, then submodule lattices of  $A$ -modules are complemented modular lattices.

# Consequences of modularity in ring theory



# Consequences of modularity in ring theory

- ① Any finite (subdirect) product of Noeth. rings or modules is Noeth.

# Consequences of modularity in ring theory

- ① Any finite (subdirect) product of Noeth. rings or modules is Noeth.

# Consequences of modularity in ring theory

- ① Any finite (subdirect) product of Noeth. rings or modules is Noeth.
- ② (Modularity unnecessary) Any quotient of a Noeth. ring or module is Noeth. Any submodule of a Noeth. module is Noeth.

# Consequences of modularity in ring theory

- ① Any finite (subdirect) product of Noeth. rings or modules is Noeth.
- ② (Modularity unnecessary) Any quotient of a Noeth. ring or module is Noeth. Any submodule of a Noeth. module is Noeth.

# Consequences of modularity in ring theory

- ① Any finite (subdirect) product of Noeth. rings or modules is Noeth.
- ② (Modularity unnecessary) Any quotient of a Noeth. ring or module is Noeth. Any submodule of a Noeth. module is Noeth.
- ③ If  $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$  is exact, then  $M$  is Noeth. iff both  $L$  and  $N$  are.

# Consequences of modularity in ring theory

- ① Any finite (subdirect) product of Noeth. rings or modules is Noeth.
- ② (Modularity unnecessary) Any quotient of a Noeth. ring or module is Noeth. Any submodule of a Noeth. module is Noeth.
- ③ If  $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$  is exact, then  $M$  is Noeth. iff both  $L$  and  $N$  are.

# Consequences of modularity in ring theory

- ① Any finite (subdirect) product of Noeth. rings or modules is Noeth.
- ② (Modularity unnecessary) Any quotient of a Noeth. ring or module is Noeth. Any submodule of a Noeth. module is Noeth.
- ③ If  $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$  is exact, then  $M$  is Noeth. iff both  $L$  and  $N$  are.
- ④ F.g. free modules over Noetherian rings are Noetherian:  
 $0 \rightarrow {}_A A \rightarrow {}_A A \oplus {}_A A \rightarrow {}_A A \rightarrow 0$  is exact.

# Consequences of modularity in ring theory

- ① Any finite (subdirect) product of Noeth. rings or modules is Noeth.
- ② (Modularity unnecessary) Any quotient of a Noeth. ring or module is Noeth. Any submodule of a Noeth. module is Noeth.
- ③ If  $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$  is exact, then  $M$  is Noeth. iff both  $L$  and  $N$  are.
- ④ F.g. free modules over Noetherian rings are Noetherian:  
 $0 \rightarrow {}_A A \rightarrow {}_A A \oplus {}_A A \rightarrow {}_A A \rightarrow 0$  is exact.



# Consequences of modularity in ring theory

- ① Any finite (subdirect) product of Noeth. rings or modules is Noeth.
- ② (Modularity unnecessary) Any quotient of a Noeth. ring or module is Noeth. Any submodule of a Noeth. module is Noeth.
- ③ If  $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$  is exact, then  $M$  is Noeth. iff both  $L$  and  $N$  are.
- ④ F.g. free modules over Noetherian rings are Noetherian:  
 $0 \rightarrow {}_A A \rightarrow {}_A A \oplus {}_A A \rightarrow {}_A A \rightarrow 0$  is exact.
- ⑤ If  $A$  is a Noeth. commutative ring, then any finitely generated  $A$ -module  $N$  is Noeth.

# Consequences of modularity in ring theory

- ① Any finite (subdirect) product of Noeth. rings or modules is Noeth.
- ② (Modularity unnecessary) Any quotient of a Noeth. ring or module is Noeth. Any submodule of a Noeth. module is Noeth.
- ③ If  $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$  is exact, then  $M$  is Noeth. iff both  $L$  and  $N$  are.
- ④ F.g. free modules over Noetherian rings are Noetherian:  
 $0 \rightarrow {}_A A \rightarrow {}_A A \oplus {}_A A \rightarrow {}_A A \rightarrow 0$  is exact.
- ⑤ If  $A$  is a Noeth. commutative ring, then any finitely generated  $A$ -module  $N$  is Noeth.

# Consequences of modularity in ring theory

- ① Any finite (subdirect) product of Noeth. rings or modules is Noeth.
- ② (Modularity unnecessary) Any quotient of a Noeth. ring or module is Noeth. Any submodule of a Noeth. module is Noeth.
- ③ If  $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$  is exact, then  $M$  is Noeth. iff both  $L$  and  $N$  are.
- ④ F.g. free modules over Noetherian rings are Noetherian:  
 $0 \rightarrow {}_A A \rightarrow {}_A A \oplus {}_A A \rightarrow {}_A A \rightarrow 0$  is exact.
- ⑤ If  $A$  is a Noeth. commutative ring, then any finitely generated  $A$ -module  $N$  is Noeth. ( $A$  Noeth.  $\Rightarrow {}_A A$  Noeth.  $\Rightarrow (\bigoplus^n {}_A A)$  Noeth.  $\Rightarrow (\bigoplus^n {}_A A)/L$  Noeth.)

# Consequences of modularity in ring theory

- ① Any finite (subdirect) product of Noeth. rings or modules is Noeth.
- ② (Modularity unnecessary) Any quotient of a Noeth. ring or module is Noeth. Any submodule of a Noeth. module is Noeth.
- ③ If  $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$  is exact, then  $M$  is Noeth. iff both  $L$  and  $N$  are.
- ④ F.g. free modules over Noetherian rings are Noetherian:  
 $0 \rightarrow {}_A A \rightarrow {}_A A \oplus {}_A A \rightarrow {}_A A \rightarrow 0$  is exact.
- ⑤ If  $A$  is a Noeth. commutative ring, then any finitely generated  $A$ -module  $N$  is Noeth. ( $A$  Noeth.  $\Rightarrow {}_A A$  Noeth.  $\Rightarrow (\bigoplus^n {}_A A)$  Noeth.  $\Rightarrow (\bigoplus^n {}_A A)/L$  Noeth.)

All remain true if ‘Noetherian’ is replaced by ‘Artinian’.





Ideal generation is an algebraic closure operator, so  $A$  Noetherian  $\Leftrightarrow$  every ideal of  $A$  is f.g.



Ideal generation is an algebraic closure operator, so  $A$  Noetherian  $\Leftrightarrow$  every ideal of  $A$  is f.g.

**The Hilbert Basis Theorem.** If  $A$  is Noetherian, then  $A[x]$  is Noetherian.



Ideal generation is an algebraic closure operator, so  $A$  Noetherian  $\Leftrightarrow$  every ideal of  $A$  is f.g.

**The Hilbert Basis Theorem.** If  $A$  is Noetherian, then  $A[x]$  is Noetherian.

*Proof.* Given  $I \triangleleft A[x]$ ,





Ideal generation is an algebraic closure operator, so  $A$  Noetherian  $\Leftrightarrow$  every ideal of  $A$  is f.g.

**The Hilbert Basis Theorem.** If  $A$  is Noetherian, then  $A[x]$  is Noetherian.

*Proof.* Given  $I \triangleleft A[x]$ , let  $L = (a_1, \dots, a_n) \triangleleft A$  be the ideal of leading coefficients of members of  $I$ ,



Ideal generation is an algebraic closure operator, so  $A$  Noetherian  $\Leftrightarrow$  every ideal of  $A$  is f.g.

**The Hilbert Basis Theorem.** If  $A$  is Noetherian, then  $A[x]$  is Noetherian.

*Proof.* Given  $I \triangleleft A[x]$ , let  $L = (a_1, \dots, a_n) \triangleleft A$  be the ideal of leading coefficients of members of  $I$ , say  $f_i = a_i x^{e_i} + \dots \in I$ .



Ideal generation is an algebraic closure operator, so  $A$  Noetherian  $\Leftrightarrow$  every ideal of  $A$  is f.g.

**The Hilbert Basis Theorem.** If  $A$  is Noetherian, then  $A[x]$  is Noetherian.

*Proof.* Given  $I \triangleleft A[x]$ , let  $L = (a_1, \dots, a_n) \triangleleft A$  be the ideal of leading coefficients of members of  $I$ , say  $f_i = a_i x^{e_i} + \dots \in I$ . Let  $e = \max\{e_i\}$ .



Ideal generation is an algebraic closure operator, so  $A$  Noetherian  $\Leftrightarrow$  every ideal of  $A$  is f.g.

**The Hilbert Basis Theorem.** If  $A$  is Noetherian, then  $A[x]$  is Noetherian.

*Proof.* Given  $I \triangleleft A[x]$ , let  $L = (a_1, \dots, a_n) \triangleleft A$  be the ideal of leading coefficients of members of  $I$ , say  $f_i = a_i x^{e_i} + \dots \in I$ . Let  $e = \max\{e_i\}$ .  
 $I = (f_1, \dots, f_n) + I \cap M$  where  $M = \langle 1, x, \dots, x^{e-1} \rangle_{A \text{ Mod}}$ .



Ideal generation is an algebraic closure operator, so  $A$  Noetherian  $\Leftrightarrow$  every ideal of  $A$  is f.g.

**The Hilbert Basis Theorem.** If  $A$  is Noetherian, then  $A[x]$  is Noetherian.

*Proof.* Given  $I \triangleleft A[x]$ , let  $L = (a_1, \dots, a_n) \triangleleft A$  be the ideal of leading coefficients of members of  $I$ , say  $f_i = a_i x^{e_i} + \dots \in I$ . Let  $e = \max\{e_i\}$ .  $I = (f_1, \dots, f_n) + I \cap M$  where  $M = \langle 1, x, \dots, x^{e-1} \rangle_{A \text{ Mod. } M}$ .  $M$  is Noetherian as an  $A$ -module,



Ideal generation is an algebraic closure operator, so  $A$  Noetherian  $\Leftrightarrow$  every ideal of  $A$  is f.g.

**The Hilbert Basis Theorem.** If  $A$  is Noetherian, then  $A[x]$  is Noetherian.

*Proof.* Given  $I \triangleleft A[x]$ , let  $L = (a_1, \dots, a_n) \triangleleft A$  be the ideal of leading coefficients of members of  $I$ , say  $f_i = a_i x^{e_i} + \dots \in I$ . Let  $e = \max\{e_i\}$ .  $I = (f_1, \dots, f_n) + I \cap M$  where  $M = \langle 1, x, \dots, x^{e-1} \rangle_{A \text{ Mod. } M}$ .  $M$  is Noetherian as an  $A$ -module, so  $I$  is f.g. as an ideal.



Ideal generation is an algebraic closure operator, so  $A$  Noetherian  $\Leftrightarrow$  every ideal of  $A$  is f.g.

**The Hilbert Basis Theorem.** If  $A$  is Noetherian, then  $A[x]$  is Noetherian.

*Proof.* Given  $I \triangleleft A[x]$ , let  $L = (a_1, \dots, a_n) \triangleleft A$  be the ideal of leading coefficients of members of  $I$ , say  $f_i = a_i x^{e_i} + \dots \in I$ . Let  $e = \max\{e_i\}$ .  $I = (f_1, \dots, f_n) + I \cap M$  where  $M = \langle 1, x, \dots, x^{e-1} \rangle_{A \text{ Mod. } M}$ .  $M$  is Noetherian as an  $A$ -module, so  $I$  is f.g. as an ideal.  $\square$

# Coordinate rings are Noetherian



# Coordinate rings are Noetherian

The equation  $x^2 + y^2 = 1$  defines a circle  $\Gamma$  in  $\mathbb{R}^2$ .

# Coordinate rings are Noetherian

The equation  $x^2 + y^2 = 1$  defines a circle  $\Gamma$  in  $\mathbb{R}^2$ .

The ring  $\mathbb{R}[x, y]$  may be viewed as the ring of polynomial functions from  $\mathbb{R}^2 \rightarrow \mathbb{R}$ .

# Coordinate rings are Noetherian

The equation  $x^2 + y^2 = 1$  defines a circle  $\Gamma$  in  $\mathbb{R}^2$ .

The ring  $\mathbb{R}[x, y]$  may be viewed as the ring of polynomial functions from  $\mathbb{R}^2 \rightarrow \mathbb{R}$ .

Elements  $g \in \mathbb{R}[x, y]$  can be restricted to  $\Gamma$  to obtain polynomial functions  $g: \Gamma \rightarrow \mathbb{R}$ .

# Coordinate rings are Noetherian

The equation  $x^2 + y^2 = 1$  defines a circle  $\Gamma$  in  $\mathbb{R}^2$ .

The ring  $\mathbb{R}[x, y]$  may be viewed as the ring of polynomial functions from  $\mathbb{R}^2 \rightarrow \mathbb{R}$ .

Elements  $g \in \mathbb{R}[x, y]$  can be restricted to  $\Gamma$  to obtain polynomial functions  $g: \Gamma \rightarrow \mathbb{R}$ . The ring of restricted functions is called the coordinate ring of  $\Gamma$ .

# Coordinate rings are Noetherian

The equation  $x^2 + y^2 = 1$  defines a circle  $\Gamma$  in  $\mathbb{R}^2$ .

The ring  $\mathbb{R}[x, y]$  may be viewed as the ring of polynomial functions from  $\mathbb{R}^2 \rightarrow \mathbb{R}$ .

Elements  $g \in \mathbb{R}[x, y]$  can be restricted to  $\Gamma$  to obtain polynomial functions  $g: \Gamma \rightarrow \mathbb{R}$ . The ring of restricted functions is called the coordinate ring of  $\Gamma$ .

In this example, the coordinate ring  $\mathbb{A}(\Gamma)$  is isomorphic to

$\mathbb{R}[x, y]/(x^2 + y^2 - 1)$ , since  $g, h \in \mathbb{R}[x, y]$  have the same restriction to  $\Gamma$  iff  $g - h$  restricts to the zero function on  $\Gamma$  iff  $g - h$  is divisible by the irreducible polynomial  $x^2 + y^2 - 1$ .

# Coordinate rings are Noetherian

The equation  $x^2 + y^2 = 1$  defines a circle  $\Gamma$  in  $\mathbb{R}^2$ .

The ring  $\mathbb{R}[x, y]$  may be viewed as the ring of polynomial functions from  $\mathbb{R}^2 \rightarrow \mathbb{R}$ .

Elements  $g \in \mathbb{R}[x, y]$  can be restricted to  $\Gamma$  to obtain polynomial functions  $g: \Gamma \rightarrow \mathbb{R}$ . The ring of restricted functions is called the coordinate ring of  $\Gamma$ .

In this example, the coordinate ring  $\mathbb{A}(\Gamma)$  is isomorphic to

$\mathbb{R}[x, y]/(x^2 + y^2 - 1)$ , since  $g, h \in \mathbb{R}[x, y]$  have the same restriction to  $\Gamma$  iff  $g - h$  restricts to the zero function on  $\Gamma$  iff  $g - h$  is divisible by the irreducible polynomial  $x^2 + y^2 - 1$ .

This can be generalized to any field  $k$  in place of  $\mathbb{R}$

# Coordinate rings are Noetherian

The equation  $x^2 + y^2 = 1$  defines a circle  $\Gamma$  in  $\mathbb{R}^2$ .

The ring  $\mathbb{R}[x, y]$  may be viewed as the ring of polynomial functions from  $\mathbb{R}^2 \rightarrow \mathbb{R}$ .

Elements  $g \in \mathbb{R}[x, y]$  can be restricted to  $\Gamma$  to obtain polynomial functions  $g: \Gamma \rightarrow \mathbb{R}$ . The ring of restricted functions is called the coordinate ring of  $\Gamma$ .

In this example, the coordinate ring  $\mathbb{A}(\Gamma)$  is isomorphic to

$\mathbb{R}[x, y]/(x^2 + y^2 - 1)$ , since  $g, h \in \mathbb{R}[x, y]$  have the same restriction to  $\Gamma$  iff  $g - h$  restricts to the zero function on  $\Gamma$  iff  $g - h$  is divisible by the irreducible polynomial  $x^2 + y^2 - 1$ .

This can be generalized to any field  $k$  in place of  $\mathbb{R}$  and any algebraic subset  $\Gamma \subseteq k^n$ .

# Coordinate rings are Noetherian

The equation  $x^2 + y^2 = 1$  defines a circle  $\Gamma$  in  $\mathbb{R}^2$ .

The ring  $\mathbb{R}[x, y]$  may be viewed as the ring of polynomial functions from  $\mathbb{R}^2 \rightarrow \mathbb{R}$ .

Elements  $g \in \mathbb{R}[x, y]$  can be restricted to  $\Gamma$  to obtain polynomial functions  $g: \Gamma \rightarrow \mathbb{R}$ . The ring of restricted functions is called the coordinate ring of  $\Gamma$ .

In this example, the coordinate ring  $\mathbb{A}(\Gamma)$  is isomorphic to

$\mathbb{R}[x, y]/(x^2 + y^2 - 1)$ , since  $g, h \in \mathbb{R}[x, y]$  have the same restriction to  $\Gamma$  iff  $g - h$  restricts to the zero function on  $\Gamma$  iff  $g - h$  is divisible by the irreducible polynomial  $x^2 + y^2 - 1$ .

This can be generalized to any field  $k$  in place of  $\mathbb{R}$  and any algebraic subset  $\Gamma \subseteq k^n$ . (Algebraic set = common zero locus of a collection of polynomials.)



# Coordinate rings are Noetherian

The equation  $x^2 + y^2 = 1$  defines a circle  $\Gamma$  in  $\mathbb{R}^2$ .

The ring  $\mathbb{R}[x, y]$  may be viewed as the ring of polynomial functions from  $\mathbb{R}^2 \rightarrow \mathbb{R}$ .

Elements  $g \in \mathbb{R}[x, y]$  can be restricted to  $\Gamma$  to obtain polynomial functions  $g: \Gamma \rightarrow \mathbb{R}$ . The ring of restricted functions is called the coordinate ring of  $\Gamma$ .

In this example, the coordinate ring  $\mathbb{A}(\Gamma)$  is isomorphic to

$\mathbb{R}[x, y]/(x^2 + y^2 - 1)$ , since  $g, h \in \mathbb{R}[x, y]$  have the same restriction to  $\Gamma$  iff  $g - h$  restricts to the zero function on  $\Gamma$  iff  $g - h$  is divisible by the irreducible polynomial  $x^2 + y^2 - 1$ .

This can be generalized to any field  $k$  in place of  $\mathbb{R}$  and any algebraic subset  $\Gamma \subseteq k^n$ . (Algebraic set = common zero locus of a collection of polynomials.)  $\mathbb{A}(\Gamma)$  is the coordinate ring of  $\Gamma$ .

# Coordinate rings are Noetherian

The equation  $x^2 + y^2 = 1$  defines a circle  $\Gamma$  in  $\mathbb{R}^2$ .

The ring  $\mathbb{R}[x, y]$  may be viewed as the ring of polynomial functions from  $\mathbb{R}^2 \rightarrow \mathbb{R}$ .

Elements  $g \in \mathbb{R}[x, y]$  can be restricted to  $\Gamma$  to obtain polynomial functions  $g: \Gamma \rightarrow \mathbb{R}$ . The ring of restricted functions is called the coordinate ring of  $\Gamma$ .

In this example, the coordinate ring  $\mathbb{A}(\Gamma)$  is isomorphic to

$\mathbb{R}[x, y]/(x^2 + y^2 - 1)$ , since  $g, h \in \mathbb{R}[x, y]$  have the same restriction to  $\Gamma$  iff  $g - h$  restricts to the zero function on  $\Gamma$  iff  $g - h$  is divisible by the irreducible polynomial  $x^2 + y^2 - 1$ .

This can be generalized to any field  $k$  in place of  $\mathbb{R}$  and any algebraic subset  $\Gamma \subseteq k^n$ . (Algebraic set = common zero locus of a collection of polynomials.)

$\mathbb{A}(\Gamma)$  is the coordinate ring of  $\Gamma$ . It is Noetherian since it is a quotient of  $k[x_1, \dots, x_n]$ .

# One last thing about Noetherian rings

# One last thing about Noetherian rings

**Thm.**

## One last thing about Noetherian rings

**Thm.** If  $A$  is Noetherian, then every ideal  $I \triangleleft A$  contains a power of its radical,  $\sqrt{I}$ .

## One last thing about Noetherian rings

**Thm.** If  $A$  is Noetherian, then every ideal  $I \triangleleft A$  contains a power of its radical,  $\sqrt{I}$ .

*Proof.*

## One last thing about Noetherian rings

**Thm.** If  $A$  is Noetherian, then every ideal  $I \triangleleft A$  contains a power of its radical,  $\sqrt{I}$ .

*Proof.*

Suffices to work in  $A/I$  and prove  $(\exists k)(\mathfrak{R}^k = (0))$ .

# One last thing about Noetherian rings

**Thm.** If  $A$  is Noetherian, then every ideal  $I \triangleleft A$  contains a power of its radical,  $\sqrt{I}$ .

*Proof.*

Suffices to work in  $A/I$  and prove  $(\exists k)(\mathfrak{R}^k = (0))$ .

If  $\mathfrak{R} = (a_1, \dots, a_n)$ , where  $a_i^{e_i} = 0$ ,



# One last thing about Noetherian rings

**Thm.** If  $A$  is Noetherian, then every ideal  $I \triangleleft A$  contains a power of its radical,  $\sqrt{I}$ .

*Proof.*

Suffices to work in  $A/I$  and prove  $(\exists k)(\mathfrak{N}^k = (0))$ .

If  $\mathfrak{N} = (a_1, \dots, a_n)$ , where  $a_i^{e_i} = 0$ , then  $\mathfrak{N}^k$  is generated by elements of the form  $a_1^{f_1} \cdots a_n^{f_n}$ ,  $\sum f_i = k$ .

# One last thing about Noetherian rings

**Thm.** If  $A$  is Noetherian, then every ideal  $I \triangleleft A$  contains a power of its radical,  $\sqrt{I}$ .

*Proof.*

Suffices to work in  $A/I$  and prove  $(\exists k)(\mathfrak{N}^k = (0))$ .

If  $\mathfrak{N} = (a_1, \dots, a_n)$ , where  $a_i^{e_i} = 0$ , then  $\mathfrak{N}^k$  is generated by elements of the form  $a_1^{f_1} \cdots a_n^{f_n}$ ,  $\sum f_i = k$ . If  $k \geq \sum e_i$ , then  $(\exists j)(f_j \geq e_j)$ ,

## One last thing about Noetherian rings

**Thm.** If  $A$  is Noetherian, then every ideal  $I \triangleleft A$  contains a power of its radical,  $\sqrt{I}$ .

*Proof.*

Suffices to work in  $A/I$  and prove  $(\exists k)(\mathfrak{R}^k = (0))$ .

If  $\mathfrak{R} = (a_1, \dots, a_n)$ , where  $a_i^{e_i} = 0$ , then  $\mathfrak{R}^k$  is generated by elements of the form  $a_1^{f_1} \cdots a_n^{f_n}$ ,  $\sum f_i = k$ . If  $k \geq \sum e_i$ , then  $(\exists j)(f_j \geq e_j)$ , so  $a_1^{f_1} \cdots a_n^{f_n} = 0$ .

## One last thing about Noetherian rings

**Thm.** If  $A$  is Noetherian, then every ideal  $I \triangleleft A$  contains a power of its radical,  $\sqrt{I}$ .

*Proof.*

Suffices to work in  $A/I$  and prove  $(\exists k)(\mathfrak{N}^k = (0))$ .

If  $\mathfrak{N} = (a_1, \dots, a_n)$ , where  $a_i^{e_i} = 0$ , then  $\mathfrak{N}^k$  is generated by elements of the form  $a_1^{f_1} \cdots a_n^{f_n}$ ,  $\sum f_i = k$ . If  $k \geq \sum e_i$ , then  $(\exists j)(f_j \geq e_j)$ , so  $a_1^{f_1} \cdots a_n^{f_n} = 0$ . Hence  $\mathfrak{N}^k = (0)$ .

# One last thing about Noetherian rings

**Thm.** If  $A$  is Noetherian, then every ideal  $I \triangleleft A$  contains a power of its radical,  $\sqrt{I}$ .

*Proof.*

Suffices to work in  $A/I$  and prove  $(\exists k)(\mathfrak{N}^k = (0))$ .

If  $\mathfrak{N} = (a_1, \dots, a_n)$ , where  $a_i^{e_i} = 0$ , then  $\mathfrak{N}^k$  is generated by elements of the form  $a_1^{f_1} \cdots a_n^{f_n}$ ,  $\sum f_i = k$ . If  $k \geq \sum e_i$ , then  $(\exists j)(f_j \geq e_j)$ , so  $a_1^{f_1} \cdots a_n^{f_n} = 0$ . Hence  $\mathfrak{N}^k = (0)$ .  $\square$

# One last thing about Noetherian rings

**Thm.** If  $A$  is Noetherian, then every ideal  $I \triangleleft A$  contains a power of its radical,  $\sqrt{I}$ .

*Proof.*

Suffices to work in  $A/I$  and prove  $(\exists k)(\mathfrak{R}^k = (0))$ .

If  $\mathfrak{R} = (a_1, \dots, a_n)$ , where  $a_i^{e_i} = 0$ , then  $\mathfrak{R}^k$  is generated by elements of the form  $a_1^{f_1} \cdots a_n^{f_n}$ ,  $\sum f_i = k$ . If  $k \geq \sum e_i$ , then  $(\exists j)(f_j \geq e_j)$ , so  $a_1^{f_1} \cdots a_n^{f_n} = 0$ . Hence  $\mathfrak{R}^k = (0)$ .  $\square$

**Example.**

# One last thing about Noetherian rings

**Thm.** If  $A$  is Noetherian, then every ideal  $I \triangleleft A$  contains a power of its radical,  $\sqrt{I}$ .

*Proof.*

Suffices to work in  $A/I$  and prove  $(\exists k)(\mathfrak{N}^k = (0))$ .

If  $\mathfrak{N} = (a_1, \dots, a_n)$ , where  $a_i^{e_i} = 0$ , then  $\mathfrak{N}^k$  is generated by elements of the form  $a_1^{f_1} \cdots a_n^{f_n}$ ,  $\sum f_i = k$ . If  $k \geq \sum e_i$ , then  $(\exists j)(f_j \geq e_j)$ , so  $a_1^{f_1} \cdots a_n^{f_n} = 0$ . Hence  $\mathfrak{N}^k = (0)$ .  $\square$

**Example.** (Noetherianness is a necessary hypothesis)

## One last thing about Noetherian rings

**Thm.** If  $A$  is Noetherian, then every ideal  $I \triangleleft A$  contains a power of its radical,  $\sqrt{I}$ .

*Proof.*

Suffices to work in  $A/I$  and prove  $(\exists k)(\mathfrak{R}^k = (0))$ .

If  $\mathfrak{R} = (a_1, \dots, a_n)$ , where  $a_i^{e_i} = 0$ , then  $\mathfrak{R}^k$  is generated by elements of the form  $a_1^{f_1} \cdots a_n^{f_n}$ ,  $\sum f_i = k$ . If  $k \geq \sum e_i$ , then  $(\exists j)(f_j \geq e_j)$ , so  $a_1^{f_1} \cdots a_n^{f_n} = 0$ . Hence  $\mathfrak{R}^k = (0)$ .  $\square$

**Example.** (Noetherianness is a necessary hypothesis)

$A = \langle x_1, x_2, x_3, \dots \mid x_1^2 = 0, x_2^2 = x_1, x_3^2 = x_2, \dots \rangle$  has  $\mathfrak{R} = (x_1, x_2, \dots)$ , and  $\mathfrak{R}^2 = \mathfrak{R} \neq (0)$ .







**Thm.** A commutative ring is Artinian iff it is a finite product of Artinian local rings.



**Thm.** A commutative ring is Artinian iff it is a finite product of Artinian local rings.

*Proof.*



**Thm.** A commutative ring is Artinian iff it is a finite product of Artinian local rings.

*Proof.*

[If] Modularity.



**Thm.** A commutative ring is Artinian iff it is a finite product of Artinian local rings.

*Proof.*

[If] Modularity.

[Only if] Main stages:



**Thm.** A commutative ring is Artinian iff it is a finite product of Artinian local rings.

*Proof.*

[If] Modularity.

[Only if] Main stages:

- 1 Stage 1: Show that any  $I \triangleleft A$  minimal for the property that it contains a nonunit, nonnilpotent element must be  $I = (e)$  for some idempotent  $e$ .



**Thm.** A commutative ring is Artinian iff it is a finite product of Artinian local rings.

*Proof.*

[If] Modularity.

[Only if] Main stages:

- 1 Stage 1: Show that any  $I \triangleleft A$  minimal for the property that it contains a nonunit, nonnilpotent element must be  $I = (e)$  for some idempotent  $e$ .
- 2 Stage 2: Show that  $A \cong A/(e) \times A/(1 - e)$  and the second factor is an Artinian local ring.



**Thm.** A commutative ring is Artinian iff it is a finite product of Artinian local rings.

*Proof.*

[If] Modularity.

[Only if] Main stages:

- 1 Stage 1: Show that any  $I \triangleleft A$  minimal for the property that it contains a nonunit, nonnilpotent element must be  $I = (e)$  for some idempotent  $e$ .
- 2 Stage 2: Show that  $A \cong A/(e) \times A/(1 - e)$  and the second factor is an Artinian local ring.
- 3 Stage 3: Continue splitting off local factors. This must terminate in finitely many steps with  $A$  isomorphic to a product of Artinian local rings.





# Stages

Stage 1:

# Stages

Stage 1: Let  $I \triangleleft A$  be minimal for the property that it contains a nonunit, nonnilpotent element,  $a$ .

# Stages

Stage 1: Let  $I \triangleleft A$  be minimal for the property that it contains a nonunit, nonnilpotent element,  $a$ .

$$I = (a) = (a^2),$$

# Stages

Stage 1: Let  $I \triangleleft A$  be minimal for the property that it contains a nonunit, nonnilpotent element,  $a$ .

$I = (a) = (a^2)$ , so  $(\exists r)(ra^2 = a)$ .

# Stages

Stage 1: Let  $I \triangleleft A$  be minimal for the property that it contains a nonunit, nonnilpotent element,  $a$ .

$I = (a) = (a^2)$ , so  $(\exists r)(ra^2 = a)$ .

If  $e = ra$ , then  $e^2 = r(ra^2) = ra = e$ .

# Stages

Stage 1: Let  $I \triangleleft A$  be minimal for the property that it contains a nonunit, nonnilpotent element,  $a$ .

$I = (a) = (a^2)$ , so  $(\exists r)(ra^2 = a)$ .

If  $e = ra$ , then  $e^2 = r(ra^2) = ra = e$ .

Moreover,  $I = (a) \supseteq (ra) = (e) \supseteq (ra^2) = (a) = I$ , so  $I = (e)$ .

# Stages

Stage 1: Let  $I \triangleleft A$  be minimal for the property that it contains a nonunit, nonnilpotent element,  $a$ .

$I = (a) = (a^2)$ , so  $(\exists r)(ra^2 = a)$ .

If  $e = ra$ , then  $e^2 = r(ra^2) = ra = e$ .

Moreover,  $I = (a) \supseteq (ra) = (e) \supseteq (ra^2) = (a) = I$ , so  $I = (e)$ .

Stage 2:



# Stages

Stage 1: Let  $I \triangleleft A$  be minimal for the property that it contains a nonunit, nonnilpotent element,  $a$ .

$I = (a) = (a^2)$ , so  $(\exists r)(ra^2 = a)$ .

If  $e = ra$ , then  $e^2 = r(ra^2) = ra = e$ .

Moreover,  $I = (a) \supseteq (ra) = (e) \supseteq (ra^2) = (a) = I$ , so  $I = (e)$ .

Stage 2:  $A \cong A/(e) \times A/(1 - e)$  since central idempotents yield direct decompositions.

Stage 1: Let  $I \triangleleft A$  be minimal for the property that it contains a nonunit, nonnilpotent element,  $a$ .

$I = (a) = (a^2)$ , so  $(\exists r)(ra^2 = a)$ .

If  $e = ra$ , then  $e^2 = r(ra^2) = ra = e$ .

Moreover,  $I = (a) \supseteq (ra) = (e) \supseteq (ra^2) = (a) = I$ , so  $I = (e)$ .

Stage 2:  $A \cong A/(e) \times A/(1 - e)$  since central idempotents yield direct decompositions. The ideal lattice of  $A/(1 - e)$  is isomorphic to the interval  $[(1 - e), A]$  in the ideal lattice of  $A$ , which is perspective with  $[0, (e)]$ .

Stage 1: Let  $I \triangleleft A$  be minimal for the property that it contains a nonunit, nonnilpotent element,  $a$ .

$I = (a) = (a^2)$ , so  $(\exists r)(ra^2 = a)$ .

If  $e = ra$ , then  $e^2 = r(ra^2) = ra = e$ .

Moreover,  $I = (a) \supseteq (ra) = (e) \supseteq (ra^2) = (a) = I$ , so  $I = (e)$ .

Stage 2:  $A \cong A/(e) \times A/(1 - e)$  since central idempotents yield direct decompositions. The ideal lattice of  $A/(1 - e)$  is isomorphic to the interval  $[(1 - e), A]$  in the ideal lattice of  $A$ , which is perspective with  $[0, (e)]$ . The latter interval has a maximal proper element  $(e) \cap \mathfrak{N}$ .

# Stages

Stage 1: Let  $I \triangleleft A$  be minimal for the property that it contains a nonunit, nonnilpotent element,  $a$ .

$I = (a) = (a^2)$ , so  $(\exists r)(ra^2 = a)$ .

If  $e = ra$ , then  $e^2 = r(ra^2) = ra = e$ .

Moreover,  $I = (a) \supseteq (ra) = (e) \supseteq (ra^2) = (a) = I$ , so  $I = (e)$ .

Stage 2:  $A \cong A/(e) \times A/(1 - e)$  since central idempotents yield direct decompositions. The ideal lattice of  $A/(1 - e)$  is isomorphic to the interval  $[(1 - e), A]$  in the ideal lattice of  $A$ , which is perspective with  $[0, (e)]$ . The latter interval has a maximal proper element  $(e) \cap \mathfrak{N}$ .

Stage 3:

# Stages

Stage 1: Let  $I \triangleleft A$  be minimal for the property that it contains a nonunit, nonnilpotent element,  $a$ .

$I = (a) = (a^2)$ , so  $(\exists r)(ra^2 = a)$ .

If  $e = ra$ , then  $e^2 = r(ra^2) = ra = e$ .

Moreover,  $I = (a) \supseteq (ra) = (e) \supseteq (ra^2) = (a) = I$ , so  $I = (e)$ .

Stage 2:  $A \cong A/(e) \times A/(1 - e)$  since central idempotents yield direct decompositions. The ideal lattice of  $A/(1 - e)$  is isomorphic to the interval  $[(1 - e), A]$  in the ideal lattice of  $A$ , which is perspective with  $[0, (e)]$ . The latter interval has a maximal proper element  $(e) \cap \mathfrak{N}$ .

Stage 3: We argue that the number of direct factors cannot exceed the number of maximal ideals, which is finite.

# Stages

Stage 1: Let  $I \triangleleft A$  be minimal for the property that it contains a nonunit, nonnilpotent element,  $a$ .

$I = (a) = (a^2)$ , so  $(\exists r)(ra^2 = a)$ .

If  $e = ra$ , then  $e^2 = r(ra^2) = ra = e$ .

Moreover,  $I = (a) \supseteq (ra) = (e) \supseteq (ra^2) = (a) = I$ , so  $I = (e)$ .

Stage 2:  $A \cong A/(e) \times A/(1 - e)$  since central idempotents yield direct decompositions. The ideal lattice of  $A/(1 - e)$  is isomorphic to the interval  $[(1 - e), A]$  in the ideal lattice of  $A$ , which is perspective with  $[0, (e)]$ . The latter interval has a maximal proper element  $(e) \cap \mathfrak{N}$ .

Stage 3: We argue that the number of direct factors cannot exceed the number of maximal ideals, which is finite. The first element of this is from calg1p4.

# Stages

Stage 1: Let  $I \triangleleft A$  be minimal for the property that it contains a nonunit, nonnilpotent element,  $a$ .

$I = (a) = (a^2)$ , so  $(\exists r)(ra^2 = a)$ .

If  $e = ra$ , then  $e^2 = r(ra^2) = ra = e$ .

Moreover,  $I = (a) \supseteq (ra) = (e) \supseteq (ra^2) = (a) = I$ , so  $I = (e)$ .

Stage 2:  $A \cong A/(e) \times A/(1 - e)$  since central idempotents yield direct decompositions. The ideal lattice of  $A/(1 - e)$  is isomorphic to the interval  $[(1 - e), A]$  in the ideal lattice of  $A$ , which is perspective with  $[0, (e)]$ . The latter interval has a maximal proper element  $(e) \cap \mathfrak{N}$ .

Stage 3: We argue that the number of direct factors cannot exceed the number of maximal ideals, which is finite. The first element of this is from calg1p4. For the second element, assume that  $\mathfrak{m}_0, \mathfrak{m}_1, \dots$  is an infinite sequence of distinct maximal ideals.

# Stages

Stage 1: Let  $I \triangleleft A$  be minimal for the property that it contains a nonunit, nonnilpotent element,  $a$ .

$I = (a) = (a^2)$ , so  $(\exists r)(ra^2 = a)$ .

If  $e = ra$ , then  $e^2 = r(ra^2) = ra = e$ .

Moreover,  $I = (a) \supseteq (ra) = (e) \supseteq (ra^2) = (a) = I$ , so  $I = (e)$ .

Stage 2:  $A \cong A/(e) \times A/(1 - e)$  since central idempotents yield direct decompositions. The ideal lattice of  $A/(1 - e)$  is isomorphic to the interval  $[(1 - e), A]$  in the ideal lattice of  $A$ , which is perspective with  $[0, (e)]$ . The latter interval has a maximal proper element  $(e) \cap \mathfrak{N}$ .

Stage 3: We argue that the number of direct factors cannot exceed the number of maximal ideals, which is finite. The first element of this is from calg1p4. For the second element, assume that  $\mathfrak{m}_0, \mathfrak{m}_1, \dots$  is an infinite sequence of distinct maximal ideals. Then  $\mathfrak{m}_0 > \mathfrak{m}_0 \cap \mathfrak{m}_1 > \mathfrak{m}_0 \cap \mathfrak{m}_1 \cap \mathfrak{m}_2 > \dots$  is an infinite strictly decreasing chain of ideals.



# More on the structure of Artinian rings

## More on the structure of Artinian rings

**Lm.** The maximal ideal of an Artinian local ring is nilpotent.

## More on the structure of Artinian rings

**Lm.** The maximal ideal of an Artinian local ring is nilpotent.

*Proof.*

## More on the structure of Artinian rings

**Lm.** The maximal ideal of an Artinian local ring is nilpotent.

*Proof.* Assume  $\mathfrak{m} \supseteq \mathfrak{m}^2 \supseteq \cdots \supseteq \mathfrak{m}^k = \mathfrak{m}^{k+1} = I \neq 0$ .

# More on the structure of Artinian rings

**Lm.** The maximal ideal of an Artinian local ring is nilpotent.

*Proof.* Assume  $\mathfrak{m} \geq \mathfrak{m}^2 \geq \cdots \geq \mathfrak{m}^k = \mathfrak{m}^{k+1} = I \neq 0$ . Choose  $a \in I$  so that  $(a)$  is minimal for  $I \cdot (a) \neq 0$ .

# More on the structure of Artinian rings

**Lm.** The maximal ideal of an Artinian local ring is nilpotent.

*Proof.* Assume  $\mathfrak{m} \geq \mathfrak{m}^2 \geq \cdots \geq \mathfrak{m}^k = \mathfrak{m}^{k+1} = I \neq 0$ . Choose  $a \in I$  so that  $(a)$  is minimal for  $I \cdot (a) \neq 0$ . Since  $I^2 = I$ ,

# More on the structure of Artinian rings

**Lm.** The maximal ideal of an Artinian local ring is nilpotent.

*Proof.* Assume  $\mathfrak{m} \geq \mathfrak{m}^2 \geq \cdots \geq \mathfrak{m}^k = \mathfrak{m}^{k+1} = I \neq 0$ . Choose  $a \in I$  so that  $(a)$  is minimal for  $I \cdot (a) \neq 0$ . Since  $I^2 = I$ ,  $I(I(a)) = I(a) \neq 0$ ,

# More on the structure of Artinian rings

**Lm.** The maximal ideal of an Artinian local ring is nilpotent.

*Proof.* Assume  $\mathfrak{m} \geq \mathfrak{m}^2 \geq \cdots \geq \mathfrak{m}^k = \mathfrak{m}^{k+1} = I \neq 0$ . Choose  $a \in I$  so that  $(a)$  is minimal for  $I \cdot (a) \neq 0$ . Since  $I^2 = I$ ,  $I(I(a)) = I(a) \neq 0$ , so  $I(a) = (a)$ .



# More on the structure of Artinian rings

**Lm.** The maximal ideal of an Artinian local ring is nilpotent.

*Proof.* Assume  $\mathfrak{m} \geq \mathfrak{m}^2 \geq \cdots \geq \mathfrak{m}^k = \mathfrak{m}^{k+1} = I \neq 0$ . Choose  $a \in I$  so that  $(a)$  is minimal for  $I \cdot (a) \neq 0$ . Since  $I^2 = I$ ,  $I(I(a)) = I(a) \neq 0$ , so  $I(a) = (a)$ . By NAK,  $(a) = 0$ ,

# More on the structure of Artinian rings

**Lm.** The maximal ideal of an Artinian local ring is nilpotent.

*Proof.* Assume  $\mathfrak{m} \supseteq \mathfrak{m}^2 \supseteq \cdots \supseteq \mathfrak{m}^k = \mathfrak{m}^{k+1} = I \neq 0$ . Choose  $a \in I$  so that  $(a)$  is minimal for  $I \cdot (a) \neq 0$ . Since  $I^2 = I$ ,  $I(I(a)) = I(a) \neq 0$ , so  $I(a) = (a)$ . By NAK,  $(a) = 0$ , contradiction.

# More on the structure of Artinian rings

**Lm.** The maximal ideal of an Artinian local ring is nilpotent.

*Proof.* Assume  $\mathfrak{m} \supseteq \mathfrak{m}^2 \supseteq \cdots \supseteq \mathfrak{m}^k = \mathfrak{m}^{k+1} = I \neq 0$ . Choose  $a \in I$  so that  $(a)$  is minimal for  $I \cdot (a) \neq 0$ . Since  $I^2 = I$ ,  $I(I(a)) = I(a) \neq 0$ , so  $I(a) = (a)$ . By NAK,  $(a) = 0$ , contradiction.  $\square$

# More on the structure of Artinian rings

**Lm.** The maximal ideal of an Artinian local ring is nilpotent.

*Proof.* Assume  $\mathfrak{m} \geq \mathfrak{m}^2 \geq \cdots \geq \mathfrak{m}^k = \mathfrak{m}^{k+1} = I \neq 0$ . Choose  $a \in I$  so that  $(a)$  is minimal for  $I \cdot (a) \neq 0$ . Since  $I^2 = I$ ,  $I(I(a)) = I(a) \neq 0$ , so  $I(a) = (a)$ . By NAK,  $(a) = 0$ , contradiction.  $\square$

**Thm.**

# More on the structure of Artinian rings

**Lm.** The maximal ideal of an Artinian local ring is nilpotent.

*Proof.* Assume  $\mathfrak{m} \supseteq \mathfrak{m}^2 \supseteq \cdots \supseteq \mathfrak{m}^k = \mathfrak{m}^{k+1} = I \neq 0$ . Choose  $a \in I$  so that  $(a)$  is minimal for  $I \cdot (a) \neq 0$ . Since  $I^2 = I$ ,  $I(I(a)) = I(a) \neq 0$ , so  $I(a) = (a)$ . By NAK,  $(a) = 0$ , contradiction.  $\square$

**Thm.** An Artinian ring has Krull dimension zero.

# More on the structure of Artinian rings

**Lm.** The maximal ideal of an Artinian local ring is nilpotent.

*Proof.* Assume  $\mathfrak{m} \supseteq \mathfrak{m}^2 \supseteq \cdots \supseteq \mathfrak{m}^k = \mathfrak{m}^{k+1} = I \neq 0$ . Choose  $a \in I$  so that  $(a)$  is minimal for  $I \cdot (a) \neq 0$ . Since  $I^2 = I$ ,  $I(I(a)) = I(a) \neq 0$ , so  $I(a) = (a)$ . By NAK,  $(a) = 0$ , contradiction.  $\square$

**Thm.** An Artinian ring has Krull dimension zero. (All primes are maximal.)

# More on the structure of Artinian rings

**Lm.** The maximal ideal of an Artinian local ring is nilpotent.

*Proof.* Assume  $\mathfrak{m} \supseteq \mathfrak{m}^2 \supseteq \cdots \supseteq \mathfrak{m}^k = \mathfrak{m}^{k+1} = I \neq 0$ . Choose  $a \in I$  so that  $(a)$  is minimal for  $I \cdot (a) \neq 0$ . Since  $I^2 = I$ ,  $I(I(a)) = I(a) \neq 0$ , so  $I(a) = (a)$ . By NAK,  $(a) = 0$ , contradiction.  $\square$

**Thm.** An Artinian ring has Krull dimension zero. (All primes are maximal.)

*Proof.*

# More on the structure of Artinian rings

**Lm.** The maximal ideal of an Artinian local ring is nilpotent.

*Proof.* Assume  $\mathfrak{m} \geq \mathfrak{m}^2 \geq \cdots \geq \mathfrak{m}^k = \mathfrak{m}^{k+1} = I \neq 0$ . Choose  $a \in I$  so that  $(a)$  is minimal for  $I \cdot (a) \neq 0$ . Since  $I^2 = I$ ,  $I(I(a)) = I(a) \neq 0$ , so  $I(a) = (a)$ . By NAK,  $(a) = 0$ , contradiction.  $\square$

**Thm.** An Artinian ring has Krull dimension zero. (All primes are maximal.)

*Proof.* Local rings with nilpotent maximal ideal have Krull dimension zero,



# More on the structure of Artinian rings

**Lm.** The maximal ideal of an Artinian local ring is nilpotent.

*Proof.* Assume  $\mathfrak{m} \geq \mathfrak{m}^2 \geq \cdots \geq \mathfrak{m}^k = \mathfrak{m}^{k+1} = I \neq 0$ . Choose  $a \in I$  so that  $(a)$  is minimal for  $I \cdot (a) \neq 0$ . Since  $I^2 = I$ ,  $I(I(a)) = I(a) \neq 0$ , so  $I(a) = (a)$ . By NAK,  $(a) = 0$ , contradiction.  $\square$

**Thm.** An Artinian ring has Krull dimension zero. (All primes are maximal.)

*Proof.* Local rings with nilpotent maximal ideal have Krull dimension zero, and, by  $\text{calg1p4}$ ,  $\dim(R \times S) = \max\{\dim(R), \dim(S)\}$ .

# More on the structure of Artinian rings

**Lm.** The maximal ideal of an Artinian local ring is nilpotent.

*Proof.* Assume  $\mathfrak{m} \geq \mathfrak{m}^2 \geq \cdots \geq \mathfrak{m}^k = \mathfrak{m}^{k+1} = I \neq 0$ . Choose  $a \in I$  so that  $(a)$  is minimal for  $I \cdot (a) \neq 0$ . Since  $I^2 = I$ ,  $I(I(a)) = I(a) \neq 0$ , so  $I(a) = (a)$ . By NAK,  $(a) = 0$ , contradiction.  $\square$

**Thm.** An Artinian ring has Krull dimension zero. (All primes are maximal.)

*Proof.* Local rings with nilpotent maximal ideal have Krull dimension zero, and, by  $\text{calg1p4}$ ,  $\dim(R \times S) = \max\{\dim(R), \dim(S)\}$ .  $\square$

# More on the structure of Artinian rings

**Lm.** The maximal ideal of an Artinian local ring is nilpotent.

*Proof.* Assume  $\mathfrak{m} \geq \mathfrak{m}^2 \geq \cdots \geq \mathfrak{m}^k = \mathfrak{m}^{k+1} = I \neq 0$ . Choose  $a \in I$  so that  $(a)$  is minimal for  $I \cdot (a) \neq 0$ . Since  $I^2 = I$ ,  $I(I(a)) = I(a) \neq 0$ , so  $I(a) = (a)$ . By NAK,  $(a) = 0$ , contradiction.  $\square$

**Thm.** An Artinian ring has Krull dimension zero. (All primes are maximal.)

*Proof.* Local rings with nilpotent maximal ideal have Krull dimension zero, and, by  $\text{calg1p4}$ ,  $\dim(R \times S) = \max\{\dim(R), \dim(S)\}$ .  $\square$

**Thm.** (Commutative version of Hopkins-Levitski)

# More on the structure of Artinian rings

**Lm.** The maximal ideal of an Artinian local ring is nilpotent.

*Proof.* Assume  $\mathfrak{m} \supseteq \mathfrak{m}^2 \supseteq \cdots \supseteq \mathfrak{m}^k = \mathfrak{m}^{k+1} = I \neq 0$ . Choose  $a \in I$  so that  $(a)$  is minimal for  $I \cdot (a) \neq 0$ . Since  $I^2 = I$ ,  $I(I(a)) = I(a) \neq 0$ , so  $I(a) = (a)$ . By NAK,  $(a) = 0$ , contradiction.  $\square$

**Thm.** An Artinian ring has Krull dimension zero. (All primes are maximal.)

*Proof.* Local rings with nilpotent maximal ideal have Krull dimension zero, and, by  $\text{calg1p4}$ ,  $\dim(R \times S) = \max\{\dim(R), \dim(S)\}$ .  $\square$

**Thm.** (Commutative version of Hopkins-Levitski)  
An Artinian ring is Noetherian.

# More on the structure of Artinian rings

**Lm.** The maximal ideal of an Artinian local ring is nilpotent.

*Proof.* Assume  $\mathfrak{m} \geq \mathfrak{m}^2 \geq \cdots \geq \mathfrak{m}^k = \mathfrak{m}^{k+1} = I \neq 0$ . Choose  $a \in I$  so that  $(a)$  is minimal for  $I \cdot (a) \neq 0$ . Since  $I^2 = I$ ,  $I(I(a)) = I(a) \neq 0$ , so  $I(a) = (a)$ . By NAK,  $(a) = 0$ , contradiction.  $\square$

**Thm.** An Artinian ring has Krull dimension zero. (All primes are maximal.)

*Proof.* Local rings with nilpotent maximal ideal have Krull dimension zero, and, by  $\text{calg1p4}$ ,  $\dim(R \times S) = \max\{\dim(R), \dim(S)\}$ .  $\square$

**Thm.** (Commutative version of Hopkins-Levitski)  
An Artinian ring is Noetherian.

*Proof.*

# More on the structure of Artinian rings

**Lm.** The maximal ideal of an Artinian local ring is nilpotent.

*Proof.* Assume  $\mathfrak{m} \geq \mathfrak{m}^2 \geq \cdots \geq \mathfrak{m}^k = \mathfrak{m}^{k+1} = I \neq 0$ . Choose  $a \in I$  so that  $(a)$  is minimal for  $I \cdot (a) \neq 0$ . Since  $I^2 = I$ ,  $I(I(a)) = I(a) \neq 0$ , so  $I(a) = (a)$ . By NAK,  $(a) = 0$ , contradiction.  $\square$

**Thm.** An Artinian ring has Krull dimension zero. (All primes are maximal.)

*Proof.* Local rings with nilpotent maximal ideal have Krull dimension zero, and, by  $\text{calg1p4}$ ,  $\dim(R \times S) = \max\{\dim(R), \dim(S)\}$ .  $\square$

**Thm.** (Commutative version of Hopkins-Levitski)  
An Artinian ring is Noetherian.

*Proof.* Suffices to prove it for local  $A$ .

# More on the structure of Artinian rings

**Lm.** The maximal ideal of an Artinian local ring is nilpotent.

*Proof.* Assume  $\mathfrak{m} \geq \mathfrak{m}^2 \geq \cdots \geq \mathfrak{m}^k = \mathfrak{m}^{k+1} = I \neq 0$ . Choose  $a \in I$  so that  $(a)$  is minimal for  $I \cdot (a) \neq 0$ . Since  $I^2 = I$ ,  $I(I(a)) = I(a) \neq 0$ , so  $I(a) = (a)$ . By NAK,  $(a) = 0$ , contradiction.  $\square$

**Thm.** An Artinian ring has Krull dimension zero. (All primes are maximal.)

*Proof.* Local rings with nilpotent maximal ideal have Krull dimension zero, and, by  $\text{calg1p4}$ ,  $\dim(R \times S) = \max\{\dim(R), \dim(S)\}$ .  $\square$

**Thm.** (Commutative version of Hopkins-Levitski)  
An Artinian ring is Noetherian.

*Proof.* Suffices to prove it for local  $A$ . The finite chain  $A \supset \mathfrak{m} \geq \mathfrak{m}^2 \geq \cdots \geq \mathfrak{m}^k = (0)$  has sections  $\mathfrak{m}^r / \mathfrak{m}^{r+1}$  that are  $A/\mathfrak{m}$ -vector spaces.

# More on the structure of Artinian rings

**Lm.** The maximal ideal of an Artinian local ring is nilpotent.

*Proof.* Assume  $\mathfrak{m} \geq \mathfrak{m}^2 \geq \cdots \geq \mathfrak{m}^k = \mathfrak{m}^{k+1} = I \neq 0$ . Choose  $a \in I$  so that  $(a)$  is minimal for  $I \cdot (a) \neq 0$ . Since  $I^2 = I$ ,  $I(I(a)) = I(a) \neq 0$ , so  $I(a) = (a)$ . By NAK,  $(a) = 0$ , contradiction.  $\square$

**Thm.** An Artinian ring has Krull dimension zero. (All primes are maximal.)

*Proof.* Local rings with nilpotent maximal ideal have Krull dimension zero, and, by  $\text{calg1p4}$ ,  $\dim(R \times S) = \max\{\dim(R), \dim(S)\}$ .  $\square$

**Thm.** (Commutative version of Hopkins-Levitski)  
An Artinian ring is Noetherian.

*Proof.* Suffices to prove it for local  $A$ . The finite chain  $A \supset \mathfrak{m} \geq \mathfrak{m}^2 \geq \cdots \geq \mathfrak{m}^k = (0)$  has sections  $\mathfrak{m}^r / \mathfrak{m}^{r+1}$  that are  $A/\mathfrak{m}$ -vector spaces. Subspace lattices are complemented modular lattices, hence satisfy ACC iff DCC.



# More on the structure of Artinian rings

**Lm.** The maximal ideal of an Artinian local ring is nilpotent.

*Proof.* Assume  $\mathfrak{m} \geq \mathfrak{m}^2 \geq \cdots \geq \mathfrak{m}^k = \mathfrak{m}^{k+1} = I \neq 0$ . Choose  $a \in I$  so that  $(a)$  is minimal for  $I \cdot (a) \neq 0$ . Since  $I^2 = I$ ,  $I(I(a)) = I(a) \neq 0$ , so  $I(a) = (a)$ . By NAK,  $(a) = 0$ , contradiction.  $\square$

**Thm.** An Artinian ring has Krull dimension zero. (All primes are maximal.)

*Proof.* Local rings with nilpotent maximal ideal have Krull dimension zero, and, by  $\text{calg1p4}$ ,  $\dim(R \times S) = \max\{\dim(R), \dim(S)\}$ .  $\square$

**Thm.** (Commutative version of Hopkins-Levitski)  
An Artinian ring is Noetherian.

*Proof.* Suffices to prove it for local  $A$ . The finite chain  $A \supset \mathfrak{m} \geq \mathfrak{m}^2 \geq \cdots \geq \mathfrak{m}^k = (0)$  has sections  $\mathfrak{m}^r / \mathfrak{m}^{r+1}$  that are  $A/\mathfrak{m}$ -vector spaces. Subspace lattices are complemented modular lattices, hence satisfy ACC iff DCC. Hence each section satisfies ACC iff DCC.

# More on the structure of Artinian rings

**Lm.** The maximal ideal of an Artinian local ring is nilpotent.

*Proof.* Assume  $\mathfrak{m} \geq \mathfrak{m}^2 \geq \cdots \geq \mathfrak{m}^k = \mathfrak{m}^{k+1} = I \neq 0$ . Choose  $a \in I$  so that  $(a)$  is minimal for  $I \cdot (a) \neq 0$ . Since  $I^2 = I$ ,  $I(I(a)) = I(a) \neq 0$ , so  $I(a) = (a)$ . By NAK,  $(a) = 0$ , contradiction.  $\square$

**Thm.** An Artinian ring has Krull dimension zero. (All primes are maximal.)

*Proof.* Local rings with nilpotent maximal ideal have Krull dimension zero, and, by  $\text{calg1p4}$ ,  $\dim(R \times S) = \max\{\dim(R), \dim(S)\}$ .  $\square$

**Thm.** (Commutative version of Hopkins-Levitski)  
An Artinian ring is Noetherian.

*Proof.* Suffices to prove it for local  $A$ . The finite chain  $A \supset \mathfrak{m} \geq \mathfrak{m}^2 \geq \cdots \geq \mathfrak{m}^k = (0)$  has sections  $\mathfrak{m}^r/\mathfrak{m}^{r+1}$  that are  $A/\mathfrak{m}$ -vector spaces. Subspace lattices are complemented modular lattices, hence satisfy ACC iff DCC. Hence each section satisfies ACC iff DCC.  $\square$

# More on the structure of Artinian rings

**Thm.**

## More on the structure of Artinian rings

**Thm.** A commutative ring is Artinian iff it is Noetherian of Krull dimension zero.

# More on the structure of Artinian rings

**Thm.** A commutative ring is Artinian iff it is Noetherian of Krull dimension zero.

*Proof.*

# More on the structure of Artinian rings

**Thm.** A commutative ring is Artinian iff it is Noetherian of Krull dimension zero.

*Proof.* Assume  $A$  is Noetherian with Krull dimension zero.

# More on the structure of Artinian rings

**Thm.** A commutative ring is Artinian iff it is Noetherian of Krull dimension zero.

*Proof.* Assume  $A$  is Noetherian with Krull dimension zero. Let  $(0) = Q_1 \cap \cdots \cap Q_k$  be a primary decomposition.



# More on the structure of Artinian rings

**Thm.** A commutative ring is Artinian iff it is Noetherian of Krull dimension zero.

*Proof.* Assume  $A$  is Noetherian with Krull dimension zero. Let  $(0) = Q_1 \cap \cdots \cap Q_k$  be a primary decomposition. Let  $\mathfrak{p}_i = \sqrt{Q_i}$  be the prime belonging to  $Q_i$ .

# More on the structure of Artinian rings

**Thm.** A commutative ring is Artinian iff it is Noetherian of Krull dimension zero.

*Proof.* Assume  $A$  is Noetherian with Krull dimension zero. Let  $(0) = Q_1 \cap \cdots \cap Q_k$  be a primary decomposition. Let  $\mathfrak{p}_i = \sqrt{Q_i}$  be the prime belonging to  $Q_i$ .

$$\mathfrak{N} = \sqrt{(0)} = \sqrt{\cap_{i=1}^k Q_i} = \cap_{i=1}^k \sqrt{Q_i} = \cap_{i=1}^k \mathfrak{p}_i,$$

# More on the structure of Artinian rings

**Thm.** A commutative ring is Artinian iff it is Noetherian of Krull dimension zero.

*Proof.* Assume  $A$  is Noetherian with Krull dimension zero. Let  $(0) = Q_1 \cap \cdots \cap Q_k$  be a primary decomposition. Let  $\mathfrak{p}_i = \sqrt{Q_i}$  be the prime belonging to  $Q_i$ .

$$\mathfrak{N} = \sqrt{(0)} = \sqrt{\cap_{i=1}^k Q_i} = \cap_{i=1}^k \sqrt{Q_i} = \cap_{i=1}^k \mathfrak{p}_i,$$

hence  $\mathfrak{N}$  equals a finite intersection of maximal ideals.

# More on the structure of Artinian rings

**Thm.** A commutative ring is Artinian iff it is Noetherian of Krull dimension zero.

*Proof.* Assume  $A$  is Noetherian with Krull dimension zero. Let  $(0) = Q_1 \cap \cdots \cap Q_k$  be a primary decomposition. Let  $\mathfrak{p}_i = \sqrt{Q_i}$  be the prime belonging to  $Q_i$ .

$$\mathfrak{N} = \sqrt{(0)} = \sqrt{\cap_{i=1}^k Q_i} = \cap_{i=1}^k \sqrt{Q_i} = \cap_{i=1}^k \mathfrak{p}_i,$$

hence  $\mathfrak{N}$  equals a finite intersection of maximal ideals. Consider  $A > \mathfrak{N} \geq \mathfrak{N}^2 \geq \cdots \geq \mathfrak{N}^r = (0)$ ;

# More on the structure of Artinian rings

**Thm.** A commutative ring is Artinian iff it is Noetherian of Krull dimension zero.

*Proof.* Assume  $A$  is Noetherian with Krull dimension zero. Let  $(0) = Q_1 \cap \cdots \cap Q_k$  be a primary decomposition. Let  $\mathfrak{p}_i = \sqrt{Q_i}$  be the prime belonging to  $Q_i$ .

$$\mathfrak{N} = \sqrt{(0)} = \sqrt{\cap_{i=1}^k Q_i} = \cap_{i=1}^k \sqrt{Q_i} = \cap_{i=1}^k \mathfrak{p}_i,$$

hence  $\mathfrak{N}$  equals a finite intersection of maximal ideals. Consider  $A > \mathfrak{N} \geq \mathfrak{N}^2 \geq \cdots \geq \mathfrak{N}^r = (0)$ ; each section  $\mathfrak{N}^i / \mathfrak{N}^{i+1}$  is an  $A/\mathfrak{N}$ -module.

# More on the structure of Artinian rings

**Thm.** A commutative ring is Artinian iff it is Noetherian of Krull dimension zero.

*Proof.* Assume  $A$  is Noetherian with Krull dimension zero. Let  $(0) = Q_1 \cap \cdots \cap Q_k$  be a primary decomposition. Let  $\mathfrak{p}_i = \sqrt{Q_i}$  be the prime belonging to  $Q_i$ .

$$\mathfrak{N} = \sqrt{(0)} = \sqrt{\cap_{i=1}^k Q_i} = \cap_{i=1}^k \sqrt{Q_i} = \cap_{i=1}^k \mathfrak{p}_i,$$

hence  $\mathfrak{N}$  equals a finite intersection of maximal ideals. Consider  $A > \mathfrak{N} \geq \mathfrak{N}^2 \geq \cdots \geq \mathfrak{N}^r = (0)$ ; each section  $\mathfrak{N}^i / \mathfrak{N}^{i+1}$  is an  $A/\mathfrak{N}$ -module. By the CRT,  $A/\mathfrak{N} \cong A/\mathfrak{p}_1 \times \cdots \times A/\mathfrak{p}_k$  is a finite product of fields.

# More on the structure of Artinian rings

**Thm.** A commutative ring is Artinian iff it is Noetherian of Krull dimension zero.

*Proof.* Assume  $A$  is Noetherian with Krull dimension zero. Let  $(0) = Q_1 \cap \cdots \cap Q_k$  be a primary decomposition. Let  $\mathfrak{p}_i = \sqrt{Q_i}$  be the prime belonging to  $Q_i$ .

$$\mathfrak{N} = \sqrt{(0)} = \sqrt{\cap_{i=1}^k Q_i} = \cap_{i=1}^k \sqrt{Q_i} = \cap_{i=1}^k \mathfrak{p}_i,$$

hence  $\mathfrak{N}$  equals a finite intersection of maximal ideals. Consider  $A > \mathfrak{N} \geq \mathfrak{N}^2 \geq \cdots \geq \mathfrak{N}^r = (0)$ ; each section  $\mathfrak{N}^i/\mathfrak{N}^{i+1}$  is an  $A/\mathfrak{N}$ -module. By the CRT,  $A/\mathfrak{N} \cong A/\mathfrak{p}_1 \times \cdots \times A/\mathfrak{p}_k$  is a finite product of fields. Hence each section  $\mathfrak{N}^i/\mathfrak{N}^{i+1}$  has ACC iff DCC.

# More on the structure of Artinian rings

**Thm.** A commutative ring is Artinian iff it is Noetherian of Krull dimension zero.

*Proof.* Assume  $A$  is Noetherian with Krull dimension zero. Let  $(0) = Q_1 \cap \cdots \cap Q_k$  be a primary decomposition. Let  $\mathfrak{p}_i = \sqrt{Q_i}$  be the prime belonging to  $Q_i$ .

$$\mathfrak{N} = \sqrt{(0)} = \sqrt{\cap_{i=1}^k Q_i} = \cap_{i=1}^k \sqrt{Q_i} = \cap_{i=1}^k \mathfrak{p}_i,$$

hence  $\mathfrak{N}$  equals a finite intersection of maximal ideals. Consider  $A > \mathfrak{N} \geq \mathfrak{N}^2 \geq \cdots \geq \mathfrak{N}^r = (0)$ ; each section  $\mathfrak{N}^i/\mathfrak{N}^{i+1}$  is an  $A/\mathfrak{N}$ -module. By the CRT,  $A/\mathfrak{N} \cong A/\mathfrak{p}_1 \times \cdots \times A/\mathfrak{p}_k$  is a finite product of fields. Hence each section  $\mathfrak{N}^i/\mathfrak{N}^{i+1}$  has ACC iff DCC. Hence  $A$  is Artinian.



# More on the structure of Artinian rings

**Thm.** A commutative ring is Artinian iff it is Noetherian of Krull dimension zero.

*Proof.* Assume  $A$  is Noetherian with Krull dimension zero. Let  $(0) = Q_1 \cap \cdots \cap Q_k$  be a primary decomposition. Let  $\mathfrak{p}_i = \sqrt{Q_i}$  be the prime belonging to  $Q_i$ .

$$\mathfrak{N} = \sqrt{(0)} = \sqrt{\cap_{i=1}^k Q_i} = \cap_{i=1}^k \sqrt{Q_i} = \cap_{i=1}^k \mathfrak{p}_i,$$

hence  $\mathfrak{N}$  equals a finite intersection of maximal ideals. Consider  $A > \mathfrak{N} \geq \mathfrak{N}^2 \geq \cdots \geq \mathfrak{N}^r = (0)$ ; each section  $\mathfrak{N}^i/\mathfrak{N}^{i+1}$  is an  $A/\mathfrak{N}$ -module. By the CRT,  $A/\mathfrak{N} \cong A/\mathfrak{p}_1 \times \cdots \times A/\mathfrak{p}_k$  is a finite product of fields. Hence each section  $\mathfrak{N}^i/\mathfrak{N}^{i+1}$  has ACC iff DCC. Hence  $A$  is Artinian.  $\square$

# Commutative Artinian rings

# Commutative Artinian rings

TFAE for a commutative ring  $A$ .

# Commutative Artinian rings

TFAE for a commutative ring  $A$ .

- 1  $A$  is Artinian.

# Commutative Artinian rings

TFAE for a commutative ring  $A$ .

- 1  $A$  is Artinian.

# Commutative Artinian rings

TFAE for a commutative ring  $A$ .

- ①  $A$  is Artinian.
- ②  $A$  is Noetherian and has Krull dimension zero.

# Commutative Artinian rings

TFAE for a commutative ring  $A$ .

- ①  $A$  is Artinian.
- ②  $A$  is Noetherian and has Krull dimension zero.

# Commutative Artinian rings

TFAE for a commutative ring  $A$ .

- 1  $A$  is Artinian.
- 2  $A$  is Noetherian and has Krull dimension zero.
- 3  $A$  is a finite product of Artinian local rings.



# Commutative Artinian rings

TFAE for a commutative ring  $A$ .

- ①  $A$  is Artinian.
- ②  $A$  is Noetherian and has Krull dimension zero.
- ③  $A$  is a finite product of Artinian local rings.

# Commutative Artinian rings

TFAE for a commutative ring  $A$ .

- ①  $A$  is Artinian.
- ②  $A$  is Noetherian and has Krull dimension zero.
- ③  $A$  is a finite product of Artinian local rings.
- ④  $A$  has finitely many prime ideals, all maximal, and their intersection  $\cap \mathfrak{m}_i = J(A)$  equals  $\mathfrak{N}_A$ , which is finitely generated.

# Commutative Artinian rings

TFAE for a commutative ring  $A$ .

- ①  $A$  is Artinian.
- ②  $A$  is Noetherian and has Krull dimension zero.
- ③  $A$  is a finite product of Artinian local rings.
- ④  $A$  has finitely many prime ideals, all maximal, and their intersection  $\cap \mathfrak{m}_i = J(A)$  equals  $\mathfrak{N}_A$ , which is finitely generated.

# Commutative Artinian rings

TFAE for a commutative ring  $A$ .

- ①  $A$  is Artinian.
- ②  $A$  is Noetherian and has Krull dimension zero.
- ③  $A$  is a finite product of Artinian local rings.
- ④  $A$  has finitely many prime ideals, all maximal, and their intersection  $\cap \mathfrak{m}_i = J(A)$  equals  $\mathfrak{N}_A$ , which is finitely generated.

**Examples.** Fields and the finite rings  $\mathbb{Z}_{p^k}$  are Artinian local rings.

# Commutative Artinian rings

TFAE for a commutative ring  $A$ .

- ①  $A$  is Artinian.
- ②  $A$  is Noetherian and has Krull dimension zero.
- ③  $A$  is a finite product of Artinian local rings.
- ④  $A$  has finitely many prime ideals, all maximal, and their intersection  $\cap \mathfrak{m}_i = J(A)$  equals  $\mathfrak{N}_A$ , which is finitely generated.

**Examples.** Fields and the finite rings  $\mathbb{Z}_{p^k}$  are Artinian local rings. A common generalization of  $\mathbb{F}_q = \text{GF}(p, m)$  and  $\mathbb{Z}_{p^k}$  is the Galois ring  $\text{GF}(p^k, m)$ :

# Commutative Artinian rings

TFAE for a commutative ring  $A$ .

- ①  $A$  is Artinian.
- ②  $A$  is Noetherian and has Krull dimension zero.
- ③  $A$  is a finite product of Artinian local rings.
- ④  $A$  has finitely many prime ideals, all maximal, and their intersection  $\cap \mathfrak{m}_i = J(A)$  equals  $\mathfrak{N}_A$ , which is finitely generated.

**Examples.** Fields and the finite rings  $\mathbb{Z}_{p^k}$  are Artinian local rings. A common generalization of  $\mathbb{F}_q = \text{GF}(p, m)$  and  $\mathbb{Z}_{p^k}$  is the Galois ring  $\text{GF}(p^k, m)$ :

$$\text{GF}(p^k, m) = \mathbb{Z}_{p^k}[x]/(f(x))$$

where  $f(x) \in \mathbb{Z}_{p^k}[x]$  is a polynomial of degree  $m$  that is irreducible modulo  $p$ .

# Commutative Artinian rings

TFAE for a commutative ring  $A$ .

- 1  $A$  is Artinian.
- 2  $A$  is Noetherian and has Krull dimension zero.
- 3  $A$  is a finite product of Artinian local rings.
- 4  $A$  has finitely many prime ideals, all maximal, and their intersection  $\cap \mathfrak{m}_i = J(A)$  equals  $\mathfrak{N}_A$ , which is finitely generated.

**Examples.** Fields and the finite rings  $\mathbb{Z}_{p^k}$  are Artinian local rings. A common generalization of  $\mathbb{F}_q = \text{GF}(p, m)$  and  $\mathbb{Z}_{p^k}$  is the Galois ring  $\text{GF}(p^k, m)$ :

$$\text{GF}(p^k, m) = \mathbb{Z}_{p^k}[x]/(f(x))$$

where  $f(x) \in \mathbb{Z}_{p^k}[x]$  is a polynomial of degree  $m$  that is irreducible modulo  $p$ . (It is not obvious that this data determines the ring uniquely, but it does.)

# Commutative Artinian rings

TFAE for a commutative ring  $A$ .

- ①  $A$  is Artinian.
- ②  $A$  is Noetherian and has Krull dimension zero.
- ③  $A$  is a finite product of Artinian local rings.
- ④  $A$  has finitely many prime ideals, all maximal, and their intersection  $\cap \mathfrak{m}_i = J(A)$  equals  $\mathfrak{N}_A$ , which is finitely generated.

**Examples.** Fields and the finite rings  $\mathbb{Z}_{p^k}$  are Artinian local rings. A common generalization of  $\mathbb{F}_q = \text{GF}(p, m)$  and  $\mathbb{Z}_{p^k}$  is the Galois ring  $\text{GF}(p^k, m)$ :

$$\text{GF}(p^k, m) = \mathbb{Z}_{p^k}[x]/(f(x))$$

where  $f(x) \in \mathbb{Z}_{p^k}[x]$  is a polynomial of degree  $m$  that is irreducible modulo  $p$ . (It is not obvious that this data determines the ring uniquely, but it does.) The smallest nontrivial example is  $\mathbb{Z}_4[x]/(x^2 + x + 1)$ .



# $\text{Spec}(A)$ when $A$ is Noetherian

# $\text{Spec}(A)$ when $A$ is Noetherian

**Kaplansky's Question.**

# $\text{Spec}(A)$ when $A$ is Noetherian

**Kaplansky's Question.** ( $\sim 1950$ )

# $\text{Spec}(A)$ when $A$ is Noetherian

**Kaplansky's Question.** ( $\sim 1950$ ) Which topological spaces arise as  $\text{Spec}(A)$  for  $A$  a commutative Noetherian ring?

# $\text{Spec}(A)$ when $A$ is Noetherian

**Kaplansky's Question.** ( $\sim 1950$ ) Which topological spaces arise as  $\text{Spec}(A)$  for  $A$  a commutative Noetherian ring?

Known properties:

# $\text{Spec}(A)$ when $A$ is Noetherian

**Kaplansky's Question.** ( $\sim 1950$ ) Which topological spaces arise as  $\text{Spec}(A)$  for  $A$  a commutative Noetherian ring?

Known properties:

- ① The topology on the set of primes and the inclusion order on the set of primes mutually determine one another.

# $\text{Spec}(A)$ when $A$ is Noetherian

**Kaplansky's Question.** ( $\sim 1950$ ) Which topological spaces arise as  $\text{Spec}(A)$  for  $A$  a commutative Noetherian ring?

Known properties:

- ① The topology on the set of primes and the inclusion order on the set of primes mutually determine one another.

# $\text{Spec}(A)$ when $A$ is Noetherian

**Kaplansky's Question.** ( $\sim 1950$ ) Which topological spaces arise as  $\text{Spec}(A)$  for  $A$  a commutative Noetherian ring?

Known properties:

- ① The topology on the set of primes and the inclusion order on the set of primes mutually determine one another. (calg2p6)



# $\text{Spec}(A)$ when $A$ is Noetherian

**Kaplansky's Question.** ( $\sim 1950$ ) Which topological spaces arise as  $\text{Spec}(A)$  for  $A$  a commutative Noetherian ring?

Known properties:

- 1 The topology on the set of primes and the inclusion order on the set of primes mutually determine one another. (calg2p6)
- 2  $P = \langle \text{Spec}(A); \subseteq \rangle$  has the following properties:

# $\text{Spec}(A)$ when $A$ is Noetherian

**Kaplansky's Question.** ( $\sim 1950$ ) Which topological spaces arise as  $\text{Spec}(A)$  for  $A$  a commutative Noetherian ring?

Known properties:

- ① The topology on the set of primes and the inclusion order on the set of primes mutually determine one another. (calg2p6)
- ②  $P = \langle \text{Spec}(A); \subseteq \rangle$  has the following properties:
  - ①  $P$  has finitely many minimal elements

# $\text{Spec}(A)$ when $A$ is Noetherian

**Kaplansky's Question.** ( $\sim 1950$ ) Which topological spaces arise as  $\text{Spec}(A)$  for  $A$  a commutative Noetherian ring?

Known properties:

- ① The topology on the set of primes and the inclusion order on the set of primes mutually determine one another. (calg2p6)
- ②  $P = \langle \text{Spec}(A); \subseteq \rangle$  has the following properties:
  - ①  $P$  has finitely many minimal elements

# $\text{Spec}(A)$ when $A$ is Noetherian

**Kaplansky's Question.** ( $\sim 1950$ ) Which topological spaces arise as  $\text{Spec}(A)$  for  $A$  a commutative Noetherian ring?

Known properties:

- ① The topology on the set of primes and the inclusion order on the set of primes mutually determine one another. (calg2p6)
- ②  $P = \langle \text{Spec}(A); \subseteq \rangle$  has the following properties:
  - ①  $P$  has finitely many minimal elements (calg2p5)

# $\text{Spec}(A)$ when $A$ is Noetherian

**Kaplansky's Question.** ( $\sim 1950$ ) Which topological spaces arise as  $\text{Spec}(A)$  for  $A$  a commutative Noetherian ring?

Known properties:

- ① The topology on the set of primes and the inclusion order on the set of primes mutually determine one another. (calg2p6)
- ②  $P = \langle \text{Spec}(A); \subseteq \rangle$  has the following properties:
  - ①  $P$  has finitely many minimal elements (calg2p5)
  - ②  $P$  satisfies ACC

# $\text{Spec}(A)$ when $A$ is Noetherian

**Kaplansky's Question.** ( $\sim 1950$ ) Which topological spaces arise as  $\text{Spec}(A)$  for  $A$  a commutative Noetherian ring?

Known properties:

- ① The topology on the set of primes and the inclusion order on the set of primes mutually determine one another. (calg2p6)
- ②  $P = \langle \text{Spec}(A); \subseteq \rangle$  has the following properties:
  - ①  $P$  has finitely many minimal elements (calg2p5)
  - ②  $P$  satisfies ACC

# $\text{Spec}(A)$ when $A$ is Noetherian

**Kaplansky's Question.** ( $\sim 1950$ ) Which topological spaces arise as  $\text{Spec}(A)$  for  $A$  a commutative Noetherian ring?

Known properties:

- ① The topology on the set of primes and the inclusion order on the set of primes mutually determine one another. (calg2p6)
- ②  $P = \langle \text{Spec}(A); \subseteq \rangle$  has the following properties:
  - ①  $P$  has finitely many minimal elements (calg2p5)
  - ②  $P$  satisfies ACC (!)

# $\text{Spec}(A)$ when $A$ is Noetherian

**Kaplansky's Question.** ( $\sim 1950$ ) Which topological spaces arise as  $\text{Spec}(A)$  for  $A$  a commutative Noetherian ring?

Known properties:

- ① The topology on the set of primes and the inclusion order on the set of primes mutually determine one another. (calg2p6)
- ②  $P = \langle \text{Spec}(A); \subseteq \rangle$  has the following properties:
  - ①  $P$  has finitely many minimal elements (calg2p5)
  - ②  $P$  satisfies ACC (!)
  - ③  $P$  satisfies DCC



# $\text{Spec}(A)$ when $A$ is Noetherian

**Kaplansky's Question.** ( $\sim 1950$ ) Which topological spaces arise as  $\text{Spec}(A)$  for  $A$  a commutative Noetherian ring?

Known properties:

- ① The topology on the set of primes and the inclusion order on the set of primes mutually determine one another. (calg2p6)
- ②  $P = \langle \text{Spec}(A); \subseteq \rangle$  has the following properties:
  - ①  $P$  has finitely many minimal elements (calg2p5)
  - ②  $P$  satisfies ACC (!)
  - ③  $P$  satisfies DCC

# $\text{Spec}(A)$ when $A$ is Noetherian

**Kaplansky's Question.** ( $\sim 1950$ ) Which topological spaces arise as  $\text{Spec}(A)$  for  $A$  a commutative Noetherian ring?

Known properties:

- ① The topology on the set of primes and the inclusion order on the set of primes mutually determine one another. (calg2p6)
- ②  $P = \langle \text{Spec}(A); \subseteq \rangle$  has the following properties:
  - ①  $P$  has finitely many minimal elements (calg2p5)
  - ②  $P$  satisfies ACC (!)
  - ③  $P$  satisfies DCC (from the Krull Height Theorem)

# $\text{Spec}(A)$ when $A$ is Noetherian

**Kaplansky's Question.** ( $\sim 1950$ ) Which topological spaces arise as  $\text{Spec}(A)$  for  $A$  a commutative Noetherian ring?

Known properties:

- ① The topology on the set of primes and the inclusion order on the set of primes mutually determine one another. (calg2p6)
- ②  $P = \langle \text{Spec}(A); \subseteq \rangle$  has the following properties:
  - ①  $P$  has finitely many minimal elements (calg2p5)
  - ②  $P$  satisfies ACC (!)
  - ③  $P$  satisfies DCC (from the Krull Height Theorem)
  - ④ The class  $\mathcal{F}$  of finitely generated order-filters is closed under arbitrary intersection.

# $\text{Spec}(A)$ when $A$ is Noetherian

**Kaplansky's Question.** ( $\sim 1950$ ) Which topological spaces arise as  $\text{Spec}(A)$  for  $A$  a commutative Noetherian ring?

Known properties:

- ① The topology on the set of primes and the inclusion order on the set of primes mutually determine one another. (calg2p6)
- ②  $P = \langle \text{Spec}(A); \subseteq \rangle$  has the following properties:
  - ①  $P$  has finitely many minimal elements (calg2p5)
  - ②  $P$  satisfies ACC (!)
  - ③  $P$  satisfies DCC (from the Krull Height Theorem)
  - ④ The class  $\mathcal{F}$  of finitely generated order-filters is closed under arbitrary intersection.

# $\text{Spec}(A)$ when $A$ is Noetherian

**Kaplansky's Question.** ( $\sim 1950$ ) Which topological spaces arise as  $\text{Spec}(A)$  for  $A$  a commutative Noetherian ring?

Known properties:

- ① The topology on the set of primes and the inclusion order on the set of primes mutually determine one another. (calg2p6)
- ②  $P = \langle \text{Spec}(A); \subseteq \rangle$  has the following properties:
  - ①  $P$  has finitely many minimal elements (calg2p5)
  - ②  $P$  satisfies ACC (!)
  - ③  $P$  satisfies DCC (from the Krull Height Theorem)
  - ④ The class  $\mathcal{F}$  of finitely generated order-filters is closed under arbitrary intersection. (calg2p6)

# $\text{Spec}(A)$ when $A$ is Noetherian

**Kaplansky's Question.** ( $\sim 1950$ ) Which topological spaces arise as  $\text{Spec}(A)$  for  $A$  a commutative Noetherian ring?

Known properties:

- ① The topology on the set of primes and the inclusion order on the set of primes mutually determine one another. (calg2p6)
- ②  $P = \langle \text{Spec}(A); \subseteq \rangle$  has the following properties:
  - ①  $P$  has finitely many minimal elements (calg2p5)
  - ②  $P$  satisfies ACC (!)
  - ③  $P$  satisfies DCC (from the Krull Height Theorem)
  - ④ The class  $\mathcal{F}$  of finitely generated order-filters is closed under arbitrary intersection. (calg2p6)
  - ⑤ Nontrivial noncovering intervals  $[\mathfrak{p}, \mathfrak{r}]$  are infinite.

# $\text{Spec}(A)$ when $A$ is Noetherian

**Kaplansky's Question.** ( $\sim 1950$ ) Which topological spaces arise as  $\text{Spec}(A)$  for  $A$  a commutative Noetherian ring?

Known properties:

- ① The topology on the set of primes and the inclusion order on the set of primes mutually determine one another. (calg2p6)
- ②  $P = \langle \text{Spec}(A); \subseteq \rangle$  has the following properties:
  - ①  $P$  has finitely many minimal elements (calg2p5)
  - ②  $P$  satisfies ACC (!)
  - ③  $P$  satisfies DCC (from the Krull Height Theorem)
  - ④ The class  $\mathcal{F}$  of finitely generated order-filters is closed under arbitrary intersection. (calg2p6)
  - ⑤ Nontrivial noncovering intervals  $[\mathfrak{p}, \mathfrak{r}]$  are infinite.

# $\text{Spec}(A)$ when $A$ is Noetherian

**Kaplansky's Question.** ( $\sim 1950$ ) Which topological spaces arise as  $\text{Spec}(A)$  for  $A$  a commutative Noetherian ring?

Known properties:

- ① The topology on the set of primes and the inclusion order on the set of primes mutually determine one another. (calg2p6)
- ②  $P = \langle \text{Spec}(A); \subseteq \rangle$  has the following properties:
  - ①  $P$  has finitely many minimal elements (calg2p5)
  - ②  $P$  satisfies ACC (!)
  - ③  $P$  satisfies DCC (from the Krull Height Theorem)
  - ④ The class  $\mathcal{F}$  of finitely generated order-filters is closed under arbitrary intersection. (calg2p6)
  - ⑤ Nontrivial noncovering intervals  $[\mathfrak{p}, \mathfrak{r}]$  are infinite. (If  $\mathfrak{p} < \mathfrak{q} < \mathfrak{r}$ , then there exist infinitely many distinct  $\mathfrak{q}_i$  such that  $\mathfrak{p} < \mathfrak{q}_i < \mathfrak{r}$ .)



# $\text{Spec}(A)$ when $A$ is Noetherian

**Kaplansky's Question.** ( $\sim 1950$ ) Which topological spaces arise as  $\text{Spec}(A)$  for  $A$  a commutative Noetherian ring?

Known properties:

- ① The topology on the set of primes and the inclusion order on the set of primes mutually determine one another. (calg2p6)
- ②  $P = \langle \text{Spec}(A); \subseteq \rangle$  has the following properties:
  - ①  $P$  has finitely many minimal elements (calg2p5)
  - ②  $P$  satisfies ACC (!)
  - ③  $P$  satisfies DCC (from the Krull Height Theorem)
  - ④ The class  $\mathcal{F}$  of finitely generated order-filters is closed under arbitrary intersection. (calg2p6)
  - ⑤ Nontrivial noncovering intervals  $[\mathfrak{p}, \mathfrak{r}]$  are infinite. (If  $\mathfrak{p} < \mathfrak{q} < \mathfrak{r}$ , then there exist infinitely many distinct  $\mathfrak{q}_i$  such that  $\mathfrak{p} < \mathfrak{q}_i < \mathfrak{r}$ .) (from Prime Avoidance+Krull Height)

# $\text{Spec}(A)$ when $A$ is Noetherian

**Kaplansky's Question.** ( $\sim 1950$ ) Which topological spaces arise as  $\text{Spec}(A)$  for  $A$  a commutative Noetherian ring?

Known properties:

- ① The topology on the set of primes and the inclusion order on the set of primes mutually determine one another. (calg2p6)
- ②  $P = \langle \text{Spec}(A); \subseteq \rangle$  has the following properties:
  - ①  $P$  has finitely many minimal elements (calg2p5)
  - ②  $P$  satisfies ACC (!)
  - ③  $P$  satisfies DCC (from the Krull Height Theorem)
  - ④ The class  $\mathcal{F}$  of finitely generated order-filters is closed under arbitrary intersection. (calg2p6)
  - ⑤ Nontrivial noncovering intervals  $[\mathfrak{p}, \mathfrak{r}]$  are infinite. (If  $\mathfrak{p} < \mathfrak{q} < \mathfrak{r}$ , then there exist infinitely many distinct  $\mathfrak{q}_i$  such that  $\mathfrak{p} < \mathfrak{q}_i < \mathfrak{r}$ .) (from Prime Avoidance+Krull Height)

**Krull Height Theorem.** If  $A$  is a commutative Noetherian ring and  $I \triangleleft A$  can be generated by  $n$  elements, then any prime minimal over  $I$  has height at most  $n$ . (In particular, every prime has finite height.)



## **Prime Avoidance Theorem.**

**Prime Avoidance Theorem.** If

**Prime Avoidance Theorem.** If

①  $I_1, \dots, I_n, J$  are ideals,

**Prime Avoidance Theorem.** If

①  $I_1, \dots, I_n, J$  are ideals,

**Prime Avoidance Theorem.** If

- ①  $I_1, \dots, I_n, J$  are ideals,
- ②  $I_j$  is prime for  $j \geq 3$ , and



**Prime Avoidance Theorem.** If

- ①  $I_1, \dots, I_n, J$  are ideals,
- ②  $I_j$  is prime for  $j \geq 3$ , and

**Prime Avoidance Theorem.** If

- ①  $I_1, \dots, I_n, J$  are ideals,
- ②  $I_j$  is prime for  $j \geq 3$ , and
- ③  $J \subseteq \cup_{i=1}^n I_i$ ,

**Prime Avoidance Theorem.** If

- ①  $I_1, \dots, I_n, J$  are ideals,
- ②  $I_j$  is prime for  $j \geq 3$ , and
- ③  $J \subseteq \cup_{i=1}^n I_i$ ,

**Prime Avoidance Theorem.** If

- ①  $I_1, \dots, I_n, J$  are ideals,
- ②  $I_j$  is prime for  $j \geq 3$ , and
- ③  $J \subseteq \cup_{i=1}^n I_i$ ,

then  $J \subseteq I_j$  for some  $j$ .

**Prime Avoidance Theorem.** If

- ①  $I_1, \dots, I_n, J$  are ideals,
- ②  $I_j$  is prime for  $j \geq 3$ , and
- ③  $J \subseteq \cup_{i=1}^n I_i$ ,

then  $J \subseteq I_j$  for some  $j$ .

*Proof.*

**Prime Avoidance Theorem.** If

- ①  $I_1, \dots, I_n, J$  are ideals,
- ②  $I_j$  is prime for  $j \geq 3$ , and
- ③  $J \subseteq \cup_{i=1}^n I_i$ ,

then  $J \subseteq I_j$  for some  $j$ .

*Proof.* Induction on  $n$ .

**Prime Avoidance Theorem.** If

- ①  $I_1, \dots, I_n, J$  are ideals,
- ②  $I_j$  is prime for  $j \geq 3$ , and
- ③  $J \subseteq \cup_{i=1}^n I_i$ ,

then  $J \subseteq I_j$  for some  $j$ .

*Proof.* Induction on  $n$ . The  $n = 1$  case is clear.

**Prime Avoidance Theorem.** If

- ①  $I_1, \dots, I_n, J$  are ideals,
- ②  $I_j$  is prime for  $j \geq 3$ , and
- ③  $J \subseteq \cup_{i=1}^n I_i$ ,

then  $J \subseteq I_j$  for some  $j$ .

*Proof.* Induction on  $n$ . The  $n = 1$  case is clear. For  $n = 2$ , choose  $x_1 \in J - I_2 \subseteq I_1$  and  $x_2 \in J - I_1 \subseteq I_2$ .



**Prime Avoidance Theorem.** If

- ①  $I_1, \dots, I_n, J$  are ideals,
- ②  $I_j$  is prime for  $j \geq 3$ , and
- ③  $J \subseteq \cup_{i=1}^n I_i$ ,

then  $J \subseteq I_j$  for some  $j$ .

*Proof.* Induction on  $n$ . The  $n = 1$  case is clear. For  $n = 2$ , choose  $x_1 \in J - I_2 \subseteq I_1$  and  $x_2 \in J - I_1 \subseteq I_2$ . Hence  $x_1 + x_2 \in J - (I_1 \cup I_2)$ ,

**Prime Avoidance Theorem.** If

- ①  $I_1, \dots, I_n, J$  are ideals,
- ②  $I_j$  is prime for  $j \geq 3$ , and
- ③  $J \subseteq \cup_{i=1}^n I_i$ ,

then  $J \subseteq I_j$  for some  $j$ .

*Proof.* Induction on  $n$ . The  $n = 1$  case is clear. For  $n = 2$ , choose  $x_1 \in J - I_2 \subseteq I_1$  and  $x_2 \in J - I_1 \subseteq I_2$ . Hence  $x_1 + x_2 \in J - (I_1 \cup I_2)$ , contradiction.

**Prime Avoidance Theorem.** If

- ①  $I_1, \dots, I_n, J$  are ideals,
- ②  $I_j$  is prime for  $j \geq 3$ , and
- ③  $J \subseteq \cup_{i=1}^n I_i$ ,

then  $J \subseteq I_j$  for some  $j$ .

*Proof.* Induction on  $n$ . The  $n = 1$  case is clear. For  $n = 2$ , choose  $x_1 \in J - I_2 \subseteq I_1$  and  $x_2 \in J - I_1 \subseteq I_2$ . Hence  $x_1 + x_2 \in J - (I_1 \cup I_2)$ , contradiction. Now assume  $n \geq 3$ .

**Prime Avoidance Theorem.** If

- ①  $I_1, \dots, I_n, J$  are ideals,
- ②  $I_j$  is prime for  $j \geq 3$ , and
- ③  $J \subseteq \cup_{i=1}^n I_i$ ,

then  $J \subseteq I_j$  for some  $j$ .

*Proof.* Induction on  $n$ . The  $n = 1$  case is clear. For  $n = 2$ , choose  $x_1 \in J - I_2 \subseteq I_1$  and  $x_2 \in J - I_1 \subseteq I_2$ . Hence  $x_1 + x_2 \in J - (I_1 \cup I_2)$ , contradiction. Now assume  $n \geq 3$ . Choose  $x_j \in (J - \cup_{i \neq j} I_i)$ .

**Prime Avoidance Theorem.** If

- ①  $I_1, \dots, I_n, J$  are ideals,
- ②  $I_j$  is prime for  $j \geq 3$ , and
- ③  $J \subseteq \cup_{i=1}^n I_i$ ,

then  $J \subseteq I_j$  for some  $j$ .

*Proof.* Induction on  $n$ . The  $n = 1$  case is clear. For  $n = 2$ , choose  $x_1 \in J - I_2 \subseteq I_1$  and  $x_2 \in J - I_1 \subseteq I_2$ . Hence  $x_1 + x_2 \in J - (I_1 \cup I_2)$ , contradiction. Now assume  $n \geq 3$ . Choose  $x_j \in (J - \cup_{i \neq j} I_i)$ . Necessarily  $x_j \in I_j$ .

**Prime Avoidance Theorem.** If

- ①  $I_1, \dots, I_n, J$  are ideals,
- ②  $I_j$  is prime for  $j \geq 3$ , and
- ③  $J \subseteq \cup_{i=1}^n I_i$ ,

then  $J \subseteq I_j$  for some  $j$ .

*Proof.* Induction on  $n$ . The  $n = 1$  case is clear. For  $n = 2$ , choose  $x_1 \in J - I_2 \subseteq I_1$  and  $x_2 \in J - I_1 \subseteq I_2$ . Hence  $x_1 + x_2 \in J - (I_1 \cup I_2)$ , contradiction. Now assume  $n \geq 3$ . Choose  $x_j \in (J - \cup_{i \neq j} I_i)$ . Necessarily  $x_j \in I_j$ . Now  $(x_1 \cdots x_{n-1}) + x_n$  belongs to  $J$  but to no  $I_j$ .

**Prime Avoidance Theorem.** If

- ①  $I_1, \dots, I_n, J$  are ideals,
- ②  $I_j$  is prime for  $j \geq 3$ , and
- ③  $J \subseteq \cup_{i=1}^n I_i$ ,

then  $J \subseteq I_j$  for some  $j$ .

*Proof.* Induction on  $n$ . The  $n = 1$  case is clear. For  $n = 2$ , choose  $x_1 \in J - I_2 \subseteq I_1$  and  $x_2 \in J - I_1 \subseteq I_2$ . Hence  $x_1 + x_2 \in J - (I_1 \cup I_2)$ , contradiction. Now assume  $n \geq 3$ . Choose  $x_j \in (J - \cup_{j \neq i} I_i)$ . Necessarily  $x_j \in I_j$ . Now  $(x_1 \cdots x_{n-1}) + x_n$  belongs to  $J$  but to no  $I_j$ , contradiction.

**Prime Avoidance Theorem.** If

- ①  $I_1, \dots, I_n, J$  are ideals,
- ②  $I_j$  is prime for  $j \geq 3$ , and
- ③  $J \subseteq \bigcup_{i=1}^n I_i$ ,

then  $J \subseteq I_j$  for some  $j$ .

*Proof.* Induction on  $n$ . The  $n = 1$  case is clear. For  $n = 2$ , choose  $x_1 \in J - I_2 \subseteq I_1$  and  $x_2 \in J - I_1 \subseteq I_2$ . Hence  $x_1 + x_2 \in J - (I_1 \cup I_2)$ , contradiction. Now assume  $n \geq 3$ . Choose  $x_j \in (J - \bigcup_{i \neq j} I_i)$ . Necessarily  $x_j \in I_j$ . Now  $(x_1 \cdots x_{n-1}) + x_n$  belongs to  $J$  but to no  $I_j$ , contradiction.  $\square$



**Prime Avoidance Theorem.** If

- ①  $I_1, \dots, I_n, J$  are ideals,
- ②  $I_j$  is prime for  $j \geq 3$ , and
- ③  $J \subseteq \cup_{i=1}^n I_i$ ,

then  $J \subseteq I_j$  for some  $j$ .

*Proof.* Induction on  $n$ . The  $n = 1$  case is clear. For  $n = 2$ , choose  $x_1 \in J - I_2 \subseteq I_1$  and  $x_2 \in J - I_1 \subseteq I_2$ . Hence  $x_1 + x_2 \in J - (I_1 \cup I_2)$ , contradiction. Now assume  $n \geq 3$ . Choose  $x_j \in (J - \cup_{j \neq i} I_i)$ . Necessarily  $x_j \in I_j$ . Now  $(x_1 \cdots x_{n-1}) + x_n$  belongs to  $J$  but to no  $I_j$ , contradiction.  $\square$

**Application.** Let  $A$  be a commutative Noetherian ring. If  $\mathfrak{p} < \mathfrak{q} < \mathfrak{r}$  are prime, then there exist infinitely many distinct primes  $\mathfrak{q}_i$  such that  $\mathfrak{p} < \mathfrak{q}_i < \mathfrak{r}$ .

**Prime Avoidance Theorem.** If

- ①  $I_1, \dots, I_n, J$  are ideals,
- ②  $I_j$  is prime for  $j \geq 3$ , and
- ③  $J \subseteq \cup_{i=1}^n I_i$ ,

then  $J \subseteq I_j$  for some  $j$ .

*Proof.* Induction on  $n$ . The  $n = 1$  case is clear. For  $n = 2$ , choose  $x_1 \in J - I_2 \subseteq I_1$  and  $x_2 \in J - I_1 \subseteq I_2$ . Hence  $x_1 + x_2 \in J - (I_1 \cup I_2)$ , contradiction. Now assume  $n \geq 3$ . Choose  $x_j \in (J - \cup_{j \neq i} I_i)$ . Necessarily  $x_j \in I_j$ . Now  $(x_1 \cdots x_{n-1}) + x_n$  belongs to  $J$  but to no  $I_j$ , contradiction.  $\square$

**Application.** Let  $A$  be a commutative Noetherian ring. If  $\mathfrak{p} < \mathfrak{q} < \mathfrak{r}$  are prime, then there exist infinitely many distinct primes  $\mathfrak{q}_i$  such that  $\mathfrak{p} < \mathfrak{q}_i < \mathfrak{r}$ .

*Proof.* Shrink interval so that  $[\mathfrak{p}, \mathfrak{r}]$  has height 2.

**Prime Avoidance Theorem.** If

- ①  $I_1, \dots, I_n, J$  are ideals,
- ②  $I_j$  is prime for  $j \geq 3$ , and
- ③  $J \subseteq \cup_{i=1}^n I_i$ ,

then  $J \subseteq I_j$  for some  $j$ .

*Proof.* Induction on  $n$ . The  $n = 1$  case is clear. For  $n = 2$ , choose  $x_1 \in J - I_2 \subseteq I_1$  and  $x_2 \in J - I_1 \subseteq I_2$ . Hence  $x_1 + x_2 \in J - (I_1 \cup I_2)$ , contradiction. Now assume  $n \geq 3$ . Choose  $x_j \in (J - \cup_{j \neq i} I_i)$ . Necessarily  $x_j \in I_j$ . Now  $(x_1 \cdots x_{n-1}) + x_n$  belongs to  $J$  but to no  $I_j$ , contradiction.  $\square$

**Application.** Let  $A$  be a commutative Noetherian ring. If  $\mathfrak{p} < \mathfrak{q} < \mathfrak{r}$  are prime, then there exist infinitely many distinct primes  $\mathfrak{q}_i$  such that  $\mathfrak{p} < \mathfrak{q}_i < \mathfrak{r}$ .

*Proof.* Shrink interval so that  $[\mathfrak{p}, \mathfrak{r}]$  has height 2. May assume that  $\mathfrak{p} = (0)$

**Prime Avoidance Theorem.** If

- ①  $I_1, \dots, I_n, J$  are ideals,
- ②  $I_j$  is prime for  $j \geq 3$ , and
- ③  $J \subseteq \bigcup_{i=1}^n I_i$ ,

then  $J \subseteq I_j$  for some  $j$ .

*Proof.* Induction on  $n$ . The  $n = 1$  case is clear. For  $n = 2$ , choose  $x_1 \in J - I_2 \subseteq I_1$  and  $x_2 \in J - I_1 \subseteq I_2$ . Hence  $x_1 + x_2 \in J - (I_1 \cup I_2)$ , contradiction. Now assume  $n \geq 3$ . Choose  $x_j \in (J - \bigcup_{i \neq j} I_i)$ . Necessarily  $x_j \in I_j$ . Now  $(x_1 \cdots x_{n-1}) + x_n$  belongs to  $J$  but to no  $I_j$ , contradiction.  $\square$

**Application.** Let  $A$  be a commutative Noetherian ring. If  $\mathfrak{p} < \mathfrak{q} < \mathfrak{r}$  are prime, then there exist infinitely many distinct primes  $\mathfrak{q}_i$  such that  $\mathfrak{p} < \mathfrak{q}_i < \mathfrak{r}$ .

*Proof.* Shrink interval so that  $[\mathfrak{p}, \mathfrak{r}]$  has height 2. May assume that  $\mathfrak{p} = (0)$  and that  $\mathfrak{r}$  is maximal.

**Prime Avoidance Theorem.** If

- ①  $I_1, \dots, I_n, J$  are ideals,
- ②  $I_j$  is prime for  $j \geq 3$ , and
- ③  $J \subseteq \bigcup_{i=1}^n I_i$ ,

then  $J \subseteq I_j$  for some  $j$ .

*Proof.* Induction on  $n$ . The  $n = 1$  case is clear. For  $n = 2$ , choose  $x_1 \in J - I_2 \subseteq I_1$  and  $x_2 \in J - I_1 \subseteq I_2$ . Hence  $x_1 + x_2 \in J - (I_1 \cup I_2)$ , contradiction. Now assume  $n \geq 3$ . Choose  $x_j \in (J - \bigcup_{i \neq j} I_i)$ . Necessarily  $x_j \in I_j$ . Now  $(x_1 \cdots x_{n-1}) + x_n$  belongs to  $J$  but to no  $I_j$ , contradiction.  $\square$

**Application.** Let  $A$  be a commutative Noetherian ring. If  $\mathfrak{p} < \mathfrak{q} < \mathfrak{r}$  are prime, then there exist infinitely many distinct primes  $\mathfrak{q}_i$  such that  $\mathfrak{p} < \mathfrak{q}_i < \mathfrak{r}$ .

*Proof.* Shrink interval so that  $[\mathfrak{p}, \mathfrak{r}]$  has height 2. May assume that  $\mathfrak{p} = (0)$  and that  $\mathfrak{r}$  is maximal. If only finitely many height-1 primes exist,  $\mathfrak{q}_1, \dots, \mathfrak{q}_n$ , then choose  $a \in \mathfrak{r} - (\bigcup_{i=1}^n \mathfrak{q}_i)$ .

**Prime Avoidance Theorem.** If

- ①  $I_1, \dots, I_n, J$  are ideals,
- ②  $I_j$  is prime for  $j \geq 3$ , and
- ③  $J \subseteq \bigcup_{i=1}^n I_i$ ,

then  $J \subseteq I_j$  for some  $j$ .

*Proof.* Induction on  $n$ . The  $n = 1$  case is clear. For  $n = 2$ , choose  $x_1 \in J - I_2 \subseteq I_1$  and  $x_2 \in J - I_1 \subseteq I_2$ . Hence  $x_1 + x_2 \in J - (I_1 \cup I_2)$ , contradiction. Now assume  $n \geq 3$ . Choose  $x_j \in (J - \bigcup_{i \neq j} I_i)$ . Necessarily  $x_j \in I_j$ . Now  $(x_1 \cdots x_{n-1}) + x_n$  belongs to  $J$  but to no  $I_j$ , contradiction.  $\square$

**Application.** Let  $A$  be a commutative Noetherian ring. If  $\mathfrak{p} < \mathfrak{q} < \mathfrak{r}$  are prime, then there exist infinitely many distinct primes  $\mathfrak{q}_i$  such that  $\mathfrak{p} < \mathfrak{q}_i < \mathfrak{r}$ .

*Proof.* Shrink interval so that  $[\mathfrak{p}, \mathfrak{r}]$  has height 2. May assume that  $\mathfrak{p} = (0)$  and that  $\mathfrak{r}$  is maximal. If only finitely many height-1 primes exist,  $\mathfrak{q}_1, \dots, \mathfrak{q}_n$ , then choose  $a \in \mathfrak{r} - (\bigcup_{i=1}^n \mathfrak{q}_i)$ . Any prime minimal over  $(a)$  has height 1 by Krull Height Theorem

**Prime Avoidance Theorem.** If

- ①  $I_1, \dots, I_n, J$  are ideals,
- ②  $I_j$  is prime for  $j \geq 3$ , and
- ③  $J \subseteq \bigcup_{i=1}^n I_i$ ,

then  $J \subseteq I_j$  for some  $j$ .

*Proof.* Induction on  $n$ . The  $n = 1$  case is clear. For  $n = 2$ , choose  $x_1 \in J - I_2 \subseteq I_1$  and  $x_2 \in J - I_1 \subseteq I_2$ . Hence  $x_1 + x_2 \in J - (I_1 \cup I_2)$ , contradiction. Now assume  $n \geq 3$ . Choose  $x_j \in (J - \bigcup_{i \neq j} I_i)$ . Necessarily  $x_j \in I_j$ . Now  $(x_1 \cdots x_{n-1}) + x_n$  belongs to  $J$  but to no  $I_j$ , contradiction.  $\square$

**Application.** Let  $A$  be a commutative Noetherian ring. If  $\mathfrak{p} < \mathfrak{q} < \mathfrak{r}$  are prime, then there exist infinitely many distinct primes  $\mathfrak{q}_i$  such that  $\mathfrak{p} < \mathfrak{q}_i < \mathfrak{r}$ .

*Proof.* Shrink interval so that  $[\mathfrak{p}, \mathfrak{r}]$  has height 2. May assume that  $\mathfrak{p} = (0)$  and that  $\mathfrak{r}$  is maximal. If only finitely many height-1 primes exist,  $\mathfrak{q}_1, \dots, \mathfrak{q}_n$ , then choose  $a \in \mathfrak{r} - (\bigcup_{i=1}^n \mathfrak{q}_i)$ . Any prime minimal over  $(a)$  has height 1 by Krull Height Theorem (needs Noetherianness),

**Prime Avoidance Theorem.** If

- ①  $I_1, \dots, I_n, J$  are ideals,
- ②  $I_j$  is prime for  $j \geq 3$ , and
- ③  $J \subseteq \bigcup_{i=1}^n I_i$ ,

then  $J \subseteq I_j$  for some  $j$ .

*Proof.* Induction on  $n$ . The  $n = 1$  case is clear. For  $n = 2$ , choose  $x_1 \in J - I_2 \subseteq I_1$  and  $x_2 \in J - I_1 \subseteq I_2$ . Hence  $x_1 + x_2 \in J - (I_1 \cup I_2)$ , contradiction. Now assume  $n \geq 3$ . Choose  $x_j \in (J - \bigcup_{i \neq j} I_i)$ . Necessarily  $x_j \in I_j$ . Now  $(x_1 \cdots x_{n-1}) + x_n$  belongs to  $J$  but to no  $I_j$ , contradiction.  $\square$

**Application.** Let  $A$  be a commutative Noetherian ring. If  $\mathfrak{p} < \mathfrak{q} < \mathfrak{r}$  are prime, then there exist infinitely many distinct primes  $\mathfrak{q}_i$  such that  $\mathfrak{p} < \mathfrak{q}_i < \mathfrak{r}$ .

*Proof.* Shrink interval so that  $[\mathfrak{p}, \mathfrak{r}]$  has height 2. May assume that  $\mathfrak{p} = (0)$  and that  $\mathfrak{r}$  is maximal. If only finitely many height-1 primes exist,  $\mathfrak{q}_1, \dots, \mathfrak{q}_n$ , then choose  $a \in \mathfrak{r} - (\bigcup_{i=1}^n \mathfrak{q}_i)$ . Any prime minimal over  $(a)$  has height 1 by Krull Height Theorem (needs Noetherianness), and cannot be any of the  $\mathfrak{q}$ 's.



**Prime Avoidance Theorem.** If

- ①  $I_1, \dots, I_n, J$  are ideals,
- ②  $I_j$  is prime for  $j \geq 3$ , and
- ③  $J \subseteq \bigcup_{i=1}^n I_i$ ,

then  $J \subseteq I_j$  for some  $j$ .

*Proof.* Induction on  $n$ . The  $n = 1$  case is clear. For  $n = 2$ , choose  $x_1 \in J - I_2 \subseteq I_1$  and  $x_2 \in J - I_1 \subseteq I_2$ . Hence  $x_1 + x_2 \in J - (I_1 \cup I_2)$ , contradiction. Now assume  $n \geq 3$ . Choose  $x_j \in (J - \bigcup_{i \neq j} I_i)$ . Necessarily  $x_j \in I_j$ . Now  $(x_1 \cdots x_{n-1}) + x_n$  belongs to  $J$  but to no  $I_j$ , contradiction.  $\square$

**Application.** Let  $A$  be a commutative Noetherian ring. If  $\mathfrak{p} < \mathfrak{q} < \mathfrak{r}$  are prime, then there exist infinitely many distinct primes  $\mathfrak{q}_i$  such that  $\mathfrak{p} < \mathfrak{q}_i < \mathfrak{r}$ .

*Proof.* Shrink interval so that  $[\mathfrak{p}, \mathfrak{r}]$  has height 2. May assume that  $\mathfrak{p} = (0)$  and that  $\mathfrak{r}$  is maximal. If only finitely many height-1 primes exist,  $\mathfrak{q}_1, \dots, \mathfrak{q}_n$ , then choose  $a \in \mathfrak{r} - (\bigcup_{i=1}^n \mathfrak{q}_i)$ . Any prime minimal over  $(a)$  has height 1 by Krull Height Theorem (needs Noetherianness), and cannot be any of the  $\mathfrak{q}$ 's.  $\square$