

Jacobson Radical, Nakayama's Lemma



Some radicals

Some radicals

The Wedderburn radical of a ring is the sum of all nilpotent ideals.

Some radicals

The Wedderburn radical of a ring is the sum of all nilpotent ideals.

Joseph Wedderburn proved that if R is a finite dimensional algebra over \mathbb{C} , then $R/W \cong M_{n_1}(\mathbb{C}) \times \cdots \times M_{n_k}(\mathbb{C})$.

Some radicals

The Wedderburn radical of a ring is the sum of all nilpotent ideals.

Joseph Wedderburn proved that if R is a finite dimensional algebra over \mathbb{C} , then $R/W \cong M_{n_1}(\mathbb{C}) \times \cdots \times M_{n_k}(\mathbb{C})$.

The Wedderburn radical was generalized in a satisfactory way to arbitrary rings/algebras by his student Nathan Jacobson.

Some radicals

The Wedderburn radical of a ring is the sum of all nilpotent ideals.

Joseph Wedderburn proved that if R is a finite dimensional algebra over \mathbb{C} , then $R/W \cong M_{n_1}(\mathbb{C}) \times \cdots \times M_{n_k}(\mathbb{C})$.

The Wedderburn radical was generalized in a satisfactory way to arbitrary rings/algebras by his student Nathan Jacobson.

An element $r \in R$ belongs to the *Jacobson radical* of R

Some radicals

The Wedderburn radical of a ring is the sum of all nilpotent ideals.

Joseph Wedderburn proved that if R is a finite dimensional algebra over \mathbb{C} , then $R/W \cong M_{n_1}(\mathbb{C}) \times \cdots \times M_{n_k}(\mathbb{C})$.

The Wedderburn radical was generalized in a satisfactory way to arbitrary rings/algebras by his student Nathan Jacobson.

An element $r \in R$ belongs to the *Jacobson radical* of R if it acts nilpotently on all left R -modules of finite length.

Some radicals

The Wedderburn radical of a ring is the sum of all nilpotent ideals.

Joseph Wedderburn proved that if R is a finite dimensional algebra over \mathbb{C} , then $R/W \cong M_{n_1}(\mathbb{C}) \times \cdots \times M_{n_k}(\mathbb{C})$.

The Wedderburn radical was generalized in a satisfactory way to arbitrary rings/algebras by his student Nathan Jacobson.

An element $r \in R$ belongs to the *Jacobson radical* of R if it acts nilpotently on all left R -modules of finite length. This is equivalent to saying that $rS = \{0\}$ whenever S is a simple R -module.

Equivalent conditions

Equivalent conditions

Theorem

The following are equivalent for $r \in R$:

Equivalent conditions

Theorem

The following are equivalent for $r \in R$:

Equivalent conditions

Theorem

The following are equivalent for $r \in R$:

- ① $rS = \{0\}$ for every simple R -module S .

Equivalent conditions

Theorem

The following are equivalent for $r \in R$:

- ① $rS = \{0\}$ for every simple R -module S .

Equivalent conditions

Theorem

The following are equivalent for $r \in R$:

- ① *$rS = \{0\}$ for every simple R -module S .*
- ② *r lies in every maximal left ideal of R .*

Equivalent conditions

Theorem

The following are equivalent for $r \in R$:

- ① *$rS = \{0\}$ for every simple R -module S .*
- ② *r lies in every maximal left ideal of R .*

Equivalent conditions

Theorem

The following are equivalent for $r \in R$:

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal of R .
- ③ $1 - sr$ is left invertible for any $s \in R$.

Equivalent conditions

Theorem

The following are equivalent for $r \in R$:

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal of R .
- ③ $1 - sr$ is left invertible for any $s \in R$.

Equivalent conditions

Theorem

The following are equivalent for $r \in R$:

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal of R .
- ③ $1 - sr$ is left invertible for any $s \in R$.
- ④ $1 - srt$ is a unit for any $s, t \in R$.

Equivalent conditions

Theorem

The following are equivalent for $r \in R$:

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal of R .
- ③ $1 - sr$ is left invertible for any $s \in R$.
- ④ $1 - srt$ is a unit for any $s, t \in R$.

Equivalent conditions

Theorem

The following are equivalent for $r \in R$:

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal of R .
- ③ $1 - sr$ is left invertible for any $s \in R$.
- ④ $1 - srt$ is a unit for any $s, t \in R$.

Equivalent conditions

Theorem

The following are equivalent for $r \in R$:

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal of R .
- ③ $1 - sr$ is left invertible for any $s \in R$.
- ④ $1 - srt$ is a unit for any $s, t \in R$.

The set of elements having these properties is the *Jacobson radical*, $J(R)$.

Equivalent conditions

Theorem

The following are equivalent for $r \in R$:

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal of R .
- ③ $1 - sr$ is left invertible for any $s \in R$.
- ④ $1 - srt$ is a unit for any $s, t \in R$.

The set of elements having these properties is the *Jacobson radical*, $J(R)$.

Facts/Terminology.

- ① $J(R) = \bigcap \text{max left ideals}$

Equivalent conditions

Theorem

The following are equivalent for $r \in R$:

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal of R .
- ③ $1 - sr$ is left invertible for any $s \in R$.
- ④ $1 - srt$ is a unit for any $s, t \in R$.

The set of elements having these properties is the *Jacobson radical*, $J(R)$.

Facts/Terminology.

- ① $J(R) = \bigcap \text{max left ideals}$

Equivalent conditions

Theorem

The following are equivalent for $r \in R$:

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal of R .
- ③ $1 - sr$ is left invertible for any $s \in R$.
- ④ $1 - srt$ is a unit for any $s, t \in R$.

The set of elements having these properties is the *Jacobson radical*, $J(R)$.

Facts/Terminology.

- ① $J(R) = \bigcap \text{max left ideals} = \text{a 2-sided ideal.}$

Equivalent conditions

Theorem

The following are equivalent for $r \in R$:

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal of R .
- ③ $1 - sr$ is left invertible for any $s \in R$.
- ④ $1 - srt$ is a unit for any $s, t \in R$.

The set of elements having these properties is the *Jacobson radical*, $J(R)$.

Facts/Terminology.

- ① $J(R) = \bigcap \text{max left ideals} = \text{a 2-sided ideal.}$
- ② $\mathfrak{N} \subset J(R)$.

Equivalent conditions

Theorem

The following are equivalent for $r \in R$:

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal of R .
- ③ $1 - sr$ is left invertible for any $s \in R$.
- ④ $1 - srt$ is a unit for any $s, t \in R$.

The set of elements having these properties is the *Jacobson radical*, $J(R)$.

Facts/Terminology.

- ① $J(R) = \bigcap \text{max left ideals} = \text{a 2-sided ideal.}$
- ② $\mathfrak{N} \subset J(R)$.
- ③ Makes sense to talk about the Jacobson radical of an ideal,

Equivalent conditions

Theorem

The following are equivalent for $r \in R$:

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal of R .
- ③ $1 - sr$ is left invertible for any $s \in R$.
- ④ $1 - srt$ is a unit for any $s, t \in R$.

The set of elements having these properties is the *Jacobson radical*, $J(R)$.

Facts/Terminology.

- ① $J(R) = \bigcap \text{max left ideals} = \text{a 2-sided ideal.}$
- ② $\mathfrak{N} \subset J(R)$.
- ③ Makes sense to talk about the Jacobson radical of an ideal,

Equivalent conditions

Theorem

The following are equivalent for $r \in R$:

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal of R .
- ③ $1 - sr$ is left invertible for any $s \in R$.
- ④ $1 - srt$ is a unit for any $s, t \in R$.

The set of elements having these properties is the *Jacobson radical*, $J(R)$.

Facts/Terminology.

- ① $J(R) = \bigcap \text{max left ideals} = \text{a 2-sided ideal.}$
- ② $\mathfrak{N} \subset J(R)$.
- ③ Makes sense to talk about the Jacobson radical of an ideal,
 $J(I) = \bigcap_{I \subseteq M, M \text{ max}} M.$

Some proofs

Some proofs

Theorem. TFAE

Some proofs

Theorem. TFAE

- 1 $rS = \{0\}$ for every simple R -module S .

Some proofs

Theorem. TFAE

- 1 $rS = \{0\}$ for every simple R -module S .

Some proofs

Theorem. TFAE

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal $M \prec R$.

Some proofs

Theorem. TFAE

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal $M \prec R$.

Some proofs

Theorem. TFAE

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal $M \prec R$.
- ③ $1 - sr$ is left invertible for any $s \in R$.

Some proofs

Theorem. TFAE

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal $M \prec R$.
- ③ $1 - sr$ is left invertible for any $s \in R$.

Some proofs

Theorem. TFAE

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal $M \prec R$.
- ③ $1 - sr$ is left invertible for any $s \in R$.

(1) \Rightarrow (2):

Some proofs

Theorem. TFAE

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal $M \prec R$.
- ③ $1 - sr$ is left invertible for any $s \in R$.

$(1) \Rightarrow (2): r(R/M) = \{0\},$

Some proofs

Theorem. TFAE

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal $M \prec R$.
- ③ $1 - sr$ is left invertible for any $s \in R$.

$(1) \Rightarrow (2)$: $r(R/M) = \{0\}$, so $\forall M \prec R (r \in M)$.

Some proofs

Theorem. TFAE

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal $M \prec R$.
- ③ $1 - sr$ is left invertible for any $s \in R$.

$(1) \Rightarrow (2)$: $r(R/M) = \{0\}$, so $\forall M \prec R (r \in M)$.

$((2) \wedge \neg(3)) \Rightarrow \perp$:

Some proofs

Theorem. TFAE

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal $M \prec R$.
- ③ $1 - sr$ is left invertible for any $s \in R$.

$(1) \Rightarrow (2)$: $r(R/M) = \{0\}$, so $\forall M \prec R (r \in M)$.

$((2) \wedge \neg(3)) \Rightarrow \perp$: If $\exists s_0 (1 - s_0 r$ not left invertible),

Some proofs

Theorem. TFAE

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal $M \prec R$.
- ③ $1 - sr$ is left invertible for any $s \in R$.

$(1) \Rightarrow (2)$: $r(R/M) = \{0\}$, so $\forall M \prec R (r \in M)$.

$((2) \wedge \neg(3)) \Rightarrow \perp$: If $\exists s_0(1 - s_0r$ not left invertible), then $R(1 - s_0r) \neq R$,

Some proofs

Theorem. TFAE

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal $M \prec R$.
- ③ $1 - sr$ is left invertible for any $s \in R$.

(1) \Rightarrow (2): $r(R/M) = \{0\}$, so $\forall M \prec R (r \in M)$.

((2) $\wedge \neg$ (3)) $\Rightarrow \perp$: If $\exists s_0(1 - s_0r$ not left invertible), then $R(1 - s_0r) \neq R$, so $\exists M^{\max}(R(1 - s_0r) \leq M)$.

Some proofs

Theorem. TFAE

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal $M \prec R$.
- ③ $1 - sr$ is left invertible for any $s \in R$.

$(1) \Rightarrow (2)$: $r(R/M) = \{0\}$, so $\forall M \prec R (r \in M)$.

$((2) \wedge \neg(3)) \Rightarrow \perp$: If $\exists s_0(1 - s_0r$ not left invertible), then $R(1 - s_0r) \neq R$, so $\exists M^{\max}(R(1 - s_0r) \leq M)$. But $r \in M$,

Some proofs

Theorem. TFAE

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal $M \prec R$.
- ③ $1 - sr$ is left invertible for any $s \in R$.

$(1) \Rightarrow (2)$: $r(R/M) = \{0\}$, so $\forall M \prec R (r \in M)$.

$((2) \wedge \neg(3)) \Rightarrow \perp$: If $\exists s_0(1 - s_0r$ not left invertible), then $R(1 - s_0r) \neq R$, so $\exists M^{\max}(R(1 - s_0r) \leq M)$. But $r \in M$, so $s_0r \in M$,

Some proofs

Theorem. TFAE

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal $M \prec R$.
- ③ $1 - sr$ is left invertible for any $s \in R$.

$(1) \Rightarrow (2)$: $r(R/M) = \{0\}$, so $\forall M \prec R (r \in M)$.

$((2) \wedge \neg(3)) \Rightarrow \perp$: If $\exists s_0(1 - s_0r$ not left invertible), then $R(1 - s_0r) \neq R$, so $\exists M^{\max}(R(1 - s_0r) \leq M)$. But $r \in M$, so $s_0r \in M$, so $(1 - s_0r) + s_0r = 1 \in M$,

Some proofs

Theorem. TFAE

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal $M \prec R$.
- ③ $1 - sr$ is left invertible for any $s \in R$.

$(1) \Rightarrow (2)$: $r(R/M) = \{0\}$, so $\forall M \prec R (r \in M)$.

$((2) \wedge \neg(3)) \Rightarrow \perp$: If $\exists s_0(1 - s_0r$ not left invertible), then $R(1 - s_0r) \neq R$, so $\exists M^{\max}(R(1 - s_0r) \leq M)$. But $r \in M$, so $s_0r \in M$, so $(1 - s_0r) + s_0r = 1 \in M$, \perp

$\neg(1) \Rightarrow \neg(3)$:

Some proofs

Theorem. TFAE

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal $M \prec R$.
- ③ $1 - sr$ is left invertible for any $s \in R$.

$(1) \Rightarrow (2)$: $r(R/M) = \{0\}$, so $\forall M \prec R (r \in M)$.

$((2) \wedge \neg(3)) \Rightarrow \perp$: If $\exists s_0(1 - s_0r$ not left invertible), then $R(1 - s_0r) \neq R$, so $\exists M^{\max}(R(1 - s_0r) \leq M)$. But $r \in M$, so $s_0r \in M$, so $(1 - s_0r) + s_0r = 1 \in M$, \perp

$\neg(1) \Rightarrow \neg(3)$: If $rS \neq \{0\}$,

Some proofs

Theorem. TFAE

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal $M \prec R$.
- ③ $1 - sr$ is left invertible for any $s \in R$.

$(1) \Rightarrow (2)$: $r(R/M) = \{0\}$, so $\forall M \prec R (r \in M)$.

$((2) \wedge \neg(3)) \Rightarrow \perp$: If $\exists s_0(1 - s_0r$ not left invertible), then $R(1 - s_0r) \neq R$, so $\exists M^{\max}(R(1 - s_0r) \leq M)$. But $r \in M$, so $s_0r \in M$, so $(1 - s_0r) + s_0r = 1 \in M$, \perp

$\neg(1) \Rightarrow \neg(3)$: If $rS \neq \{0\}$, $\exists x \in S(rx \neq 0)$,

Some proofs

Theorem. TFAE

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal $M \prec R$.
- ③ $1 - sr$ is left invertible for any $s \in R$.

$(1) \Rightarrow (2)$: $r(R/M) = \{0\}$, so $\forall M \prec R (r \in M)$.

$((2) \wedge \neg(3)) \Rightarrow \perp$: If $\exists s_0(1 - s_0r$ not left invertible), then $R(1 - s_0r) \neq R$, so $\exists M^{\max}(R(1 - s_0r) \leq M)$. But $r \in M$, so $s_0r \in M$, so $(1 - s_0r) + s_0r = 1 \in M$, \perp

$\neg(1) \Rightarrow \neg(3)$: If $rS \neq \{0\}$, $\exists x \in S(rx \neq 0)$, then since $RrS = S$

Some proofs

Theorem. TFAE

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal $M \prec R$.
- ③ $1 - sr$ is left invertible for any $s \in R$.

$(1) \Rightarrow (2)$: $r(R/M) = \{0\}$, so $\forall M \prec R (r \in M)$.

$((2) \wedge \neg(3)) \Rightarrow \perp$: If $\exists s_0(1 - s_0r$ not left invertible), then $R(1 - s_0r) \neq R$, so $\exists M^{\max}(R(1 - s_0r) \leq M)$. But $r \in M$, so $s_0r \in M$, so $(1 - s_0r) + s_0r = 1 \in M$, \perp

$\neg(1) \Rightarrow \neg(3)$: If $rS \neq \{0\}$, $\exists x \in S(rx \neq 0)$, then since $RrS = S \exists s \in R(srx = x)$,

Some proofs

Theorem. TFAE

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal $M \prec R$.
- ③ $1 - sr$ is left invertible for any $s \in R$.

$(1) \Rightarrow (2)$: $r(R/M) = \{0\}$, so $\forall M \prec R (r \in M)$.

$((2) \wedge \neg(3)) \Rightarrow \perp$: If $\exists s_0(1 - s_0r$ not left invertible), then $R(1 - s_0r) \neq R$, so $\exists M^{\max}(R(1 - s_0r) \leq M)$. But $r \in M$, so $s_0r \in M$, so $(1 - s_0r) + s_0r = 1 \in M$, \perp

$\neg(1) \Rightarrow \neg(3)$: If $rS \neq \{0\}$, $\exists x \in S(rx \neq 0)$, then since $RrS = S \exists s \in R(srx = x)$, implying $(1 - sr)x = 0$,

Some proofs

Theorem. TFAE

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal $M \prec R$.
- ③ $1 - sr$ is left invertible for any $s \in R$.

$(1) \Rightarrow (2)$: $r(R/M) = \{0\}$, so $\forall M \prec R (r \in M)$.

$((2) \wedge \neg(3)) \Rightarrow \perp$: If $\exists s_0(1 - s_0r$ not left invertible), then $R(1 - s_0r) \neq R$, so $\exists M^{\max}(R(1 - s_0r) \leq M)$. But $r \in M$, so $s_0r \in M$, so $(1 - s_0r) + s_0r = 1 \in M$, \perp

$\neg(1) \Rightarrow \neg(3)$: If $rS \neq \{0\}$, $\exists x \in S(rx \neq 0)$, then since $RrS = S \exists s \in R(srx = x)$, implying $(1 - sr)x = 0$, so $(1 - sr)$ cannot be left invertible.

Some proofs

Theorem. TFAE

- ① $rS = \{0\}$ for every simple R -module S .
- ② r lies in every maximal left ideal $M \prec R$.
- ③ $1 - sr$ is left invertible for any $s \in R$.

$(1) \Rightarrow (2)$: $r(R/M) = \{0\}$, so $\forall M \prec R (r \in M)$.

$((2) \wedge \neg(3)) \Rightarrow \perp$: If $\exists s_0(1 - s_0r$ not left invertible), then $R(1 - s_0r) \neq R$, so $\exists M^{\max}(R(1 - s_0r) \leq M)$. But $r \in M$, so $s_0r \in M$, so $(1 - s_0r) + s_0r = 1 \in M$, \perp

$\neg(1) \Rightarrow \neg(3)$: If $rS \neq \{0\}$, $\exists x \in S(rx \neq 0)$, then since $RrS = S \exists s \in R(srx = x)$, implying $(1 - sr)x = 0$, so $(1 - sr)$ cannot be left invertible. \square

Jacobson radical of a module

Jacobson radical of a module

Df. If M is an R -module, then $J(M) := \bigcap_{N \prec M} N$.

Jacobson radical of a module

Df. If M is an R -module, then $J(M) := \bigcap_{N \prec M} N$.

Hence the Jacobson radical of R is the same whether R is considered as a ring or as the R -module ${}_R R$.

Jacobson radical of a module

Df. If M is an R -module, then $J(M) := \bigcap_{N \prec M} N$.

Hence the Jacobson radical of R is the same whether R is considered as a ring or as the R -module ${}_R R$.

Examples. ($R = \mathbb{Z}$)

Jacobson radical of a module

Df. If M is an R -module, then $J(M) := \bigcap_{N \prec M} N$.

Hence the Jacobson radical of R is the same whether R is considered as a ring or as the R -module ${}_R R$.

Examples. ($R = \mathbb{Z}$)

$$J(\mathbb{Z}) = \{0\};$$

Jacobson radical of a module

Df. If M is an R -module, then $J(M) := \bigcap_{N \prec M} N$.

Hence the Jacobson radical of R is the same whether R is considered as a ring or as the R -module ${}_R R$.

Examples. ($R = \mathbb{Z}$)

$$J(\mathbb{Z}) = \{0\}; \quad J(\mathbb{Z}_4) = 2\mathbb{Z}_4;$$

Jacobson radical of a module

Df. If M is an R -module, then $J(M) := \bigcap_{N \prec M} N$.

Hence the Jacobson radical of R is the same whether R is considered as a ring or as the R -module ${}_R R$.

Examples. ($R = \mathbb{Z}$)

$$J(\mathbb{Z}) = \{0\}; \quad J(\mathbb{Z}_4) = 2\mathbb{Z}_4; \quad J(\mathbb{Q}) = \mathbb{Q}.$$

Jacobson radical of a module

Df. If M is an R -module, then $J(M) := \bigcap_{N \prec M} N$.

Hence the Jacobson radical of R is the same whether R is considered as a ring or as the R -module ${}_R R$.

Examples. ($R = \mathbb{Z}$)

$$J(\mathbb{Z}) = \{0\}; \quad J(\mathbb{Z}_4) = 2\mathbb{Z}_4; \quad J(\mathbb{Q}) = \mathbb{Q}.$$

Lemma. $JM \subseteq J(M)$.

Jacobson radical of a module

Df. If M is an R -module, then $J(M) := \bigcap_{N \prec M} N$.

Hence the Jacobson radical of R is the same whether R is considered as a ring or as the R -module ${}_R R$.

Examples. ($R = \mathbb{Z}$)

$$J(\mathbb{Z}) = \{0\}; \quad J(\mathbb{Z}_4) = 2\mathbb{Z}_4; \quad J(\mathbb{Q}) = \mathbb{Q}.$$

Lemma. $JM \subseteq J(M)$. If M is finitely generated and $\neq \{0\}$, then $J(M) \subsetneq M$.

Jacobson radical of a module

Df. If M is an R -module, then $J(M) := \bigcap_{N \prec M} N$.

Hence the Jacobson radical of R is the same whether R is considered as a ring or as the R -module ${}_R R$.

Examples. ($R = \mathbb{Z}$)

$$J(\mathbb{Z}) = \{0\}; \quad J(\mathbb{Z}_4) = 2\mathbb{Z}_4; \quad J(\mathbb{Q}) = \mathbb{Q}.$$

Lemma. $JM \subseteq J(M)$. If M is finitely generated and $\neq \{0\}$, then $J(M) \subsetneq M$.

Proof.

(1st claim)

Jacobson radical of a module

Df. If M is an R -module, then $J(M) := \bigcap_{N \prec M} N$.

Hence the Jacobson radical of R is the same whether R is considered as a ring or as the R -module ${}_R R$.

Examples. ($R = \mathbb{Z}$)

$$J(\mathbb{Z}) = \{0\}; \quad J(\mathbb{Z}_4) = 2\mathbb{Z}_4; \quad J(\mathbb{Q}) = \mathbb{Q}.$$

Lemma. $JM \subseteq J(M)$. If M is finitely generated and $\neq \{0\}$, then $J(M) \subsetneq M$.

Proof.

(1st claim) If $N \prec M$,

Jacobson radical of a module

Df. If M is an R -module, then $J(M) := \bigcap_{N \prec M} N$.

Hence the Jacobson radical of R is the same whether R is considered as a ring or as the R -module ${}_R R$.

Examples. ($R = \mathbb{Z}$)

$$J(\mathbb{Z}) = \{0\}; \quad J(\mathbb{Z}_4) = 2\mathbb{Z}_4; \quad J(\mathbb{Q}) = \mathbb{Q}.$$

Lemma. $JM \subseteq J(M)$. If M is finitely generated and $\neq \{0\}$, then $J(M) \subsetneq M$.

Proof.

(1st claim) If $N \prec M$, then M/N is simple,

Jacobson radical of a module

Df. If M is an R -module, then $J(M) := \bigcap_{N \prec M} N$.

Hence the Jacobson radical of R is the same whether R is considered as a ring or as the R -module ${}_R R$.

Examples. ($R = \mathbb{Z}$)

$$J(\mathbb{Z}) = \{0\}; \quad J(\mathbb{Z}_4) = 2\mathbb{Z}_4; \quad J(\mathbb{Q}) = \mathbb{Q}.$$

Lemma. $JM \subseteq J(M)$. If M is finitely generated and $\neq \{0\}$, then $J(M) \subsetneq M$.

Proof.

(1st claim) If $N \prec M$, then M/N is simple, so $J \cdot M/N = 0 = J(M/N)$,

Jacobson radical of a module

Df. If M is an R -module, then $J(M) := \bigcap_{N \prec M} N$.

Hence the Jacobson radical of R is the same whether R is considered as a ring or as the R -module ${}_R R$.

Examples. ($R = \mathbb{Z}$)

$$J(\mathbb{Z}) = \{0\}; \quad J(\mathbb{Z}_4) = 2\mathbb{Z}_4; \quad J(\mathbb{Q}) = \mathbb{Q}.$$

Lemma. $JM \subseteq J(M)$. If M is finitely generated and $\neq \{0\}$, then $J(M) \subsetneq M$.

Proof.

(1st claim) If $N \prec M$, then M/N is simple, so $J \cdot M/N = 0 = J(M/N)$, so $JM \subseteq N$.

Jacobson radical of a module

Df. If M is an R -module, then $J(M) := \bigcap_{N \prec M} N$.

Hence the Jacobson radical of R is the same whether R is considered as a ring or as the R -module ${}_R R$.

Examples. ($R = \mathbb{Z}$)

$$J(\mathbb{Z}) = \{0\}; \quad J(\mathbb{Z}_4) = 2\mathbb{Z}_4; \quad J(\mathbb{Q}) = \mathbb{Q}.$$

Lemma. $JM \subseteq J(M)$. If M is finitely generated and $\neq \{0\}$, then $J(M) \subsetneq M$.

Proof.

(1st claim) If $N \prec M$, then M/N is simple, so $J \cdot M/N = 0 = J(M/N)$, so $JM \subseteq N$. N was arbitrary, so $JM \subseteq \bigcap_{N \prec M} N = J(M)$.

Jacobson radical of a module

Df. If M is an R -module, then $J(M) := \bigcap_{N \prec M} N$.

Hence the Jacobson radical of R is the same whether R is considered as a ring or as the R -module ${}_R R$.

Examples. ($R = \mathbb{Z}$)

$$J(\mathbb{Z}) = \{0\}; \quad J(\mathbb{Z}_4) = 2\mathbb{Z}_4; \quad J(\mathbb{Q}) = \mathbb{Q}.$$

Lemma. $JM \subseteq J(M)$. If M is finitely generated and $\neq \{0\}$, then $J(M) \subsetneq M$.

Proof.

(1st claim) If $N \prec M$, then M/N is simple, so $J \cdot M/N = 0 = J(M/N)$, so $JM \subseteq N$. N was arbitrary, so $JM \subseteq \bigcap_{N \prec M} N = J(M)$. \square

Jacobson radical of a module

Df. If M is an R -module, then $J(M) := \bigcap_{N \prec M} N$.

Hence the Jacobson radical of R is the same whether R is considered as a ring or as the R -module ${}_R R$.

Examples. ($R = \mathbb{Z}$)

$$J(\mathbb{Z}) = \{0\}; \quad J(\mathbb{Z}_4) = 2\mathbb{Z}_4; \quad J(\mathbb{Q}) = \mathbb{Q}.$$

Lemma. $JM \subseteq J(M)$. If M is finitely generated and $\neq \{0\}$, then $J(M) \subsetneq M$.

Proof.

(1st claim) If $N \prec M$, then M/N is simple, so $J \cdot M/N = 0 = J(M/N)$, so $JM \subseteq N$. N was arbitrary, so $JM \subseteq \bigcap_{N \prec M} N = J(M)$. \square

(2nd claim)

Jacobson radical of a module

Df. If M is an R -module, then $J(M) := \bigcap_{N \prec M} N$.

Hence the Jacobson radical of R is the same whether R is considered as a ring or as the R -module ${}_R R$.

Examples. ($R = \mathbb{Z}$)

$$J(\mathbb{Z}) = \{0\}; \quad J(\mathbb{Z}_4) = 2\mathbb{Z}_4; \quad J(\mathbb{Q}) = \mathbb{Q}.$$

Lemma. $JM \subseteq J(M)$. If M is finitely generated and $\neq \{0\}$, then $J(M) \subsetneq M$.

Proof.

(1st claim) If $N \prec M$, then M/N is simple, so $J \cdot M/N = 0 = J(M/N)$, so $JM \subseteq N$. N was arbitrary, so $JM \subseteq \bigcap_{N \prec M} N = J(M)$. \square

(2nd claim) If $\{m_1, \dots, m_n\}$ is a minimal generating set for M

Jacobson radical of a module

Df. If M is an R -module, then $J(M) := \bigcap_{N \prec M} N$.

Hence the Jacobson radical of R is the same whether R is considered as a ring or as the R -module ${}_R R$.

Examples. ($R = \mathbb{Z}$)

$$J(\mathbb{Z}) = \{0\}; \quad J(\mathbb{Z}_4) = 2\mathbb{Z}_4; \quad J(\mathbb{Q}) = \mathbb{Q}.$$

Lemma. $JM \subseteq J(M)$. If M is finitely generated and $\neq \{0\}$, then $J(M) \subsetneq M$.

Proof.

(1st claim) If $N \prec M$, then M/N is simple, so $J \cdot M/N = 0 = J(M/N)$, so $JM \subseteq N$. N was arbitrary, so $JM \subseteq \bigcap_{N \prec M} N = J(M)$. \square

(2nd claim) If $\{m_1, \dots, m_n\}$ is a minimal generating set for M and $N = \langle m_1, \dots, m_{n-1} \rangle$,

Jacobson radical of a module

Df. If M is an R -module, then $J(M) := \bigcap_{N \prec M} N$.

Hence the Jacobson radical of R is the same whether R is considered as a ring or as the R -module ${}_R R$.

Examples. ($R = \mathbb{Z}$)

$$J(\mathbb{Z}) = \{0\}; \quad J(\mathbb{Z}_4) = 2\mathbb{Z}_4; \quad J(\mathbb{Q}) = \mathbb{Q}.$$

Lemma. $JM \subseteq J(M)$. If M is finitely generated and $\neq \{0\}$, then $J(M) \subsetneq M$.

Proof.

(1st claim) If $N \prec M$, then M/N is simple, so $J \cdot M/N = 0 = J(M/N)$, so $JM \subseteq N$. N was arbitrary, so $JM \subseteq \bigcap_{N \prec M} N = J(M)$. \square

(2nd claim) If $\{m_1, \dots, m_n\}$ is a minimal generating set for M and $N = \langle m_1, \dots, m_{n-1} \rangle$, then there is a submodule N' satisfying $N \leq N' \prec M$.

Jacobson radical of a module

Df. If M is an R -module, then $J(M) := \bigcap_{N \prec M} N$.

Hence the Jacobson radical of R is the same whether R is considered as a ring or as the R -module ${}_R R$.

Examples. ($R = \mathbb{Z}$)

$$J(\mathbb{Z}) = \{0\}; \quad J(\mathbb{Z}_4) = 2\mathbb{Z}_4; \quad J(\mathbb{Q}) = \mathbb{Q}.$$

Lemma. $JM \subseteq J(M)$. If M is finitely generated and $\neq \{0\}$, then $J(M) \subsetneq M$.

Proof.

(1st claim) If $N \prec M$, then M/N is simple, so $J \cdot M/N = 0 = J(M/N)$, so $JM \subseteq N$. N was arbitrary, so $JM \subseteq \bigcap_{N \prec M} N = J(M)$. \square

(2nd claim) If $\{m_1, \dots, m_n\}$ is a minimal generating set for M and $N = \langle m_1, \dots, m_{n-1} \rangle$, then there is a submodule N' satisfying $N \leq N' \prec M$. Hence $J(M) \leq N' \subsetneq M$.

Jacobson radical of a module

Df. If M is an R -module, then $J(M) := \bigcap_{N \prec M} N$.

Hence the Jacobson radical of R is the same whether R is considered as a ring or as the R -module ${}_R R$.

Examples. ($R = \mathbb{Z}$)

$$J(\mathbb{Z}) = \{0\}; \quad J(\mathbb{Z}_4) = 2\mathbb{Z}_4; \quad J(\mathbb{Q}) = \mathbb{Q}.$$

Lemma. $JM \subseteq J(M)$. If M is finitely generated and $\neq \{0\}$, then $J(M) \subsetneq M$.

Proof.

(1st claim) If $N \prec M$, then M/N is simple, so $J \cdot M/N = 0 = J(M/N)$, so $JM \subseteq N$. N was arbitrary, so $JM \subseteq \bigcap_{N \prec M} N = J(M)$. \square

(2nd claim) If $\{m_1, \dots, m_n\}$ is a minimal generating set for M and $N = \langle m_1, \dots, m_{n-1} \rangle$, then there is a submodule N' satisfying $N \leq N' \prec M$. Hence $J(M) \leq N' \subsetneq M$. \square

Nakayama's Lemma = Krull-Azumaya Theorem

Nakayama's Lemma = Krull-Azumaya Theorem

Nakayama's Lemma. If M is a finitely generated R -module and $JM = M$, then $M = \{0\}$.

Nakayama's Lemma = Krull-Azumaya Theorem

Nakayama's Lemma. If M is a finitely generated R -module and $JM = M$, then $M = \{0\}$.

Nontraditional Proof.

Nakayama's Lemma = Krull-Azumaya Theorem

Nakayama's Lemma. If M is a finitely generated R -module and $JM = M$, then $M = \{0\}$.

Nontraditional Proof. $M \neq \{0\}$, then $JM \leq J(M) \subsetneq M$. \square

Nakayama's Lemma = Krull-Azumaya Theorem

Nakayama's Lemma. If M is a finitely generated R -module and $JM = M$, then $M = \{0\}$.

Nontraditional Proof. $M \neq \{0\}$, then $JM \leq J(M) \subsetneq M$. \square

The traditional proof of Nakayama's Lemma is based on the Cayley-Hamilton Theorem for commutative rings.

Nakayama's Lemma = Krull-Azumaya Theorem

Nakayama's Lemma. If M is a finitely generated R -module and $JM = M$, then $M = \{0\}$.

Nontraditional Proof. $M \neq \{0\}$, then $JM \leq J(M) \subsetneq M$. \square

The traditional proof of Nakayama's Lemma is based on the Cayley-Hamilton Theorem for commutative rings.

Cayley-Hamilton Theorem

Cayley-Hamilton Theorem

Cayley-Hamilton Theorem for fields. Any $M \in M_n(\mathbb{F})$

Cayley-Hamilton Theorem

Cayley-Hamilton Theorem for fields. Any $M \in M_n(\mathbb{F})$ satisfies its own characteristic polynomial.

Cayley-Hamilton Theorem

Cayley-Hamilton Theorem for fields. Any $M \in M_n(\mathbb{F})$ satisfies its own characteristic polynomial.

Example.

Cayley-Hamilton Theorem

Cayley-Hamilton Theorem for fields. Any $M \in M_n(\mathbb{F})$ satisfies its own characteristic polynomial.

Example.

$$\text{If } M = \begin{bmatrix} a & b \\ c & d \end{bmatrix},$$

Cayley-Hamilton Theorem

Cayley-Hamilton Theorem for fields. Any $M \in M_n(\mathbb{F})$ satisfies its own characteristic polynomial.

Example.

If $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, then $\chi_M(t) = t^2 - \operatorname{tr}(M)t + \det(M)$.

Cayley-Hamilton Theorem

Cayley-Hamilton Theorem for fields. Any $M \in M_n(\mathbb{F})$ satisfies its own characteristic polynomial.

Example.

If $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, then $\chi_M(t) = t^2 - \operatorname{tr}(M)t + \det(M)$. C-H asserts that $M^2 - \operatorname{tr}(M)M + \det(M)I = 0$.

Cayley-Hamilton Theorem

Cayley-Hamilton Theorem for fields. Any $M \in M_n(\mathbb{F})$ satisfies its own characteristic polynomial.

Example.

If $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, then $\chi_M(t) = t^2 - \operatorname{tr}(M)t + \det(M)$. C-H asserts that $M^2 - \operatorname{tr}(M)M + \det(M)I = 0$. More explicitly,

$$\begin{bmatrix} a^2 + bc & ab + bd \\ ac + cd & bc + d^2 \end{bmatrix} - (a + d) \begin{bmatrix} a & b \\ c & d \end{bmatrix} + (ad - bc) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}.$$

Cayley-Hamilton Theorem

Cayley-Hamilton Theorem for fields. Any $M \in M_n(\mathbb{F})$ satisfies its own characteristic polynomial.

Example.

If $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, then $\chi_M(t) = t^2 - \operatorname{tr}(M)t + \det(M)$. C-H asserts that $M^2 - \operatorname{tr}(M)M + \det(M)I = 0$. More explicitly,

$$\begin{bmatrix} a^2 + bc & ab + bd \\ ac + cd & bc + d^2 \end{bmatrix} - (a + d) \begin{bmatrix} a & b \\ c & d \end{bmatrix} + (ad - bc) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}.$$

C-H for 2×2 matrices is equivalent to

Cayley-Hamilton Theorem

Cayley-Hamilton Theorem for fields. Any $M \in M_n(\mathbb{F})$ satisfies its own characteristic polynomial.

Example.

If $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, then $\chi_M(t) = t^2 - \text{tr}(M)t + \det(M)$. C-H asserts that $M^2 - \text{tr}(M)M + \det(M)I = 0$. More explicitly,

$$\begin{bmatrix} a^2 + bc & ab + bd \\ ac + cd & bc + d^2 \end{bmatrix} - (a + d) \begin{bmatrix} a & b \\ c & d \end{bmatrix} + (ad - bc) \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}.$$

C-H for 2×2 matrices is equivalent to

$$\begin{aligned} &(\forall a)(\forall b)(\forall c)(\forall d) \\ &\quad ((a^2 + bc) - (a + d)a + (ad - bc)1 = 0) \\ &\quad \wedge ((ab + bd) - (a + d)b + (ad - bc)0 = 0) \\ &\quad \wedge ((ac + cd) - (a + d)c + (ad - bc)0 = 0) \\ &\quad \wedge ((bc + d^2) - (a + d)d + (ad - bc)1 = 0) \end{aligned}$$

Transfer Theorem (calg1p8: Toby, Ezzeddine, Howie)

Transfer Theorem (calg1p8: Toby, Ezzeddine, Howie)

Thm. Any positive

Transfer Theorem (calg1p8: Toby, Ezzeddine, Howie)

Thm. Any positive universal

Thm. Any positive universal first-order

Thm. Any positive universal first-order sentence

Transfer Theorem (calg1p8: Toby, Ezzeddine, Howie)

Thm. Any positive universal first-order sentence true in all fields

Transfer Theorem (calg1p8: Toby, Ezzeddine, Howie)

Thm. Any positive universal first-order sentence true in all fields is true in all commutative rings.

Transfer Theorem (calg1p8: Toby, Ezzeddine, Howie)

Thm. Any positive universal first-order sentence true in all fields is true in all commutative rings.

Roughly speaking:

Thm. Any positive universal first-order sentence true in all fields is true in all commutative rings.

Roughly speaking:

① first-order =

Thm. Any positive universal first-order sentence true in all fields is true in all commutative rings.

Roughly speaking:

① first-order =

Transfer Theorem (calg1p8: Toby, Ezzeddine, Howie)

Thm. Any positive universal first-order sentence true in all fields is true in all commutative rings.

Roughly speaking:

- ① first-order = well-formed, finite length, quantification over elements of the first order only.

Transfer Theorem (calg1p8: Toby, Ezzeddine, Howie)

Thm. Any positive universal first-order sentence true in all fields is true in all commutative rings.

Roughly speaking:

- ① first-order = well-formed, finite length, quantification over elements of the first order only.
(Can be written: $(\text{quantifiers})[\bigvee(\bigwedge \pm \text{atomic})]$.)
- ② universal =

Transfer Theorem (calg1p8: Toby, Ezzeddine, Howie)

Thm. Any positive universal first-order sentence true in all fields is true in all commutative rings.

Roughly speaking:

- ① first-order = well-formed, finite length, quantification over elements of the first order only.
(Can be written: $(\text{quantifiers})[\bigvee(\bigwedge \pm \text{atomic})]$.)
- ② universal =

Thm. Any positive universal first-order sentence true in all fields is true in all commutative rings.

Roughly speaking:

- ① first-order = well-formed, finite length, quantification over elements of the first order only.
(Can be written: (quantifiers)[$\forall(\wedge \pm \text{atomic})$].)
- ② universal = all quantifiers are \forall .
- ③ positive =

Thm. Any positive universal first-order sentence true in all fields is true in all commutative rings.

Roughly speaking:

- ① first-order = well-formed, finite length, quantification over elements of the first order only.
(Can be written: (quantifiers)[$\forall(\wedge \pm \text{atomic})$].)
- ② universal = all quantifiers are \forall .
- ③ positive =

Thm. Any positive universal first-order sentence true in all fields is true in all commutative rings.

Roughly speaking:

- ① first-order = well-formed, finite length, quantification over elements of the first order only.
(Can be written: (quantifiers)[$\forall(\wedge \pm\text{atomic})$].)
- ② universal = all quantifiers are \forall .
- ③ positive = no occurrence of \neg , when written in prenex form.
($\pm\text{atomic}$ is always $+\text{atomic}$.)
- ④ sentence =

Thm. Any positive universal first-order sentence true in all fields is true in all commutative rings.

Roughly speaking:

- ① first-order = well-formed, finite length, quantification over elements of the first order only.
(Can be written: (quantifiers)[$\forall(\wedge \pm\text{atomic})$].)
- ② universal = all quantifiers are \forall .
- ③ positive = no occurrence of \neg , when written in prenex form.
($\pm\text{atomic}$ is always $+\text{atomic}$.)
- ④ sentence =

Thm. Any positive universal first-order sentence true in all fields is true in all commutative rings.

Roughly speaking:

- ① first-order = well-formed, finite length, quantification over elements of the first order only.
(Can be written: (quantifiers)[$\forall(\wedge \pm\text{atomic})$].)
- ② universal = all quantifiers are \forall .
- ③ positive = no occurrence of \neg , when written in prenex form.
($\pm\text{atomic}$ is always $+\text{atomic}$.)
- ④ sentence = all variables bound by quantifiers.

Traditional proof of Nakayama's Lemma

Traditional proof of Nakayama's Lemma

Assume that $M = \langle m_1, \dots, m_m \rangle$ and $M = IM$ for some ideal I .

Traditional proof of Nakayama's Lemma

Assume that $M = \langle m_1, \dots, m_m \rangle$ and $M = IM$ for some ideal I .

The identity function on M is representable as an $n \times n$ matrix A with coefficients in I .

Traditional proof of Nakayama's Lemma

Assume that $M = \langle m_1, \dots, m_m \rangle$ and $M = IM$ for some ideal I .

The identity function on M is representable as an $n \times n$ matrix A with coefficients in I .

If $m_1 = i_1 m_1 + \dots + i_n m_n$, then $A = \begin{bmatrix} i_1 & \dots & \dots \\ i_2 & \dots & \dots \\ \vdots & & \\ i_n & \dots & \dots \end{bmatrix}.$

Traditional proof of Nakayama's Lemma

Assume that $M = \langle m_1, \dots, m_m \rangle$ and $M = IM$ for some ideal I .

The identity function on M is representable as an $n \times n$ matrix A with coefficients in I .

If $m_1 = i_1 m_1 + \dots + i_n m_n$, then $A = \begin{bmatrix} i_1 & \dots & \dots \\ i_2 & \dots & \dots \\ \vdots & & \\ i_n & \dots & \dots \end{bmatrix}.$

Assume $\chi_A(t) = t^n - a_1 t^{n-1} + \dots + (-1)^n a_0$,

Traditional proof of Nakayama's Lemma

Assume that $M = \langle m_1, \dots, m_m \rangle$ and $M = IM$ for some ideal I .

The identity function on M is representable as an $n \times n$ matrix A with coefficients in I .

If $m_1 = i_1 m_1 + \dots + i_n m_n$, then $A = \begin{bmatrix} i_1 & \dots & \dots \\ i_2 & \dots & \dots \\ \vdots & & \\ i_n & \dots & \dots \end{bmatrix}$.

Assume $\chi_A(t) = t^n - a_1 t^{n-1} + \dots + (-1)^n a_0$, $a_i \in I$.

Traditional proof of Nakayama's Lemma

Assume that $M = \langle m_1, \dots, m_m \rangle$ and $M = IM$ for some ideal I .

The identity function on M is representable as an $n \times n$ matrix A with coefficients in I .

If $m_1 = i_1 m_1 + \dots + i_n m_n$, then $A = \begin{bmatrix} i_1 & \dots & \dots \\ i_2 & \dots & \dots \\ \vdots & & \\ i_n & \dots & \dots \end{bmatrix}$.

Assume $\chi_A(t) = t^n - a_1 t^{n-1} + \dots + (-1)^n a_0$, $a_i \in I$. (C-H)
 $0 = \chi_A(\text{id})$

Traditional proof of Nakayama's Lemma

Assume that $M = \langle m_1, \dots, m_m \rangle$ and $M = IM$ for some ideal I .

The identity function on M is representable as an $n \times n$ matrix A with coefficients in I .

If $m_1 = i_1 m_1 + \dots + i_n m_n$, then $A = \begin{bmatrix} i_1 & \dots & \dots \\ i_2 & \dots & \dots \\ \vdots & & \\ i_n & \dots & \dots \end{bmatrix}$.

Assume $\chi_A(t) = t^n - a_1 t^{n-1} + \dots + (-1)^n a_0$, $a_i \in I$. (C-H)
 $0 = \chi_A(\text{id}) = \chi_A(\lambda_1)$

Traditional proof of Nakayama's Lemma

Assume that $M = \langle m_1, \dots, m_m \rangle$ and $M = IM$ for some ideal I .

The identity function on M is representable as an $n \times n$ matrix A with coefficients in I .

If $m_1 = i_1 m_1 + \dots + i_n m_n$, then $A = \begin{bmatrix} i_1 & \dots & \dots \\ i_2 & \dots & \dots \\ \vdots & & \\ i_n & \dots & \dots \end{bmatrix}$.

Assume $\chi_A(t) = t^n - a_1 t^{n-1} + \dots + (-1)^n a_0$, $a_i \in I$. (C-H)
 $0 = \chi_A(\text{id}) = \chi_A(\lambda_1) = \lambda_1^n - a_1 \lambda_1^{n-1} + \dots + (-1)^n a_0$,

Traditional proof of Nakayama's Lemma

Assume that $M = \langle m_1, \dots, m_m \rangle$ and $M = IM$ for some ideal I .

The identity function on M is representable as an $n \times n$ matrix A with coefficients in I .

If $m_1 = i_1 m_1 + \dots + i_n m_n$, then $A = \begin{bmatrix} i_1 & \dots & \dots \\ i_2 & \dots & \dots \\ \vdots & & \\ i_n & \dots & \dots \end{bmatrix}$.

Assume $\chi_A(t) = t^n - a_1 t^{n-1} + \dots + (-1)^n a_0$, $a_i \in I$. (C-H)

$0 = \chi_A(\text{id}) = \chi_A(\lambda_1) = \lambda_1^n - a_1 \lambda_1^{n-1} + \dots + (-1)^n a_0 = \lambda_{\chi_A(1)}$.

Traditional proof of Nakayama's Lemma

Assume that $M = \langle m_1, \dots, m_m \rangle$ and $M = IM$ for some ideal I .

The identity function on M is representable as an $n \times n$ matrix A with coefficients in I .

If $m_1 = i_1 m_1 + \dots + i_n m_n$, then $A = \begin{bmatrix} i_1 & \dots & \dots \\ i_2 & \dots & \dots \\ \vdots & & \\ i_n & \dots & \dots \end{bmatrix}$.

Assume $\chi_A(t) = t^n - a_1 t^{n-1} + \dots + (-1)^n a_0$, $a_i \in I$. (C-H)

$0 = \chi_A(\text{id}) = \chi_A(\lambda_1) = \lambda_1^n - a_1 \lambda_1^{n-1} + \dots + (-1)^n a_0 = \lambda_{\chi_A(1)}$.

Traditional proof of Nakayama's Lemma

Assume that $M = \langle m_1, \dots, m_m \rangle$ and $M = IM$ for some ideal I .

The identity function on M is representable as an $n \times n$ matrix A with coefficients in I .

If $m_1 = i_1 m_1 + \dots + i_n m_n$, then $A = \begin{bmatrix} i_1 & \dots & \dots \\ i_2 & \dots & \dots \\ \vdots & & \\ i_n & \dots & \dots \end{bmatrix}$.

Assume $\chi_A(t) = t^n - a_1 t^{n-1} + \dots + (-1)^n a_0$, $a_i \in I$. (C-H)

$0 = \chi_A(\text{id}) = \chi_A(\lambda_1) = \lambda_1^n - a_1 \lambda_1^{n-1} + \dots + (-1)^n a_0 = \lambda_{\chi_A(1)}$.

Now $\chi_A(1) = 1 - i$ for some $i \in I$,

Traditional proof of Nakayama's Lemma

Assume that $M = \langle m_1, \dots, m_m \rangle$ and $M = IM$ for some ideal I .

The identity function on M is representable as an $n \times n$ matrix A with coefficients in I .

If $m_1 = i_1 m_1 + \dots + i_n m_n$, then $A = \begin{bmatrix} i_1 & \dots & \dots \\ i_2 & \dots & \dots \\ \vdots & & \\ i_n & \dots & \dots \end{bmatrix}$.

Assume $\chi_A(t) = t^n - a_1 t^{n-1} + \dots + (-1)^n a_0$, $a_i \in I$. (C-H)

$0 = \chi_A(\text{id}) = \chi_A(\lambda_1) = \lambda_1^n - a_1 \lambda_1^{n-1} + \dots + (-1)^n a_0 = \lambda_{\chi_A(1)}$.

Now $\chi_A(1) = 1 - i$ for some $i \in I$, so $(1 - i)M = 0$ for some $i \in I$.

Traditional proof of Nakayama's Lemma

Assume that $M = \langle m_1, \dots, m_m \rangle$ and $M = IM$ for some ideal I .

The identity function on M is representable as an $n \times n$ matrix A with coefficients in I .

If $m_1 = i_1 m_1 + \dots + i_n m_n$, then $A = \begin{bmatrix} i_1 & \cdots & \cdots \\ i_2 & \cdots & \cdots \\ \vdots & & \\ i_n & \cdots & \cdots \end{bmatrix}$.

Assume $\chi_A(t) = t^n - a_1 t^{n-1} + \dots + (-1)^n a_0$, $a_i \in I$. (C-H)

$$0 = \chi_A(\text{id}) = \chi_A(\lambda_1) = \lambda_1^n - a_1 \lambda_1^{n-1} + \dots + (-1)^n a_0 = \lambda_{\chi_A(1)}.$$

Now $\chi_A(1) = 1 - i$ for some $i \in I$, so $(1 - i)M = 0$ for some $i \in I$.

If $I = J$, then $1 - i$ is a unit that acts like 0 on M .

Traditional proof of Nakayama's Lemma

Assume that $M = \langle m_1, \dots, m_m \rangle$ and $M = IM$ for some ideal I .

The identity function on M is representable as an $n \times n$ matrix A with coefficients in I .

If $m_1 = i_1 m_1 + \dots + i_n m_n$, then $A = \begin{bmatrix} i_1 & \dots & \dots \\ i_2 & \dots & \dots \\ \vdots & & \\ i_n & \dots & \dots \end{bmatrix}$.

Assume $\chi_A(t) = t^n - a_1 t^{n-1} + \dots + (-1)^n a_0$, $a_i \in I$. (C-H)

$$0 = \chi_A(\text{id}) = \chi_A(\lambda_1) = \lambda_1^n - a_1 \lambda_1^{n-1} + \dots + (-1)^n a_0 = \lambda_{\chi_A(1)}.$$

Now $\chi_A(1) = 1 - i$ for some $i \in I$, so $(1 - i)M = 0$ for some $i \in I$.

If $I = J$, then $1 - i$ is a unit that acts like 0 on M . Conclusion: $M = 0$.

Traditional proof of Nakayama's Lemma

Assume that $M = \langle m_1, \dots, m_m \rangle$ and $M = IM$ for some ideal I .

The identity function on M is representable as an $n \times n$ matrix A with coefficients in I .

If $m_1 = i_1 m_1 + \dots + i_n m_n$, then $A = \begin{bmatrix} i_1 & \cdots & \cdots \\ i_2 & \cdots & \cdots \\ \vdots & & \\ i_n & \cdots & \cdots \end{bmatrix}$.

Assume $\chi_A(t) = t^n - a_1 t^{n-1} + \dots + (-1)^n a_0$, $a_i \in I$. (C-H)

$$0 = \chi_A(\text{id}) = \chi_A(\lambda_1) = \lambda_1^n - a_1 \lambda_1^{n-1} + \dots + (-1)^n a_0 = \lambda_{\chi_A(1)}.$$

Now $\chi_A(1) = 1 - i$ for some $i \in I$, so $(1 - i)M = 0$ for some $i \in I$.

If $I = J$, then $1 - i$ is a unit that acts like 0 on M . Conclusion: $M = 0$. \square

Extra mileage from the nontraditional proof

Extra mileage from the nontraditional proof

Extra Mileage Theorem.

Extra mileage from the nontraditional proof

Extra Mileage Theorem. Let M be a finitely generated R -module.

Extra mileage from the nontraditional proof

Extra Mileage Theorem. Let M be a finitely generated R -module. If $f : M \rightarrow M$ is a surjective homomorphism, then f must be injective.

Extra mileage from the nontraditional proof

Extra Mileage Theorem. Let M be a finitely generated R -module. If $f : M \rightarrow M$ is a surjective homomorphism, then f must be injective.

Proof.

Extra mileage from the nontraditional proof

Extra Mileage Theorem. Let M be a finitely generated R -module. If $f : M \rightarrow M$ is a surjective homomorphism, then f must be injective.

Proof.

Make M an $R[t]$ -module by defining $t \cdot m = f(m)$.

Extra mileage from the nontraditional proof

Extra Mileage Theorem. Let M be a finitely generated R -module. If $f : M \rightarrow M$ is a surjective homomorphism, then f must be injective.

Proof.

Make M an $R[t]$ -module by defining $t \cdot m = f(m)$.

Let $I = (t) \triangleleft R[t]$.

Extra mileage from the nontraditional proof

Extra Mileage Theorem. Let M be a finitely generated R -module. If $f : M \rightarrow M$ is a surjective homomorphism, then f must be injective.

Proof.

Make M an $R[t]$ -module by defining $t \cdot m = f(m)$.

Let $I = (t) \triangleleft R[t]$. $M = IM$, since f is surjective.

Extra mileage from the nontraditional proof

Extra Mileage Theorem. Let M be a finitely generated R -module. If $f : M \rightarrow M$ is a surjective homomorphism, then f must be injective.

Proof.

Make M an $R[t]$ -module by defining $t \cdot m = f(m)$.

Let $I = (t) \triangleleft R[t]$. $M = IM$, since f is surjective.

By argument on previous page, $\exists i \in I = (t)$ such that $(1 - i)M = \{0\}$.

Extra mileage from the nontraditional proof

Extra Mileage Theorem. Let M be a finitely generated R -module. If $f : M \rightarrow M$ is a surjective homomorphism, then f must be injective.

Proof.

Make M an $R[t]$ -module by defining $t \cdot m = f(m)$.

Let $I = (t) \triangleleft R[t]$. $M = IM$, since f is surjective.

By argument on previous page, $\exists i \in I = (t)$ such that $(1 - i)M = \{0\}$.

Assume $i = st$ for some $s \in R[t]$.

Extra mileage from the nontraditional proof

Extra Mileage Theorem. Let M be a finitely generated R -module. If $f : M \rightarrow M$ is a surjective homomorphism, then f must be injective.

Proof.

Make M an $R[t]$ -module by defining $t \cdot m = f(m)$.

Let $I = (t) \triangleleft R[t]$. $M = IM$, since f is surjective.

By argument on previous page, $\exists i \in I = (t)$ such that $(1 - i)M = \{0\}$.

Assume $i = st$ for some $s \in R[t]$.

Choose $m \in \ker(f)$.

Extra mileage from the nontraditional proof

Extra Mileage Theorem. Let M be a finitely generated R -module. If $f : M \rightarrow M$ is a surjective homomorphism, then f must be injective.

Proof.

Make M an $R[t]$ -module by defining $t \cdot m = f(m)$.

Let $I = (t) \triangleleft R[t]$. $M = IM$, since f is surjective.

By argument on previous page, $\exists i \in I = (t)$ such that $(1 - i)M = \{0\}$.

Assume $i = st$ for some $s \in R[t]$.

Choose $m \in \ker(f)$. $0 = (1 - st)m$

Extra mileage from the nontraditional proof

Extra Mileage Theorem. Let M be a finitely generated R -module. If $f : M \rightarrow M$ is a surjective homomorphism, then f must be injective.

Proof.

Make M an $R[t]$ -module by defining $t \cdot m = f(m)$.

Let $I = (t) \triangleleft R[t]$. $M = IM$, since f is surjective.

By argument on previous page, $\exists i \in I = (t)$ such that $(1 - i)M = \{0\}$.

Assume $i = st$ for some $s \in R[t]$.

Choose $m \in \ker(f)$. $0 = (1 - st)m = m - s(f)(f(m))$

Extra mileage from the nontraditional proof

Extra Mileage Theorem. Let M be a finitely generated R -module. If $f : M \rightarrow M$ is a surjective homomorphism, then f must be injective.

Proof.

Make M an $R[t]$ -module by defining $t \cdot m = f(m)$.

Let $I = (t) \triangleleft R[t]$. $M = IM$, since f is surjective.

By argument on previous page, $\exists i \in I = (t)$ such that $(1 - i)M = \{0\}$.

Assume $i = st$ for some $s \in R[t]$.

Choose $m \in \ker(f)$. $0 = (1 - st)m = m - s(f)(f(m)) = m$,

Extra mileage from the nontraditional proof

Extra Mileage Theorem. Let M be a finitely generated R -module. If $f : M \rightarrow M$ is a surjective homomorphism, then f must be injective.

Proof.

Make M an $R[t]$ -module by defining $t \cdot m = f(m)$.

Let $I = (t) \triangleleft R[t]$. $M = IM$, since f is surjective.

By argument on previous page, $\exists i \in I = (t)$ such that $(1 - i)M = \{0\}$.

Assume $i = st$ for some $s \in R[t]$.

Choose $m \in \ker(f)$. $0 = (1 - st)m = m - s(f)(f(m)) = m$, so $\ker(f) = 0$.

Extra mileage from the nontraditional proof

Extra Mileage Theorem. Let M be a finitely generated R -module. If $f : M \rightarrow M$ is a surjective homomorphism, then f must be injective.

Proof.

Make M an $R[t]$ -module by defining $t \cdot m = f(m)$.

Let $I = (t) \triangleleft R[t]$. $M = IM$, since f is surjective.

By argument on previous page, $\exists i \in I = (t)$ such that $(1 - i)M = \{0\}$.

Assume $i = st$ for some $s \in R[t]$.

Choose $m \in \ker(f)$. $0 = (1 - st)m = m - s(f)(f(m)) = m$, so $\ker(f) = 0$.

□

Jacobson rings

If R is a commutative ring and $I \triangleleft R$, then $r(I) \subseteq J(I)$.

Jacobson rings

If R is a commutative ring and $I \triangleleft R$, then $r(I) \subseteq J(I)$. If equality holds for all I , then R is a *Jacobson ring*.

Jacobson rings

If R is a commutative ring and $I \triangleleft R$, then $r(I) \subseteq J(I)$. If equality holds for all I , then R is a *Jacobson ring*.

Examples.

① \mathbb{Z} .

Jacobson rings

If R is a commutative ring and $I \triangleleft R$, then $r(I) \subseteq J(I)$. If equality holds for all I , then R is a *Jacobson ring*.

Examples.

① \mathbb{Z} .

Jacobson rings

If R is a commutative ring and $I \triangleleft R$, then $r(I) \subseteq J(I)$. If equality holds for all I , then R is a *Jacobson ring*.

Examples.

- 1 \mathbb{Z} .
- 2 Any field.

Jacobson rings

If R is a commutative ring and $I \triangleleft R$, then $r(I) \subseteq J(I)$. If equality holds for all I , then R is a *Jacobson ring*.

Examples.

- 1 \mathbb{Z} .
- 2 Any field.

If R is a commutative ring and $I \triangleleft R$, then $r(I) \subseteq J(I)$. If equality holds for all I , then R is a *Jacobson ring*.

Examples.

- 1 \mathbb{Z} .
- 2 Any field.
- 3 $R[x]$, if R is Jacobson.

If R is a commutative ring and $I \triangleleft R$, then $r(I) \subseteq J(I)$. If equality holds for all I , then R is a *Jacobson ring*.

Examples.

- 1 \mathbb{Z} .
- 2 Any field.
- 3 $R[x]$, if R is Jacobson.

If R is a commutative ring and $I \triangleleft R$, then $r(I) \subseteq J(I)$. If equality holds for all I , then R is a *Jacobson ring*.

Examples.

- 1 \mathbb{Z} .
- 2 Any field.
- 3 $R[x]$, if R is Jacobson.
- 4 R/I , if R is Jacobson.

If R is a commutative ring and $I \triangleleft R$, then $r(I) \subseteq J(I)$. If equality holds for all I , then R is a *Jacobson ring*.

Examples.

- 1 \mathbb{Z} .
- 2 Any field.
- 3 $R[x]$, if R is Jacobson.
- 4 R/I , if R is Jacobson.

Jacobson rings

If R is a commutative ring and $I \triangleleft R$, then $r(I) \subseteq J(I)$. If equality holds for all I , then R is a *Jacobson ring*.

Examples.

- 1 \mathbb{Z} .
- 2 Any field.
- 3 $R[x]$, if R is Jacobson.
- 4 R/I , if R is Jacobson.
- 5 Any ring whose prime ideals are intersections of maximals.

If R is a commutative ring and $I \triangleleft R$, then $r(I) \subseteq J(I)$. If equality holds for all I , then R is a *Jacobson ring*.

Examples.

- 1 \mathbb{Z} .
- 2 Any field.
- 3 $R[x]$, if R is Jacobson.
- 4 R/I , if R is Jacobson.
- 5 Any ring whose prime ideals are intersections of maximals.

Jacobson rings

If R is a commutative ring and $I \triangleleft R$, then $r(I) \subseteq J(I)$. If equality holds for all I , then R is a *Jacobson ring*.

Examples.

- 1 \mathbb{Z} .
- 2 Any field.
- 3 $R[x]$, if R is Jacobson.
- 4 R/I , if R is Jacobson.
- 5 Any ring whose prime ideals are intersections of maximals.