

Uniserial domains

A domain D is *uniserial* if its ideal lattice is a chain.

A domain D is *uniserial* if its ideal lattice is a chain.

Uniserial domains

A domain D is *uniserial* if its ideal lattice is a chain.

Uniseriality induces a linear quasiorder on the multiplicative monoid

$D^\circ = \langle D - \{0\}; \cdot, 1 \rangle$ of D :

Uniserial domains

A domain D is *uniserial* if its ideal lattice is a chain.

Uniseriality induces a linear quasiorder on the multiplicative monoid

$D^\circ = \langle D - \{0\}; \cdot, 1 \rangle$ of D : $a \leq b$

Uniserial domains

A domain D is *uniserial* if its ideal lattice is a chain.

Uniseriality induces a linear quasiorder on the multiplicative monoid

$D^\circ = \langle D - \{0\}; \cdot, 1 \rangle$ of D : $a \leq b$ iff $a \mid b$

Uniserial domains

A domain D is *uniserial* if its ideal lattice is a chain.

Uniseriality induces a linear quasiorder on the multiplicative monoid

$D^\circ = \langle D - \{0\}; \cdot, 1 \rangle$ of D : $a \leq b$ iff $a \mid b$ iff $(a) \supseteq (b)$.

A domain D is *uniserial* if its ideal lattice is a chain.

Uniseriality induces a linear quasiorder on the multiplicative monoid $D^\circ = \langle D - \{0\}; \cdot, 1 \rangle$ of D : $a \leq b$ iff $a \mid b$ iff $(a) \supseteq (b)$.

If we write $a \sim b$ to mean $a \leq b$ and $b \leq a$, then $a \sim b$ holds when $(a) = (b)$,

A domain D is *uniserial* if its ideal lattice is a chain.

Uniseriality induces a linear quasiorder on the multiplicative monoid $D^\circ = \langle D - \{0\}; \cdot, 1 \rangle$ of D : $a \leq b$ iff $a \mid b$ iff $(a) \supseteq (b)$.

If we write $a \sim b$ to mean $a \leq b$ and $b \leq a$, then $a \sim b$ holds when $(a) = (b)$, which holds iff a and b differ by a unit of D .

A domain D is *uniserial* if its ideal lattice is a chain.

Uniseriality induces a linear quasiorder on the multiplicative monoid $D^\circ = \langle D - \{0\}; \cdot, 1 \rangle$ of D : $a \leq b$ iff $a \mid b$ iff $(a) \supseteq (b)$.

If we write $a \sim b$ to mean $a \leq b$ and $b \leq a$, then $a \sim b$ holds when $(a) = (b)$, which holds iff a and b differ by a unit of D . The quotient D° / D^\times is a linearly ordered, commutative, cancellative monoid.

A domain D is *uniserial* if its ideal lattice is a chain.

Uniseriality induces a linear quasiorder on the multiplicative monoid $D^\circ = \langle D - \{0\}; \cdot, 1 \rangle$ of D : $a \leq b$ iff $a \mid b$ iff $(a) \supseteq (b)$.

If we write $a \sim b$ to mean $a \leq b$ and $b \leq a$, then $a \sim b$ holds when $(a) = (b)$, which holds iff a and b differ by a unit of D . The quotient D° / D^\times is a linearly ordered, commutative, cancellative monoid.

We can extend the linear quasiorder on D° to K^\times .

A domain D is *uniserial* if its ideal lattice is a chain.

Uniseriality induces a linear quasiorder on the multiplicative monoid $D^\circ = \langle D - \{0\}; \cdot, 1 \rangle$ of D : $a \leq b$ iff $a \mid b$ iff $(a) \supseteq (b)$.

If we write $a \sim b$ to mean $a \leq b$ and $b \leq a$, then $a \sim b$ holds when $(a) = (b)$, which holds iff a and b differ by a unit of D . The quotient D° / D^\times is a linearly ordered, commutative, cancellative monoid.

We can extend the linear quasiorder on D° to K^\times . For $x, y \in K^\times$, define $x \leq y$ iff $(\exists d)(dx = y)$

A domain D is *uniserial* if its ideal lattice is a chain.

Uniseriality induces a linear quasiorder on the multiplicative monoid $D^\circ = \langle D - \{0\}; \cdot, 1 \rangle$ of D : $a \leq b$ iff $a \mid b$ iff $(a) \supseteq (b)$.

If we write $a \sim b$ to mean $a \leq b$ and $b \leq a$, then $a \sim b$ holds when $(a) = (b)$, which holds iff a and b differ by a unit of D . The quotient D° / D^\times is a linearly ordered, commutative, cancellative monoid.

We can extend the linear quasiorder on D° to K^\times . For $x, y \in K^\times$, define $x \leq y$ iff $(\exists d)(dx = y)$ iff $\langle x \rangle_D \supseteq \langle y \rangle_D$.

A domain D is *uniserial* if its ideal lattice is a chain.

Uniseriality induces a linear quasiorder on the multiplicative monoid $D^\circ = \langle D - \{0\}; \cdot, 1 \rangle$ of D : $a \leq b$ iff $a \mid b$ iff $(a) \supseteq (b)$.

If we write $a \sim b$ to mean $a \leq b$ and $b \leq a$, then $a \sim b$ holds when $(a) = (b)$, which holds iff a and b differ by a unit of D . The quotient D° / D^\times is a linearly ordered, commutative, cancellative monoid.

We can extend the linear quasiorder on D° to K^\times . For $x, y \in K^\times$, define $x \leq y$ iff $(\exists d)(dx = y)$ iff $\langle x \rangle_D \supseteq \langle y \rangle_D$. If $x = a/b \in K^\times$, then $b \mid a$ is equivalent to $x \in D$ while $a \mid b$ is equivalent to $x^{-1} \in D$.

A domain D is *uniserial* if its ideal lattice is a chain.

Uniseriality induces a linear quasiorder on the multiplicative monoid

$D^\circ = \langle D - \{0\}; \cdot, 1 \rangle$ of D : $a \leq b$ iff $a \mid b$ iff $(a) \supseteq (b)$.

If we write $a \sim b$ to mean $a \leq b$ and $b \leq a$, then $a \sim b$ holds when $(a) = (b)$, which holds iff a and b differ by a unit of D . The quotient D° / D^\times is a linearly ordered, commutative, cancellative monoid.

We can extend the linear quasiorder on D° to K^\times . For $x, y \in K^\times$, define $x \leq y$ iff $(\exists d)(dx = y)$ iff $\langle x \rangle_D \supseteq \langle y \rangle_D$. If $x = a/b \in K^\times$, then $b \mid a$ is equivalent to $x \in D$ while $a \mid b$ is equivalent to $x^{-1} \in D$. Uniseriality of D is reflected by this property of K :

Uniserial domains

A domain D is *uniserial* if its ideal lattice is a chain.

Uniseriality induces a linear quasiorder on the multiplicative monoid

$D^\circ = \langle D - \{0\}; \cdot, 1 \rangle$ of D : $a \leq b$ iff $a \mid b$ iff $(a) \supseteq (b)$.

If we write $a \sim b$ to mean $a \leq b$ and $b \leq a$, then $a \sim b$ holds when $(a) = (b)$, which holds iff a and b differ by a unit of D . The quotient D° / D^\times is a linearly ordered, commutative, cancellative monoid.

We can extend the linear quasiorder on D° to K^\times . For $x, y \in K^\times$, define $x \leq y$ iff $(\exists d)(dx = y)$ iff $\langle x \rangle_D \supseteq \langle y \rangle_D$. If $x = a/b \in K^\times$, then $b \mid a$ is equivalent to $x \in D$ while $a \mid b$ is equivalent to $x^{-1} \in D$. Uniseriality of D is reflected by this property of K : $(\forall x \in K)((x \in D) \text{ or } (x^{-1} \in D))$.

Uniserial domains

A domain D is *uniserial* if its ideal lattice is a chain.

Uniseriality induces a linear quasiorder on the multiplicative monoid

$D^\circ = \langle D - \{0\}; \cdot, 1 \rangle$ of D : $a \leq b$ iff $a \mid b$ iff $(a) \supseteq (b)$.

If we write $a \sim b$ to mean $a \leq b$ and $b \leq a$, then $a \sim b$ holds when $(a) = (b)$, which holds iff a and b differ by a unit of D . The quotient D° / D^\times is a linearly ordered, commutative, cancellative monoid.

We can extend the linear quasiorder on D° to K^\times . For $x, y \in K^\times$, define $x \leq y$ iff $(\exists d)(dx = y)$ iff $\langle x \rangle_D \supseteq \langle y \rangle_D$. If $x = a/b \in K^\times$, then $b \mid a$ is equivalent to $x \in D$ while $a \mid b$ is equivalent to $x^{-1} \in D$. Uniseriality of D is reflected by this property of K : $(\forall x \in K)((x \in D) \text{ or } (x^{-1} \in D))$. (calg1p2)

Uniserial domains

A domain D is *uniserial* if its ideal lattice is a chain.

Uniseriality induces a linear quasiorder on the multiplicative monoid $D^\circ = \langle D - \{0\}; \cdot, 1 \rangle$ of D : $a \leq b$ iff $a \mid b$ iff $(a) \supseteq (b)$.

If we write $a \sim b$ to mean $a \leq b$ and $b \leq a$, then $a \sim b$ holds when $(a) = (b)$, which holds iff a and b differ by a unit of D . The quotient D° / D^\times is a linearly ordered, commutative, cancellative monoid.

We can extend the linear quasiorder on D° to K^\times . For $x, y \in K^\times$, define $x \leq y$ iff $(\exists d)(dx = y)$ iff $\langle x \rangle_D \supseteq \langle y \rangle_D$. If $x = a/b \in K^\times$, then $b \mid a$ is equivalent to $x \in D$ while $a \mid b$ is equivalent to $x^{-1} \in D$. Uniseriality of D is reflected by this property of K : $(\forall x \in K)((x \in D) \text{ or } (x^{-1} \in D))$. (calg1p2) (D is a valuation subring of K .)

Uniserial domains

A domain D is *uniserial* if its ideal lattice is a chain.

Uniseriality induces a linear quasiorder on the multiplicative monoid

$D^\circ = \langle D - \{0\}; \cdot, 1 \rangle$ of D : $a \leq b$ iff $a \mid b$ iff $(a) \supseteq (b)$.

If we write $a \sim b$ to mean $a \leq b$ and $b \leq a$, then $a \sim b$ holds when $(a) = (b)$, which holds iff a and b differ by a unit of D . The quotient D° / D^\times is a linearly ordered, commutative, cancellative monoid.

We can extend the linear quasiorder on D° to K^\times . For $x, y \in K^\times$, define $x \leq y$ iff $(\exists d)(dx = y)$ iff $\langle x \rangle_D \supseteq \langle y \rangle_D$. If $x = a/b \in K^\times$, then $b \mid a$ is equivalent to $x \in D$ while $a \mid b$ is equivalent to $x^{-1} \in D$. Uniseriality of D is reflected by this property of K : $(\forall x \in K)((x \in D) \text{ or } (x^{-1} \in D))$. (cald1p2) (D is a valuation subring of K .)

The observations lead us to define a “value group” $G = K^\times / D^\times$, which has a total order given by divisibility $\bar{x} \leq \bar{y}$ iff $y/x \in D$.

Uniserial domains

A domain D is *uniserial* if its ideal lattice is a chain.

Uniseriality induces a linear quasiorder on the multiplicative monoid

$D^\circ = \langle D - \{0\}; \cdot, 1 \rangle$ of D : $a \leq b$ iff $a \mid b$ iff $(a) \supseteq (b)$.

If we write $a \sim b$ to mean $a \leq b$ and $b \leq a$, then $a \sim b$ holds when $(a) = (b)$, which holds iff a and b differ by a unit of D . The quotient D° / D^\times is a linearly ordered, commutative, cancellative monoid.

We can extend the linear quasiorder on D° to K^\times . For $x, y \in K^\times$, define $x \leq y$ iff $(\exists d)(dx = y)$ iff $\langle x \rangle_D \supseteq \langle y \rangle_D$. If $x = a/b \in K^\times$, then $b \mid a$ is equivalent to $x \in D$ while $a \mid b$ is equivalent to $x^{-1} \in D$. Uniseriality of D is reflected by this property of K : $(\forall x \in K)((x \in D) \text{ or } (x^{-1} \in D))$. (cald1p2) (D is a valuation subring of K .)

The observations lead us to define a “value group” $G = K^\times / D^\times$, which has a total order given by divisibility $\bar{x} \leq \bar{y}$ iff $y/x \in D$. (Or, the order on G is given by the positive cone D° / D^\times .)

Uniserial domains

A domain D is *uniserial* if its ideal lattice is a chain.

Uniseriality induces a linear quasiorder on the multiplicative monoid

$D^\circ = \langle D - \{0\}; \cdot, 1 \rangle$ of D : $a \leq b$ iff $a \mid b$ iff $(a) \supseteq (b)$.

If we write $a \sim b$ to mean $a \leq b$ and $b \leq a$, then $a \sim b$ holds when $(a) = (b)$, which holds iff a and b differ by a unit of D . The quotient D° / D^\times is a linearly ordered, commutative, cancellative monoid.

We can extend the linear quasiorder on D° to K^\times . For $x, y \in K^\times$, define $x \leq y$ iff $(\exists d)(dx = y)$ iff $\langle x \rangle_D \supseteq \langle y \rangle_D$. If $x = a/b \in K^\times$, then $b \mid a$ is equivalent to $x \in D$ while $a \mid b$ is equivalent to $x^{-1} \in D$. Uniseriality of D is reflected by this property of K : $(\forall x \in K)((x \in D) \text{ or } (x^{-1} \in D))$. (cald1p2) (D is a valuation subring of K .)

The observations lead us to define a “value group” $G = K^\times / D^\times$, which has a total order given by divisibility $\bar{x} \leq \bar{y}$ iff $y/x \in D$. (Or, the order on G is given by the positive cone D° / D^\times .) The function $v: K^\times \rightarrow G: x \mapsto \bar{x} = x/D^\times$ is a valuation on K whose valuation ring is D .

Examples

Example 1.

Example 1. $D = (\mathbb{C}[x])_{(x)}$.

Example 1. $D = (\mathbb{C}[x])_{(x)}$. The ideals of D are $D \supseteq (x) \supseteq (x^2) \supseteq \cdots \supseteq (0)$. The value group is $\cong \mathbb{Z}$.

Example 1. $D = (\mathbb{C}[x])_{(x)}$. The ideals of D are $D \supseteq (x) \supseteq (x^2) \supseteq \cdots \supseteq (0)$. The value group is $\cong \mathbb{Z}$. All Noetherian examples look like this example.

Examples

Example 1. $D = (\mathbb{C}[x])_{(x)}$. The ideals of D are $D \supseteq (x) \supseteq (x^2) \supseteq \cdots \supseteq (0)$. The value group is $\cong \mathbb{Z}$. All Noetherian examples look like this example.

Example 2.

Examples

Example 1. $D = (\mathbb{C}[x])_{(x)}$. The ideals of D are $D \supseteq (x) \supseteq (x^2) \supseteq \cdots \supseteq (0)$. The value group is $\cong \mathbb{Z}$. All Noetherian examples look like this example.

Example 2. Let $A = \mathbb{C}[x, x^{\frac{1}{2}}, x^{\frac{1}{4}}, \dots]$,

Examples

Example 1. $D = (\mathbb{C}[x])_{(x)}$. The ideals of D are $D \supseteq (x) \supseteq (x^2) \supseteq \cdots \supseteq (0)$. The value group is $\cong \mathbb{Z}$. All Noetherian examples look like this example.

Example 2. Let $A = \mathbb{C}[x, x^{\frac{1}{2}}, x^{\frac{1}{4}}, \dots]$, $\mathfrak{m} = (x, x^{\frac{1}{2}}, x^{\frac{1}{4}}, \dots)$,

Examples

Example 1. $D = (\mathbb{C}[x])_{(x)}$. The ideals of D are $D \supseteq (x) \supseteq (x^2) \supseteq \cdots \supseteq (0)$. The value group is $\cong \mathbb{Z}$. All Noetherian examples look like this example.

Example 2. Let $A = \mathbb{C}[x, x^{\frac{1}{2}}, x^{\frac{1}{4}}, \dots]$, $\mathfrak{m} = (x, x^{\frac{1}{2}}, x^{\frac{1}{4}}, \dots)$, and $D = A_{\mathfrak{m}}$.

Examples

Example 1. $D = (\mathbb{C}[x])_{(x)}$. The ideals of D are $D \supseteq (x) \supseteq (x^2) \supseteq \cdots \supseteq (0)$. The value group is $\cong \mathbb{Z}$. All Noetherian examples look like this example.

Example 2. Let $A = \mathbb{C}[x, x^{\frac{1}{2}}, x^{\frac{1}{4}}, \dots]$, $\mathfrak{m} = (x, x^{\frac{1}{2}}, x^{\frac{1}{4}}, \dots)$, and $D = A_{\mathfrak{m}}$. Any f.g. ideal of D has the form (x^q) where q is a positive dyadic rational.

Examples

Example 1. $D = (\mathbb{C}[x])_{(x)}$. The ideals of D are $D \supseteq (x) \supseteq (x^2) \supseteq \cdots \supseteq (0)$. The value group is $\cong \mathbb{Z}$. All Noetherian examples look like this example.

Example 2. Let $A = \mathbb{C}[x, x^{\frac{1}{2}}, x^{\frac{1}{4}}, \dots]$, $\mathfrak{m} = (x, x^{\frac{1}{2}}, x^{\frac{1}{4}}, \dots)$, and $D = A_{\mathfrak{m}}$. Any f.g. ideal of D has the form (x^q) where q is a positive dyadic rational. A general ideal is determined by a filter of positive dyadic rationals.

Examples

Example 1. $D = (\mathbb{C}[x])_{(x)}$. The ideals of D are $D \supseteq (x) \supseteq (x^2) \supseteq \cdots \supseteq (0)$. The value group is $\cong \mathbb{Z}$. All Noetherian examples look like this example.

Example 2. Let $A = \mathbb{C}[x, x^{\frac{1}{2}}, x^{\frac{1}{4}}, \dots]$, $\mathfrak{m} = (x, x^{\frac{1}{2}}, x^{\frac{1}{4}}, \dots)$, and $D = A_{\mathfrak{m}}$. Any f.g. ideal of D has the form (x^q) where q is a positive dyadic rational. A general ideal is determined by a filter of positive dyadic rationals. E.g., for $\alpha \in \mathbb{R}_{\geq 0}$,

Examples

Example 1. $D = (\mathbb{C}[x])_{(x)}$. The ideals of D are $D \supseteq (x) \supseteq (x^2) \supseteq \cdots \supseteq (0)$. The value group is $\cong \mathbb{Z}$. All Noetherian examples look like this example.

Example 2. Let $A = \mathbb{C}[x, x^{\frac{1}{2}}, x^{\frac{1}{4}}, \dots]$, $\mathfrak{m} = (x, x^{\frac{1}{2}}, x^{\frac{1}{4}}, \dots)$, and $D = A_{\mathfrak{m}}$. Any f.g. ideal of D has the form (x^q) where q is a positive dyadic rational. A general ideal is determined by a filter of positive dyadic rationals. E.g., for $\alpha \in \mathbb{R}_{\geq 0}$, $I_{\alpha} = \langle x^q \mid q > \alpha \rangle$

Examples

Example 1. $D = (\mathbb{C}[x])_{(x)}$. The ideals of D are $D \supseteq (x) \supseteq (x^2) \supseteq \cdots \supseteq (0)$. The value group is $\cong \mathbb{Z}$. All Noetherian examples look like this example.

Example 2. Let $A = \mathbb{C}[x, x^{\frac{1}{2}}, x^{\frac{1}{4}}, \dots]$, $\mathfrak{m} = (x, x^{\frac{1}{2}}, x^{\frac{1}{4}}, \dots)$, and $D = A_{\mathfrak{m}}$. Any f.g. ideal of D has the form (x^q) where q is a positive dyadic rational. A general ideal is determined by a filter of positive dyadic rationals. E.g., for $\alpha \in \mathbb{R}_{\geq 0}$, $I_{\alpha} = \langle x^q \mid q > \alpha \rangle$ and (x^q) , $q > 0$.

Examples

Example 1. $D = (\mathbb{C}[x])_{(x)}$. The ideals of D are $D \supseteq (x) \supseteq (x^2) \supseteq \cdots \supseteq (0)$. The value group is $\cong \mathbb{Z}$. All Noetherian examples look like this example.

Example 2. Let $A = \mathbb{C}[x, x^{\frac{1}{2}}, x^{\frac{1}{4}}, \dots]$, $\mathfrak{m} = (x, x^{\frac{1}{2}}, x^{\frac{1}{4}}, \dots)$, and $D = A_{\mathfrak{m}}$. Any f.g. ideal of D has the form (x^q) where q is a positive dyadic rational. A general ideal is determined by a filter of positive dyadic rationals. E.g., for $\alpha \in \mathbb{R}_{\geq 0}$, $I_{\alpha} = \langle x^q \mid q > \alpha \rangle$ and (x^q) , $q > 0$. The value group has size continuum.

Examples

Example 1. $D = (\mathbb{C}[x])_{(x)}$. The ideals of D are $D \supseteq (x) \supseteq (x^2) \supseteq \cdots \supseteq (0)$. The value group is $\cong \mathbb{Z}$. All Noetherian examples look like this example.

Example 2. Let $A = \mathbb{C}[x, x^{\frac{1}{2}}, x^{\frac{1}{4}}, \dots]$, $\mathfrak{m} = (x, x^{\frac{1}{2}}, x^{\frac{1}{4}}, \dots)$, and $D = A_{\mathfrak{m}}$. Any f.g. ideal of D has the form (x^q) where q is a positive dyadic rational. A general ideal is determined by a filter of positive dyadic rationals. E.g., for $\alpha \in \mathbb{R}_{\geq 0}$, $I_{\alpha} = \langle x^q \mid q > \alpha \rangle$ and (x^q) , $q > 0$. The value group has size continuum.