


# Modules





Richard Dedekind introduced the name “module” into mathematics.

Richard Dedekind introduced the name “module” into mathematics. (From “modulus of a congruence  $a \equiv b \pmod{p}$ ”)

Richard Dedekind introduced the name “module” into mathematics. (From “modulus of a congruence  $a \equiv b \pmod{p}$ ” to “modulus of a congruence  $a \equiv b \pmod{\mathfrak{p}}$ ”)

Richard Dedekind introduced the name “module” into mathematics. (From “modulus of a congruence  $a \equiv b \pmod{p}$ ” to “modulus of a congruence  $a \equiv b \pmod{\mathfrak{p}}$ ” to “additive subgroup of  $\mathbb{C}$ ”

Richard Dedekind introduced the name “module” into mathematics. (From “modulus of a congruence  $a \equiv b \pmod{p}$ ” to “modulus of a congruence  $a \equiv b \pmod{\mathfrak{p}}$ ” to “additive subgroup of  $\mathbb{C}$ ” to “module”.)

Richard Dedekind introduced the name “module” into mathematics. (From “modulus of a congruence  $a \equiv b \pmod{p}$ ” to “modulus of a congruence  $a \equiv b \pmod{\mathfrak{p}}$ ” to “additive subgroup of  $\mathbb{C}$ ” to “module”.)

**Df.**



Richard Dedekind introduced the name “module” into mathematics. (From “modulus of a congruence  $a \equiv b \pmod{p}$ ” to “modulus of a congruence  $a \equiv b \pmod{\mathfrak{p}}$ ” to “additive subgroup of  $\mathbb{C}$ ” to “module”.)

**Df.** ( $R$  a ring.)

Richard Dedekind introduced the name “module” into mathematics. (From “modulus of a congruence  $a \equiv b \pmod{p}$ ” to “modulus of a congruence  $a \equiv b \pmod{\mathfrak{p}}$ ” to “additive subgroup of  $\mathbb{C}$ ” to “module”.)

**Df.** ( $R$  a ring.) An  $R$ -module is a structure  $\langle M; +, -, 0, \{r(x) \mid r \in R\} \rangle$  such that

# Modules, I

Richard Dedekind introduced the name “module” into mathematics. (From “modulus of a congruence  $a \equiv b \pmod{p}$ ” to “modulus of a congruence  $a \equiv b \pmod{\mathfrak{p}}$ ” to “additive subgroup of  $\mathbb{C}$ ” to “module”.)

**Df.** ( $R$  a ring.) An  $R$ -module is a structure  $\langle M; +, -, 0, \{r(x) \mid r \in R\} \rangle$  such that  $\langle M; +, -, 0 \rangle$  is an abelian group and

Richard Dedekind introduced the name “module” into mathematics. (From “modulus of a congruence  $a \equiv b \pmod{p}$ ” to “modulus of a congruence  $a \equiv b \pmod{\mathfrak{p}}$ ” to “additive subgroup of  $\mathbb{C}$ ” to “module”.)

**Df.** ( $R$  a ring.) An  $R$ -module is a structure  $\langle M; +, -, 0, \{r(x) \mid r \in R\} \rangle$  such that  $\langle M; +, -, 0 \rangle$  is an abelian group and  $\rho : R \rightarrow \text{End}_{\mathbb{Z}}(M) : r \mapsto r(x)$  is a ring homomorphism.

Richard Dedekind introduced the name “module” into mathematics. (From “modulus of a congruence  $a \equiv b \pmod{p}$ ” to “modulus of a congruence  $a \equiv b \pmod{\mathfrak{p}}$ ” to “additive subgroup of  $\mathbb{C}$ ” to “module”.)

**Df.** ( $R$  a ring.) An  $R$ -module is a structure  $\langle M; +, -, 0, \{r(x) \mid r \in R\} \rangle$  such that  $\langle M; +, -, 0 \rangle$  is an abelian group and  $\rho : R \rightarrow \text{End}_{\mathbb{Z}}(M) : r \mapsto r(x)$  is a ring homomorphism.

**Df.**

Richard Dedekind introduced the name “module” into mathematics. (From “modulus of a congruence  $a \equiv b \pmod{p}$ ” to “modulus of a congruence  $a \equiv b \pmod{\mathfrak{p}}$ ” to “additive subgroup of  $\mathbb{C}$ ” to “module”.)

**Df.** ( $R$  a ring.) An  $R$ -module is a structure  $\langle M; +, -, 0, \{r(x) \mid r \in R\} \rangle$  such that  $\langle M; +, -, 0 \rangle$  is an abelian group and  $\rho : R \rightarrow \text{End}_{\mathbb{Z}}(M) : r \mapsto r(x)$  is a ring homomorphism.

**Df.** ( $R$  a  $k$ -algebra

Richard Dedekind introduced the name “module” into mathematics. (From “modulus of a congruence  $a \equiv b \pmod{p}$ ” to “modulus of a congruence  $a \equiv b \pmod{\mathfrak{p}}$ ” to “additive subgroup of  $\mathbb{C}$ ” to “module”.)

**Df.** ( $R$  a ring.) An  $R$ -module is a structure  $\langle M; +, -, 0, \{r(x) \mid r \in R\} \rangle$  such that  $\langle M; +, -, 0 \rangle$  is an abelian group and  $\rho : R \rightarrow \text{End}_{\mathbb{Z}}(M) : r \mapsto r(x)$  is a ring homomorphism.

**Df.** ( $R$  a  $k$ -algebra defined by  $k \rightarrow R : \alpha \mapsto \alpha \cdot 1$ .)

Richard Dedekind introduced the name “module” into mathematics. (From “modulus of a congruence  $a \equiv b \pmod{p}$ ” to “modulus of a congruence  $a \equiv b \pmod{\mathfrak{p}}$ ” to “additive subgroup of  $\mathbb{C}$ ” to “module”.)

**Df.** ( $R$  a ring.) An  $R$ -module is a structure  $\langle M; +, -, 0, \{r(x) \mid r \in R\} \rangle$  such that  $\langle M; +, -, 0 \rangle$  is an abelian group and  $\rho : R \rightarrow \text{End}_{\mathbb{Z}}(M) : r \mapsto r(x)$  is a ring homomorphism.

**Df.** ( $R$  a  $k$ -algebra defined by  $k \rightarrow R : \alpha \mapsto \alpha \cdot 1$ .) An  $R$ -module is a structure  $\langle V; +, -, 0, \{r(x) \mid r \in R\} \rangle$  such that



Richard Dedekind introduced the name “module” into mathematics. (From “modulus of a congruence  $a \equiv b \pmod{p}$ ” to “modulus of a congruence  $a \equiv b \pmod{\mathfrak{p}}$ ” to “additive subgroup of  $\mathbb{C}$ ” to “module”.)

**Df.** ( $R$  a ring.) An  $R$ -module is a structure  $\langle M; +, -, 0, \{r(x) \mid r \in R\} \rangle$  such that  $\langle M; +, -, 0 \rangle$  is an abelian group and  $\rho : R \rightarrow \text{End}_{\mathbb{Z}}(M) : r \mapsto r(x)$  is a ring homomorphism.

**Df.** ( $R$  a  $k$ -algebra defined by  $k \rightarrow R : \alpha \mapsto \alpha \cdot 1$ .) An  $R$ -module is a structure  $\langle V; +, -, 0, \{r(x) \mid r \in R\} \rangle$  such that  $\langle V; +, -, 0, \{\alpha(x) \mid \alpha \in k\} \rangle$  is  $k$ -vector space and

Richard Dedekind introduced the name “module” into mathematics. (From “modulus of a congruence  $a \equiv b \pmod{p}$ ” to “modulus of a congruence  $a \equiv b \pmod{\mathfrak{p}}$ ” to “additive subgroup of  $\mathbb{C}$ ” to “module”.)

**Df.** ( $R$  a ring.) An  $R$ -module is a structure  $\langle M; +, -, 0, \{r(x) \mid r \in R\} \rangle$  such that  $\langle M; +, -, 0 \rangle$  is an abelian group and  $\rho : R \rightarrow \text{End}_{\mathbb{Z}}(M) : r \mapsto r(x)$  is a ring homomorphism.

**Df.** ( $R$  a  $k$ -algebra defined by  $k \rightarrow R : \alpha \mapsto \alpha \cdot 1$ .) An  $R$ -module is a structure  $\langle V; +, -, 0, \{r(x) \mid r \in R\} \rangle$  such that  $\langle V; +, -, 0, \{\alpha(x) \mid \alpha \in k\} \rangle$  is  $k$ -vector space and  $\rho : R \rightarrow \text{End}_k(V) : r \mapsto r(x)$  is a  $k$ -algebra homomorphism.

Richard Dedekind introduced the name “module” into mathematics. (From “modulus of a congruence  $a \equiv b \pmod{p}$ ” to “modulus of a congruence  $a \equiv b \pmod{\mathfrak{p}}$ ” to “additive subgroup of  $\mathbb{C}$ ” to “module”.)

**Df.** ( $R$  a ring.) An  $R$ -module is a structure  $\langle M; +, -, 0, \{r(x) \mid r \in R\} \rangle$  such that  $\langle M; +, -, 0 \rangle$  is an abelian group and  $\rho : R \rightarrow \text{End}_{\mathbb{Z}}(M) : r \mapsto r(x)$  is a ring homomorphism.

**Df.** ( $R$  a  $k$ -algebra defined by  $k \rightarrow R : \alpha \mapsto \alpha \cdot 1$ .) An  $R$ -module is a structure  $\langle V; +, -, 0, \{r(x) \mid r \in R\} \rangle$  such that  $\langle V; +, -, 0, \{\alpha(x) \mid \alpha \in k\} \rangle$  is  $k$ -vector space and  $\rho : R \rightarrow \text{End}_k(V) : r \mapsto r(x)$  is a  $k$ -algebra homomorphism.

The class of all  $R$ -modules is equationally definable,

Richard Dedekind introduced the name “module” into mathematics. (From “modulus of a congruence  $a \equiv b \pmod{p}$ ” to “modulus of a congruence  $a \equiv b \pmod{\mathfrak{p}}$ ” to “additive subgroup of  $\mathbb{C}$ ” to “module”.)

**Df.** ( $R$  a ring.) An  $R$ -module is a structure  $\langle M; +, -, 0, \{r(x) \mid r \in R\} \rangle$  such that  $\langle M; +, -, 0 \rangle$  is an abelian group and  $\rho : R \rightarrow \text{End}_{\mathbb{Z}}(M) : r \mapsto r(x)$  is a ring homomorphism.

**Df.** ( $R$  a  $k$ -algebra defined by  $k \rightarrow R : \alpha \mapsto \alpha \cdot 1$ .) An  $R$ -module is a structure  $\langle V; +, -, 0, \{r(x) \mid r \in R\} \rangle$  such that  $\langle V; +, -, 0, \{\alpha(x) \mid \alpha \in k\} \rangle$  is  $k$ -vector space and  $\rho : R \rightarrow \text{End}_k(V) : r \mapsto r(x)$  is a  $k$ -algebra homomorphism.

The class of all  $R$ -modules is equationally definable, hence forms (the object class of) a complete and cocomplete category.

Richard Dedekind introduced the name “module” into mathematics. (From “modulus of a congruence  $a \equiv b \pmod{p}$ ” to “modulus of a congruence  $a \equiv b \pmod{\mathfrak{p}}$ ” to “additive subgroup of  $\mathbb{C}$ ” to “module”.)

**Df.** ( $R$  a ring.) An  $R$ -module is a structure  $\langle M; +, -, 0, \{r(x) \mid r \in R\} \rangle$  such that  $\langle M; +, -, 0 \rangle$  is an abelian group and  $\rho : R \rightarrow \text{End}_{\mathbb{Z}}(M) : r \mapsto r(x)$  is a ring homomorphism.

**Df.** ( $R$  a  $k$ -algebra defined by  $k \rightarrow R : \alpha \mapsto \alpha \cdot 1$ .) An  $R$ -module is a structure  $\langle V; +, -, 0, \{r(x) \mid r \in R\} \rangle$  such that  $\langle V; +, -, 0, \{\alpha(x) \mid \alpha \in k\} \rangle$  is  $k$ -vector space and  $\rho : R \rightarrow \text{End}_k(V) : r \mapsto r(x)$  is a  $k$ -algebra homomorphism.

The class of all  $R$ -modules is equationally definable, hence forms (the object class of) a complete and cocomplete category. The study of this category is the representation theory of  $R$ .

# Simplest examples

# Simplest examples

①  $(R = \mathbb{Z}) :$

# Simplest examples

①  $(R = \mathbb{Z}) :$



# Simplest examples

- ①  $(R = \mathbb{Z})$  : A  $\mathbb{Z}$ -module is an abelian group  $M$  equipped with scalar multiplications

$$n(x) = \underbrace{x + \cdots + x}_{n \text{ times}}, \quad n \in \mathbb{Z}$$

# Simplest examples

- ①  $(R = \mathbb{Z})$  : A  $\mathbb{Z}$ -module is an abelian group  $M$  equipped with scalar multiplications

$$n(x) = \underbrace{x + \cdots + x}_{n \text{ times}}, \quad n \in \mathbb{Z}$$

- ②  $(R = k, \text{ a field})$  :

# Simplest examples

- ①  $(R = \mathbb{Z})$  : A  $\mathbb{Z}$ -module is an abelian group  $M$  equipped with scalar multiplications

$$n(x) = \underbrace{x + \cdots + x}_{n \text{ times}}, \quad n \in \mathbb{Z}$$

- ②  $(R = k, \text{ a field})$  :

# Simplest examples

- ①  $(R = \mathbb{Z})$  : A  $\mathbb{Z}$ -module is an abelian group  $M$  equipped with scalar multiplications

$$n(x) = \underbrace{x + \cdots + x}_{n \text{ times}}, \quad n \in \mathbb{Z}$$

- ②  $(R = k, \text{ a field})$  : A  $k$ -module is a  $k$ -vector space  $V$ .

# Simplest examples

- ①  $(R = \mathbb{Z})$  : A  $\mathbb{Z}$ -module is an abelian group  $M$  equipped with scalar multiplications

$$n(x) = \underbrace{x + \cdots + x}_{n \text{ times}}, \quad n \in \mathbb{Z}$$

- ②  $(R = k, \text{ a field})$  : A  $k$ -module is a  $k$ -vector space  $V$ . So the representations of a field  $k$  are classified by dimension.

# Simplest examples

- ①  $(R = \mathbb{Z})$  : A  $\mathbb{Z}$ -module is an abelian group  $M$  equipped with scalar multiplications

$$n(x) = \underbrace{x + \cdots + x}_{n \text{ times}}, \quad n \in \mathbb{Z}$$

- ②  $(R = k, \text{ a field})$  : A  $k$ -module is a  $k$ -vector space  $V$ . So the representations of a field  $k$  are classified by dimension.
- ③  $(R = M_2(\mathbb{R}))$  :

# Simplest examples

- ①  $(R = \mathbb{Z})$  : A  $\mathbb{Z}$ -module is an abelian group  $M$  equipped with scalar multiplications

$$n(x) = \underbrace{x + \cdots + x}_{n \text{ times}}, \quad n \in \mathbb{Z}$$

- ②  $(R = k, \text{ a field})$  : A  $k$ -module is a  $k$ -vector space  $V$ . So the representations of a field  $k$  are classified by dimension.
- ③  $(R = M_2(\mathbb{R}))$  :

# Simplest examples

- ①  $(R = \mathbb{Z})$  : A  $\mathbb{Z}$ -module is an abelian group  $M$  equipped with scalar multiplications

$$n(x) = \underbrace{x + \cdots + x}_{n \text{ times}}, \quad n \in \mathbb{Z}$$

- ②  $(R = k, \text{ a field})$  : A  $k$ -module is a  $k$ -vector space  $V$ . So the representations of a field  $k$  are classified by dimension.
- ③  $(R = M_2(\mathbb{R}))$  :  $\mathbb{R}^2$  is a simple  $M_2(\mathbb{R})$ -module.



# Simplest examples

- ①  $(R = \mathbb{Z})$  : A  $\mathbb{Z}$ -module is an abelian group  $M$  equipped with scalar multiplications

$$n(x) = \underbrace{x + \cdots + x}_{n \text{ times}}, \quad n \in \mathbb{Z}$$

- ②  $(R = k, \text{ a field})$  : A  $k$ -module is a  $k$ -vector space  $V$ . So the representations of a field  $k$  are classified by dimension.
- ③  $(R = M_2(\mathbb{R}))$  :  $\mathbb{R}^2$  is a simple  $M_2(\mathbb{R})$ -module. Every  $M_2(\mathbb{R})$ -module is a direct sum of a uniquely determined number of copies of this simple module.

# Simplest examples

- ①  $(R = \mathbb{Z})$  : A  $\mathbb{Z}$ -module is an abelian group  $M$  equipped with scalar multiplications

$$n(x) = \underbrace{x + \cdots + x}_{n \text{ times}}, \quad n \in \mathbb{Z}$$

- ②  $(R = k, \text{ a field})$  : A  $k$ -module is a  $k$ -vector space  $V$ . So the representations of a field  $k$  are classified by dimension.
- ③  $(R = M_2(\mathbb{R}))$  :  $\mathbb{R}^2$  is a simple  $M_2(\mathbb{R})$ -module. Every  $M_2(\mathbb{R})$ -module is a direct sum of a uniquely determined number of copies of this simple module. The representations of  $M_2(\mathbb{R})$  are also classified by a single cardinal parameter.

# Simplest examples

- 1  $(R = \mathbb{Z})$  : A  $\mathbb{Z}$ -module is an abelian group  $M$  equipped with scalar multiplications

$$n(x) = \underbrace{x + \cdots + x}_{n \text{ times}}, \quad n \in \mathbb{Z}$$

- 2  $(R = k, \text{ a field})$  : A  $k$ -module is a  $k$ -vector space  $V$ . So the representations of a field  $k$  are classified by dimension.
- 3  $(R = M_2(\mathbb{R}))$  :  $\mathbb{R}^2$  is a simple  $M_2(\mathbb{R})$ -module. Every  $M_2(\mathbb{R})$ -module is a direct sum of a uniquely determined number of copies of this simple module. The representations of  $M_2(\mathbb{R})$  are also classified by a single cardinal parameter.
- 4  $(R = R)$  :

# Simplest examples

- 1  $(R = \mathbb{Z})$  : A  $\mathbb{Z}$ -module is an abelian group  $M$  equipped with scalar multiplications

$$n(x) = \underbrace{x + \cdots + x}_{n \text{ times}}, \quad n \in \mathbb{Z}$$

- 2  $(R = k, \text{ a field})$  : A  $k$ -module is a  $k$ -vector space  $V$ . So the representations of a field  $k$  are classified by dimension.
- 3  $(R = M_2(\mathbb{R}))$  :  $\mathbb{R}^2$  is a simple  $M_2(\mathbb{R})$ -module. Every  $M_2(\mathbb{R})$ -module is a direct sum of a uniquely determined number of copies of this simple module. The representations of  $M_2(\mathbb{R})$  are also classified by a single cardinal parameter.
- 4  $(R = R)$  :

# Simplest examples

- 1  $(R = \mathbb{Z})$  : A  $\mathbb{Z}$ -module is an abelian group  $M$  equipped with scalar multiplications

$$n(x) = \underbrace{x + \cdots + x}_{n \text{ times}}, \quad n \in \mathbb{Z}$$

- 2  $(R = k, \text{ a field})$  : A  $k$ -module is a  $k$ -vector space  $V$ . So the representations of a field  $k$  are classified by dimension.
- 3  $(R = M_2(\mathbb{R}))$  :  $\mathbb{R}^2$  is a simple  $M_2(\mathbb{R})$ -module. Every  $M_2(\mathbb{R})$ -module is a direct sum of a uniquely determined number of copies of this simple module. The representations of  $M_2(\mathbb{R})$  are also classified by a single cardinal parameter.
- 4  $(R = R)$  :  $R$  is an  $R$ -module.

# Simplest examples

- ①  $(R = \mathbb{Z})$  : A  $\mathbb{Z}$ -module is an abelian group  $M$  equipped with scalar multiplications

$$n(x) = \underbrace{x + \cdots + x}_{n \text{ times}}, \quad n \in \mathbb{Z}$$

- ②  $(R = k, \text{ a field})$  : A  $k$ -module is a  $k$ -vector space  $V$ . So the representations of a field  $k$  are classified by dimension.
- ③  $(R = M_2(\mathbb{R}))$  :  $\mathbb{R}^2$  is a simple  $M_2(\mathbb{R})$ -module. Every  $M_2(\mathbb{R})$ -module is a direct sum of a uniquely determined number of copies of this simple module. The representations of  $M_2(\mathbb{R})$  are also classified by a single cardinal parameter.
- ④  $(R = R)$  :  $R$  is an  $R$ -module. (Write  ${}_R R$ .)

# Simplest examples

- ①  $(R = \mathbb{Z})$  : A  $\mathbb{Z}$ -module is an abelian group  $M$  equipped with scalar multiplications

$$n(x) = \underbrace{x + \cdots + x}_{n \text{ times}}, \quad n \in \mathbb{Z}$$

- ②  $(R = k, \text{ a field})$  : A  $k$ -module is a  $k$ -vector space  $V$ . So the representations of a field  $k$  are classified by dimension.
- ③  $(R = M_2(\mathbb{R}))$  :  $\mathbb{R}^2$  is a simple  $M_2(\mathbb{R})$ -module. Every  $M_2(\mathbb{R})$ -module is a direct sum of a uniquely determined number of copies of this simple module. The representations of  $M_2(\mathbb{R})$  are also classified by a single cardinal parameter.
- ④  $(R = R)$  :  $R$  is an  $R$ -module. (Write  ${}_R R$ .) The submodules of  ${}_R R$  are the (left) ideals of  $R$ .

# Topics to discuss



# Topics to discuss

- 1 Products, coproducts.

# Topics to discuss

- ① Products, coproducts.
  - (a) Universal properties.

# Topics to discuss

- ① Products, coproducts.
  - (a) Universal properties.
  - (b) Constructions of products and coproducts.

# Topics to discuss

- ① Products, coproducts.
  - (a) Universal properties.
  - (b) Constructions of products and coproducts.
- ② Free objects.

# Topics to discuss

- ① Products, coproducts.
  - (a) Universal properties.
  - (b) Constructions of products and coproducts.
- ② Free objects.
  - (a) Construction.

# Topics to discuss

- ① Products, coproducts.
  - (a) Universal properties.
  - (b) Constructions of products and coproducts.
- ② Free objects.
  - (a) Construction.
  - (b) Presentations.

# Topics to discuss

- ① Products, coproducts.
  - (a) Universal properties.
  - (b) Constructions of products and coproducts.
- ② Free objects.
  - (a) Construction.
  - (b) Presentations.
- ③ Tensor product.

# Topics to discuss

- ① Products, coproducts.
  - (a) Universal properties.
  - (b) Constructions of products and coproducts.
- ② Free objects.
  - (a) Construction.
  - (b) Presentations.
- ③ Tensor product.
  - (a) Representable functors.



# Topics to discuss

- ① Products, coproducts.
  - (a) Universal properties.
  - (b) Constructions of products and coproducts.
- ② Free objects.
  - (a) Construction.
  - (b) Presentations.
- ③ Tensor product.
  - (a) Representable functors.
  - (b) Coalgebras.

# Topics to discuss

- ① Products, coproducts.
  - (a) Universal properties.
  - (b) Constructions of products and coproducts.
- ② Free objects.
  - (a) Construction.
  - (b) Presentations.
- ③ Tensor product.
  - (a) Representable functors.
  - (b) Coalgebras.
  - (c) Presentation of tensor product.

# Topics to discuss

- ① Products, coproducts.
  - (a) Universal properties.
  - (b) Constructions of products and coproducts.
- ② Free objects.
  - (a) Construction.
  - (b) Presentations.
- ③ Tensor product.
  - (a) Representable functors.
  - (b) Coalgebras.
  - (c) Presentation of tensor product.
- ④ Restriction and extension of scalars.

# Topics to discuss

- ① Products, coproducts.
  - (a) Universal properties.
  - (b) Constructions of products and coproducts.
- ② Free objects.
  - (a) Construction.
  - (b) Presentations.
- ③ Tensor product.
  - (a) Representable functors.
  - (b) Coalgebras.
  - (c) Presentation of tensor product.
- ④ Restriction and extension of scalars.
- ⑤ Exact sequences.

# Topics to discuss

- ① Products, coproducts.
  - (a) Universal properties.
  - (b) Constructions of products and coproducts.
- ② Free objects.
  - (a) Construction.
  - (b) Presentations.
- ③ Tensor product.
  - (a) Representable functors.
  - (b) Coalgebras.
  - (c) Presentation of tensor product.
- ④ Restriction and extension of scalars.
- ⑤ Exact sequences.
  - (a) Exact functors.

# Topics to discuss

- ① Products, coproducts.
  - (a) Universal properties.
  - (b) Constructions of products and coproducts.
- ② Free objects.
  - (a) Construction.
  - (b) Presentations.
- ③ Tensor product.
  - (a) Representable functors.
  - (b) Coalgebras.
  - (c) Presentation of tensor product.
- ④ Restriction and extension of scalars.
- ⑤ Exact sequences.
  - (a) Exact functors.
  - (b) Projectives and injectives.

# Topics to discuss

- ① Products, coproducts.
  - (a) Universal properties.
  - (b) Constructions of products and coproducts.
- ② Free objects.
  - (a) Construction.
  - (b) Presentations.
- ③ Tensor product.
  - (a) Representable functors.
  - (b) Coalgebras.
  - (c) Presentation of tensor product.
- ④ Restriction and extension of scalars.
- ⑤ Exact sequences.
  - (a) Exact functors.
  - (b) Projectives and injectives.
  - (c) Flatness.

# Topics to discuss

- ① Products, coproducts.
  - (a) Universal properties.
  - (b) Constructions of products and coproducts.
- ② Free objects.
  - (a) Construction.
  - (b) Presentations.
- ③ Tensor product.
  - (a) Representable functors.
  - (b) Coalgebras.
  - (c) Presentation of tensor product.
- ④ Restriction and extension of scalars.
- ⑤ Exact sequences.
  - (a) Exact functors.
  - (b) Projectives and injectives.
  - (c) Flatness.
- ⑥ Localization.



# Topics to discuss

- ① Products, coproducts.
  - (a) Universal properties.
  - (b) Constructions of products and coproducts.
- ② Free objects.
  - (a) Construction.
  - (b) Presentations.
- ③ Tensor product.
  - (a) Representable functors.
  - (b) Coalgebras.
  - (c) Presentation of tensor product.
- ④ Restriction and extension of scalars.
- ⑤ Exact sequences.
  - (a) Exact functors.
  - (b) Projectives and injectives.
  - (c) Flatness.
- ⑥ Localization.

# Topics to discuss

- ① Products, coproducts.
  - (a) Universal properties.
  - (b) Constructions of products and coproducts.
- ② Free objects.
  - (a) Construction.
  - (b) Presentations.
- ③ Tensor product.
  - (a) Representable functors.
  - (b) Coalgebras.
  - (c) Presentation of tensor product.
- ④ Restriction and extension of scalars.
- ⑤ Exact sequences.
  - (a) Exact functors.
  - (b) Projectives and injectives.
  - (c) Flatness.
- ⑥ Localization.

Please read A-M, Chapters 2 & 3.

# Universal morphisms

# Universal morphisms

Given a functor  $F: \mathcal{C} \rightarrow \mathcal{D}$ ,

# Universal morphisms


Given a functor  $F: \mathcal{C} \rightarrow \mathcal{D}$ , a *universal morphism* **from**  $X \in \text{Ob}(\mathcal{D})$  **to**  $F$  is a pair  $(A, f) \in \text{Ob}(\mathcal{C}) \times \text{Mor}(\mathcal{D})$  that is “universal” among all such pairs for the property that  $f: X \rightarrow F(A)$ .

# Universal morphisms

Given a functor  $F: \mathcal{C} \rightarrow \mathcal{D}$ , a *universal morphism* **from**  $X \in \text{Ob}(\mathcal{D})$  **to**  $F$  is a pair  $(A, f) \in \text{Ob}(\mathcal{C}) \times \text{Mor}(\mathcal{D})$  that is “universal” among all such pairs for the property that  $f: X \rightarrow F(A)$ . The universality means that if  $(B, g)$  is another pair with  $g: X \rightarrow F(B)$ , then there is a unique  $h: A \rightarrow B$  such that  $g = F(h) \circ f$ .


# Universal morphisms

Given a functor  $F: \mathcal{C} \rightarrow \mathcal{D}$ , a *universal morphism* **from**  $X \in \text{Ob}(\mathcal{D})$  **to**  $F$  is a pair  $(A, f) \in \text{Ob}(\mathcal{C}) \times \text{Mor}(\mathcal{D})$  that is “universal” among all such pairs for the property that  $f: X \rightarrow F(A)$ . The universality means that if  $(B, g)$  is another pair with  $g: X \rightarrow F(B)$ , then there is a unique  $h: A \rightarrow B$  such that  $g = F(h) \circ f$ .



# Universal morphisms


Given a functor  $F: \mathcal{C} \rightarrow \mathcal{D}$ , a *universal morphism* **from**  $X \in \text{Ob}(\mathcal{D})$  **to**  $F$  is a pair  $(A, f) \in \text{Ob}(\mathcal{C}) \times \text{Mor}(\mathcal{D})$  that is “universal” among all such pairs for the property that  $f: X \rightarrow F(A)$ . The universality means that if  $(B, g)$  is another pair with  $g: X \rightarrow F(B)$ , then there is a unique  $h: A \rightarrow B$  such that  $g = F(h) \circ f$ .






# Universal morphisms

Given a functor  $F: \mathcal{C} \rightarrow \mathcal{D}$ , a *universal morphism* **from**  $X \in \text{Ob}(\mathcal{D})$  **to**  $F$  is a pair  $(A, f) \in \text{Ob}(\mathcal{C}) \times \text{Mor}(\mathcal{D})$  that is “universal” among all such pairs for the property that  $f: X \rightarrow F(A)$ . The universality means that if  $(B, g)$  is another pair with  $g: X \rightarrow F(B)$ , then there is a unique  $h: A \rightarrow B$  such that  $g = F(h) \circ f$ .




# Universal morphisms

Given a functor  $F: \mathcal{C} \rightarrow \mathcal{D}$ , a *universal morphism* **from**  $X \in \text{Ob}(\mathcal{D})$  **to**  $F$  is a pair  $(A, f) \in \text{Ob}(\mathcal{C}) \times \text{Mor}(\mathcal{D})$  that is “universal” among all such pairs for the property that  $f: X \rightarrow F(A)$ . The universality means that if  $(B, g)$  is another pair with  $g: X \rightarrow F(B)$ , then there is a unique  $h: A \rightarrow B$  such that  $g = F(h) \circ f$ .




# Universal morphisms

Given a functor  $F: \mathcal{C} \rightarrow \mathcal{D}$ , a *universal morphism* **from**  $X \in \text{Ob}(\mathcal{D})$  **to**  $F$  is a pair  $(A, f) \in \text{Ob}(\mathcal{C}) \times \text{Mor}(\mathcal{D})$  that is “universal” among all such pairs for the property that  $f: X \rightarrow F(A)$ . The universality means that if  $(B, g)$  is another pair with  $g: X \rightarrow F(B)$ , then there is a unique  $h: A \rightarrow B$  such that  $g = F(h) \circ f$ .




# Universal morphisms

Given a functor  $F: \mathcal{C} \rightarrow \mathcal{D}$ , a *universal morphism* **from**  $X \in \text{Ob}(\mathcal{D})$  **to**  $F$  is a pair  $(A, f) \in \text{Ob}(\mathcal{C}) \times \text{Mor}(\mathcal{D})$  that is “universal” among all such pairs for the property that  $f: X \rightarrow F(A)$ . The universality means that if  $(B, g)$  is another pair with  $g: X \rightarrow F(B)$ , then there is a unique  $h: A \rightarrow B$  such that  $g = F(h) \circ f$ .




# Universal morphisms

Given a functor  $F: \mathcal{C} \rightarrow \mathcal{D}$ , a *universal morphism* **from**  $X \in \text{Ob}(\mathcal{D})$  **to**  $F$  is a pair  $(A, f) \in \text{Ob}(\mathcal{C}) \times \text{Mor}(\mathcal{D})$  that is “universal” among all such pairs for the property that  $f: X \rightarrow F(A)$ . The universality means that if  $(B, g)$  is another pair with  $g: X \rightarrow F(B)$ , then there is a unique  $h: A \rightarrow B$  such that  $g = F(h) \circ f$ .




# Universal morphisms

Given a functor  $F: \mathcal{C} \rightarrow \mathcal{D}$ , a *universal morphism* **from**  $X \in \text{Ob}(\mathcal{D})$  **to**  $F$  is a pair  $(A, f) \in \text{Ob}(\mathcal{C}) \times \text{Mor}(\mathcal{D})$  that is “universal” among all such pairs for the property that  $f: X \rightarrow F(A)$ . The universality means that if  $(B, g)$  is another pair with  $g: X \rightarrow F(B)$ , then there is a unique  $h: A \rightarrow B$  such that  $g = F(h) \circ f$ .



# Universal morphisms

Given a functor  $F: \mathcal{C} \rightarrow \mathcal{D}$ , a *universal morphism from*  $X \in \text{Ob}(\mathcal{D})$  **to**  $F$  is a pair  $(A, f) \in \text{Ob}(\mathcal{C}) \times \text{Mor}(\mathcal{D})$  that is “universal” among all such pairs for the property that  $f: X \rightarrow F(A)$ . The universality means that if  $(B, g)$  is another pair with  $g: X \rightarrow F(B)$ , then there is a unique  $h: A \rightarrow B$  such that  $g = F(h) \circ f$ .



For universal morphism **to**  $X$ , **from**  $F$  reverse the directions of the morphisms.

# Universal properties



# Universal properties

The statement that  $(A, f)$  is a universal morphism is the *universal property* of this pair.

# Universal properties

The statement that  $(A, f)$  is a universal morphism is the *universal property* of this pair.

**Examples.**

The statement that  $(A, f)$  is a universal morphism is the *universal property* of this pair.

## Examples.

- 1 Let  $\mathcal{C} \times \mathcal{C}$  be the category whose objects are pairs  $(A, B)$  where  $A, B \in \text{Ob}(\mathcal{C})$  and whose morphisms are pairs  $(f, g)$  where  $f, g \in \text{Mor}(\mathcal{C})$ .

The statement that  $(A, f)$  is a universal morphism is the *universal property* of this pair.

## Examples.

- 1 Let  $\mathcal{C} \times \mathcal{C}$  be the category whose objects are pairs  $(A, B)$  where  $A, B \in \text{Ob}(\mathcal{C})$  and whose morphisms are pairs  $(f, g)$  where  $f, g \in \text{Mor}(\mathcal{C})$ .

The statement that  $(A, f)$  is a universal morphism is the *universal property* of this pair.

## Examples.

- 1 Let  $\mathcal{C} \times \mathcal{C}$  be the category whose objects are pairs  $(A, B)$  where  $A, B \in \text{Ob}(\mathcal{C})$  and whose morphisms are pairs  $(f, g)$  where  $f, g \in \text{Mor}(\mathcal{C})$ . Now let  $\Delta: \mathcal{C} \rightarrow \mathcal{C} \times \mathcal{C}$  be the functor  $C \mapsto (C, C)$ ,  $e \mapsto (e, e)$ .

The statement that  $(A, f)$  is a universal morphism is the *universal property* of this pair.

## Examples.

- 1 Let  $\mathcal{C} \times \mathcal{C}$  be the category whose objects are pairs  $(A, B)$  where  $A, B \in \text{Ob}(\mathcal{C})$  and whose morphisms are pairs  $(f, g)$  where  $f, g \in \text{Mor}(\mathcal{C})$ . Now let  $\Delta: \mathcal{C} \rightarrow \mathcal{C} \times \mathcal{C}$  be the functor  $C \mapsto (C, C)$ ,  $e \mapsto (e, e)$ . For any  $X = (A, B)$  in this category, a universal morphism **from  $\Delta$  to  $X$**  is a *product*  $(P, (\pi_A, \pi_B))$  of  $A$  and  $B$ .

The statement that  $(A, f)$  is a universal morphism is the *universal property* of this pair.

## Examples.

- 1 Let  $\mathcal{C} \times \mathcal{C}$  be the category whose objects are pairs  $(A, B)$  where  $A, B \in \text{Ob}(\mathcal{C})$  and whose morphisms are pairs  $(f, g)$  where  $f, g \in \text{Mor}(\mathcal{C})$ . Now let  $\Delta: \mathcal{C} \rightarrow \mathcal{C} \times \mathcal{C}$  be the functor  $C \mapsto (C, C)$ ,  $e \mapsto (e, e)$ . For any  $X = (A, B)$  in this category, a universal morphism **from  $\Delta$  to  $X$**  is a *product*  $(P, (\pi_A, \pi_B))$  of  $A$  and  $B$ .
- 2 A universal morphism **to  $\Delta$  from  $X$**  is a *coproduct* of  $A$  and  $B$ .

The statement that  $(A, f)$  is a universal morphism is the *universal property* of this pair.

## Examples.

- 1 Let  $\mathcal{C} \times \mathcal{C}$  be the category whose objects are pairs  $(A, B)$  where  $A, B \in \text{Ob}(\mathcal{C})$  and whose morphisms are pairs  $(f, g)$  where  $f, g \in \text{Mor}(\mathcal{C})$ . Now let  $\Delta: \mathcal{C} \rightarrow \mathcal{C} \times \mathcal{C}$  be the functor  $C \mapsto (C, C)$ ,  $e \mapsto (e, e)$ . For any  $X = (A, B)$  in this category, a universal morphism **from  $\Delta$  to  $X$**  is a *product*  $(P, (\pi_A, \pi_B))$  of  $A$  and  $B$ .
- 2 A universal morphism **to  $\Delta$  from  $X$**  is a *coproduct* of  $A$  and  $B$ .



The statement that  $(A, f)$  is a universal morphism is the *universal property* of this pair.

## Examples.

- 1 Let  $\mathcal{C} \times \mathcal{C}$  be the category whose objects are pairs  $(A, B)$  where  $A, B \in \text{Ob}(\mathcal{C})$  and whose morphisms are pairs  $(f, g)$  where  $f, g \in \text{Mor}(\mathcal{C})$ . Now let  $\Delta: \mathcal{C} \rightarrow \mathcal{C} \times \mathcal{C}$  be the functor  $C \mapsto (C, C)$ ,  $e \mapsto (e, e)$ . For any  $X = (A, B)$  in this category, a universal morphism **from  $\Delta$  to  $X$**  is a *product*  $(P, (\pi_A, \pi_B))$  of  $A$  and  $B$ .
- 2 A universal morphism **to  $\Delta$  from  $X$**  is a *coproduct* of  $A$  and  $B$ .