Counting

Two basic counting principles

Two basic counting principles

Sum Rule.

Two basic counting principles

Sum Rule. (Or “Additive counting principle”)

Two basic counting principles

Sum Rule. (Or "Additive counting principle") If A and B are disjoint finite sets, then $|A \cup B| = |A| + |B|$.

Two basic counting principles

Sum Rule. (Or "Additive counting principle") If A and B are disjoint finite sets, then $|A \cup B| = |A| + |B|$.

Product Rule.

Two basic counting principles

Sum Rule. (Or "Additive counting principle") If A and B are disjoint finite sets, then $|A \cup B| = |A| + |B|$.

Product Rule. (Or "Multiplicative counting principle")

Two basic counting principles

Sum Rule. (Or "Additive counting principle") If A and B are disjoint finite sets, then $|A \cup B| = |A| + |B|$.

Product Rule. (Or "Multiplicative counting principle") If A and B are finite sets, then $|A \times B| = |A| \cdot |B|$.

Two basic counting principles

Sum Rule. (Or "Additive counting principle") If A and B are disjoint finite sets, then $|A \cup B| = |A| + |B|$.

Product Rule. (Or "Multiplicative counting principle") If A and B are finite sets, then $|A \times B| = |A| \cdot |B|$.

In general, the Sum Rule is suggested when (exclusive) "OR" is being counted, while the Product Rule is suggested when (independent) "AND" is being counted.

Size of a power set using the Sum Rule

Size of a power set using the Sum Rule

Theorem.

Size of a power set using the Sum Rule

Theorem. If $|X| = n$, then $|\mathcal{P}(X)| = 2^n$.

Size of a power set using the Sum Rule

Theorem. If $|X| = n$, then $|\mathcal{P}(X)| = 2^n$.

Proof.

Size of a power set using the Sum Rule

Theorem. If $|X| = n$, then $|\mathcal{P}(X)| = 2^n$.

Proof. Induction on $|X|$.

Size of a power set using the Sum Rule

Theorem. If $|X| = n$, then $|\mathcal{P}(X)| = 2^n$.

Proof. Induction on $|X|$.

(Base case, $n = 0$.)

Size of a power set using the Sum Rule

Theorem. If $|X| = n$, then $|\mathcal{P}(X)| = 2^n$.

Proof. Induction on $|X|$.

(Base case, $n = 0$.) $|X| = 0 \Rightarrow X = \emptyset$,

Size of a power set using the Sum Rule

Theorem. If $|X| = n$, then $|\mathcal{P}(X)| = 2^n$.

Proof. Induction on $|X|$.

(Base case, $n = 0$.) $|X| = 0 \Rightarrow X = \emptyset, \mathcal{P}(X) = \{\emptyset\}$,

Size of a power set using the Sum Rule

Theorem. If $|X| = n$, then $|\mathcal{P}(X)| = 2^n$.

Proof. Induction on $|X|$.

(Base case, $n = 0$.) $|X| = 0 \Rightarrow X = \emptyset$, $\mathcal{P}(X) = \{\emptyset\}$, $|\mathcal{P}(X)| = 1$

Size of a power set using the Sum Rule

Theorem. If $|X| = n$, then $|\mathcal{P}(X)| = 2^n$.

Proof. Induction on $|X|$.

(Base case, $n = 0$.) $|X| = 0 \Rightarrow X = \emptyset$, $\mathcal{P}(X) = \{\emptyset\}$, $|\mathcal{P}(X)| = 1 = 2^0$.

Size of a power set using the Sum Rule

Theorem. If $|X| = n$, then $|\mathcal{P}(X)| = 2^n$.

Proof. Induction on $|X|$.

(Base case, $n = 0$.) $|X| = 0 \Rightarrow X = \emptyset$, $\mathcal{P}(X) = \{\emptyset\}$, $|\mathcal{P}(X)| = 1 = 2^0$.

(Inductive step.)

Size of a power set using the Sum Rule

Theorem. If $|X| = n$, then $|\mathcal{P}(X)| = 2^n$.

Proof. Induction on $|X|$.

(Base case, $n = 0$.) $|X| = 0 \Rightarrow X = \emptyset$, $\mathcal{P}(X) = \{\emptyset\}$, $|\mathcal{P}(X)| = 1 = 2^0$.

(Inductive step.) Assume the theorem is true for sets of size n, and let's prove it for some set $X_{n+1} = \{x_1, \ldots, x_n, x_{n+1}\}$ of size $n + 1$.

Size of a power set using the Sum Rule

Theorem. If $|X| = n$, then $|\mathcal{P}(X)| = 2^n$.

Proof. Induction on $|X|$.

(Base case, $n = 0$.) $|X| = 0 \Rightarrow X = \emptyset$, $\mathcal{P}(X) = \{\emptyset\}$, $|\mathcal{P}(X)| = 1 = 2^0$.

(Inductive step.) Assume the theorem is true for sets of size n, and let's prove it for some set $X_{n+1} = \{x_1, \ldots, x_n, x_{n+1}\}$ of size $n + 1$.

Let $A \subseteq \mathcal{P}(X_{n+1})$ be the set of those subsets of X that do not contain the last element, x_{n+1}, and let B be the set of those subsets of X that do contain the last element, x_{n+1}.

Size of a power set using the Sum Rule

Theorem. If $|X| = n$, then $|\mathcal{P}(X)| = 2^n$.

Proof. Induction on $|X|$.

(Base case, $n = 0$.) $|X| = 0 \Rightarrow X = \emptyset$, $\mathcal{P}(X) = \{\emptyset\}$, $|\mathcal{P}(X)| = 1 = 2^0$.

(Inductive step.) Assume the theorem is true for sets of size n, and let's prove it for some set $X_{n+1} = \{x_1, \ldots, x_n, x_{n+1}\}$ of size $n + 1$.

Let $A \subseteq \mathcal{P}(X_{n+1})$ be the set of those subsets of X that do not contain the last element, x_{n+1}, and let B be the set of those subsets of X that do contain the last element, x_{n+1}. $|A| = |B| = |\mathcal{P}(X_n)| = 2^n$.

Size of a power set using the Sum Rule

Theorem. If $|X| = n$, then $|\mathcal{P}(X)| = 2^n$.

Proof. Induction on $|X|$.

(Base case, $n = 0$.) $|X| = 0 \Rightarrow X = \emptyset$, $\mathcal{P}(X) = \{\emptyset\}$, $|\mathcal{P}(X)| = 1 = 2^0$.

(Inductive step.) Assume the theorem is true for sets of size n, and let's prove it for some set $X_{n+1} = \{x_1, \ldots, x_n, x_{n+1}\}$ of size $n + 1$.

Let $A \subseteq \mathcal{P}(X_{n+1})$ be the set of those subsets of X that do not contain the last element, x_{n+1}, and let B be the set of those subsets of X that do contain the last element, x_{n+1}. $|A| = |B| = |\mathcal{P}(X_n)| = 2^n$. $\mathcal{P}(X_{n+1})$ is the disjoint union of A and B.

Size of a power set using the Sum Rule

Theorem. If $|X| = n$, then $|\mathcal{P}(X)| = 2^n$.

Proof. Induction on $|X|$.

(Base case, $n = 0$.) $|X| = 0 \Rightarrow X = \emptyset$, $\mathcal{P}(X) = \{\emptyset\}$, $|\mathcal{P}(X)| = 1 = 2^0$.

(Inductive step.) Assume the theorem is true for sets of size n, and let's prove it for some set $X_{n+1} = \{x_1, \ldots, x_n, x_{n+1}\}$ of size $n + 1$.

Let $A \subseteq \mathcal{P}(X_{n+1})$ be the set of those subsets of X that do not contain the last element, x_{n+1}, and let B be the set of those subsets of X that do contain the last element, x_{n+1}. $|A| = |B| = |\mathcal{P}(X_n)| = 2^n$. $\mathcal{P}(X_{n+1})$ is the disjoint union of A and B. (A subset of X_{n+1} lies in either A OR B,

Size of a power set using the Sum Rule

Theorem. If $|X| = n$, then $|\mathcal{P}(X)| = 2^n$.

Proof. Induction on $|X|$.

(Base case, $n = 0$.) $|X| = 0 \Rightarrow X = \emptyset, \mathcal{P}(X) = \{\emptyset\}, |\mathcal{P}(X)| = 1 = 2^0$.

(Inductive step.) Assume the theorem is true for sets of size n, and let's prove it for some set $X_{n+1} = \{x_1, \ldots, x_n, x_{n+1}\}$ of size $n + 1$.

Let $A \subseteq \mathcal{P}(X_{n+1})$ be the set of those subsets of X that do not contain the last element, x_{n+1}, and let B be the set of those subsets of X that do contain the last element, x_{n+1}. $|A| = |B| = |\mathcal{P}(X_n)| = 2^n$. $\mathcal{P}(X_{n+1})$ is the disjoint union of A and B. (A subset of X_{n+1} lies in either A OR B, but not both).

Size of a power set using the Sum Rule

Theorem. If $|X| = n$, then $|\mathcal{P}(X)| = 2^n$.

Proof. Induction on $|X|$.

(Base case, $n = 0$.) $|X| = 0 \Rightarrow X = \emptyset, \mathcal{P}(X) = \{\emptyset\}, |\mathcal{P}(X)| = 1 = 2^0$.

(Inductive step.) Assume the theorem is true for sets of size n, and let's prove it for some set $X_{n+1} = \{x_1, \ldots, x_n, x_{n+1}\}$ of size $n + 1$.

Let $A \subseteq \mathcal{P}(X_{n+1})$ be the set of those subsets of X that do not contain the last element, x_{n+1}, and let B be the set of those subsets of X that do contain the last element, x_{n+1}. $|A| = |B| = |\mathcal{P}(X_n)| = 2^n$. $\mathcal{P}(X_{n+1})$ is the disjoint union of A and B. (A subset of X_{n+1} lies in either A OR B, but not both). By the Sum Rule,

Size of a power set using the Sum Rule

Theorem. If $|X| = n$, then $|\mathcal{P}(X)| = 2^n$.

Proof. Induction on $|X|$.

(Base case, $n = 0$.) $|X| = 0 \Rightarrow X = \emptyset$, $\mathcal{P}(X) = \{\emptyset\}$, $|\mathcal{P}(X)| = 1 = 2^0$.

(Inductive step.) Assume the theorem is true for sets of size n, and let's prove it for some set $X_{n+1} = \{x_1, \ldots, x_n, x_{n+1}\}$ of size $n + 1$.

Let $A \subseteq \mathcal{P}(X_{n+1})$ be the set of those subsets of X that do not contain the last element, x_{n+1}, and let B be the set of those subsets of X that do contain the last element, x_{n+1}. $|A| = |B| = |\mathcal{P}(X_n)| = 2^n$. $\mathcal{P}(X_{n+1})$ is the disjoint union of A and B. (A subset of X_{n+1} lies in either A OR B, but not both). By the Sum Rule, $|\mathcal{P}(X_{n+1})| = |A| + |B| = 2^n + 2^n = 2^{n+1}$.

Size of a power set using the Sum Rule

Theorem. If $|X| = n$, then $|\mathcal{P}(X)| = 2^n$.

Proof. Induction on $|X|$.

(Base case, $n = 0$.) $|X| = 0 \Rightarrow X = \emptyset, \mathcal{P}(X) = \{\emptyset\}, |\mathcal{P}(X)| = 1 = 2^0$.

(Inductive step.) Assume the theorem is true for sets of size n, and let's prove it for some set $X_{n+1} = \{x_1, \ldots, x_n, x_{n+1}\}$ of size $n + 1$.

Let $A \subseteq \mathcal{P}(X_{n+1})$ be the set of those subsets of X that do not contain the last element, x_{n+1}, and let B be the set of those subsets of X that do contain the last element, x_{n+1}. $|A| = |B| = |\mathcal{P}(X_n)| = 2^n$. $\mathcal{P}(X_{n+1})$ is the disjoint union of A and B. (A subset of X_{n+1} lies in either A OR B, but not both). By the Sum Rule, $|\mathcal{P}(X_{n+1})| = |A| + |B| = 2^n + 2^n = 2^{n+1}$. $\square$

Size of a power set using the Product Rule

Size of a power set using the Product Rule

Theorem.

Size of a power set using the Product Rule

Theorem. If $|X| = n$, then $|\mathcal{P}(X)| = 2^n$.

Size of a power set using the Product Rule

Theorem. If $|X| = n$, then $|\mathcal{P}(X)| = 2^n$.

Proof.

Size of a power set using the Product Rule

Theorem. If $|X| = n$, then $|\mathcal{P}(X)| = 2^n$.

Proof. We count the number of subsets of X by counting the number of "descriptions" of subsets.

Size of a power set using the Product Rule

Theorem. If $|X| = n$, then $|\mathcal{P}(X)| = 2^n$.

Proof. We count the number of subsets of X by counting the number of "descriptions" of subsets. This means we will count the number of characteristic functions $c\colon X \to \{0, 1\}$.

Size of a power set using the Product Rule

Theorem. If $|X| = n$, then $|\mathcal{P}(X)| = 2^n$.

Proof. We count the number of subsets of X by counting the number of "descriptions" of subsets. This means we will count the number of characteristic functions $c\colon X \to \{0, 1\}$. A subset $S = \{2, 3, 5\}$ of $X = \{1, 2, 3, \ldots, n\}$ may be "described" by its characteristic function:

Size of a power set using the Product Rule

Theorem. If $|X| = n$, then $|\mathcal{P}(X)| = 2^n$.

Proof. We count the number of subsets of X by counting the number of "descriptions" of subsets. This means we will count the number of characteristic functions $c\colon X \to \{0, 1\}$. A subset $S = \{2, 3, 5\}$ of $X = \{1, 2, 3, \ldots, n\}$ may be "described" by its characteristic function: $(c(x) = 1 \text{ iff } x \in S)$

Size of a power set using the Product Rule

Theorem. If $|X| = n$, then $|\mathcal{P}(X)| = 2^n$.

Proof. We count the number of subsets of X by counting the number of "descriptions" of subsets. This means we will count the number of characteristic functions $c\colon X \to \{0, 1\}$. A subset $S = \{2, 3, 5\}$ of $X = \{1, 2, 3, \ldots, n\}$ may be "described" by its characteristic function: $(c(x) = 1 \text{ iff } x \in S)$

x	1	2	3	4	5	$\cdots$	n
$c(x)$	0	1	1	0	1	$\cdots$	0

Size of a power set using the Product Rule

Theorem. If $|X| = n$, then $|\mathcal{P}(X)| = 2^n$.

Proof. We count the number of subsets of X by counting the number of "descriptions" of subsets. This means we will count the number of characteristic functions $c\colon X \to \{0, 1\}$. A subset $S = \{2, 3, 5\}$ of $X = \{1, 2, 3, \ldots, n\}$ may be "described" by its characteristic function: $(c(x) = 1 \text{ iff } x \in S)$

x	1	2	3	4	5	$\cdots$	n
$c(x)$	0	1	1	0	1	$\cdots$	0

There are 2 choices for $c(1)$, 2 independent choices for $c(2)$, $\ldots$, 2 independent choices for $c(n)$, so $|X| = 2 \cdot 2 \cdots 2 = 2^n$.

Size of a power set using the Product Rule

Theorem. If $|X| = n$, then $|\mathcal{P}(X)| = 2^n$.

Proof. We count the number of subsets of X by counting the number of "descriptions" of subsets. This means we will count the number of characteristic functions $c\colon X \to \{0, 1\}$. A subset $S = \{2, 3, 5\}$ of $X = \{1, 2, 3, \ldots, n\}$ may be "described" by its characteristic function: $(c(x) = 1 \text{ iff } x \in S)$

x	1	2	3	4	5	$\cdots$	n
$c(x)$	0	1	1	0	1	$\cdots$	0

There are 2 choices for $c(1)$, 2 independent choices for $c(2)$, $\ldots$, 2 independent choices for $c(n)$, so $|X| = 2 \cdot 2 \cdots 2 = 2^n$. $\square$

Size of a power set using the Product Rule

Theorem. If $|X| = n$, then $|\mathcal{P}(X)| = 2^n$.

Proof. We count the number of subsets of X by counting the number of "descriptions" of subsets. This means we will count the number of characteristic functions $c\colon X \to \{0, 1\}$. A subset $S = \{2, 3, 5\}$ of $X = \{1, 2, 3, \ldots, n\}$ may be "described" by its characteristic function: $(c(x) = 1 \text{ iff } x \in S)$

x	1	2	3	4	5	$\cdots$	n
$c(x)$	0	1	1	0	1	$\cdots$	0

There are 2 choices for $c(1)$, 2 independent choices for $c(2)$, $\ldots$, 2 independent choices for $c(n)$, so $|X| = 2 \cdot 2 \cdots 2 = 2^n$. $\square$

Here we used the fact that a subset $S \subseteq X$ can be described by specifying whether $1 \in S$ AND specifying whether $2 \in X$, etc.

Exercise! Colorado license plates

Exercise! Colorado license plates

Exercise! Colorado license plates

From 2000-2015, CO license plates followed the pattern 000-AAA.

Exercise! Colorado license plates

From 2000-2015, CO license plates followed the pattern 000-AAA.
From 2015-2018, CO license plates followed the pattern AAA-000.

Exercise! Colorado license plates

From 2000-2015, CO license plates followed the pattern 000-AAA.
From 2015-2018, CO license plates followed the pattern AAA-000.
From 2018-present, CO license plates follow the pattern AAA-A00.

Exercise! Colorado license plates

From 2000-2015, CO license plates followed the pattern 000-AAA.
From 2015-2018, CO license plates followed the pattern AAA-000.
From 2018-present, CO license plates follow the pattern AAA-A00.







Exercise! Colorado license plates

From 2000-2015, CO license plates followed the pattern 000-AAA.
From 2015-2018, CO license plates followed the pattern AAA-000.
From 2018-present, CO license plates follow the pattern AAA-A00.







Questions.

Exercise! Colorado license plates

From 2000-2015, CO license plates followed the pattern 000-AAA.
From 2015-2018, CO license plates followed the pattern AAA-000.
From 2018-present, CO license plates follow the pattern AAA-A00.







Questions.

1. How many possible plates for 2000-2015?

Exercise! Colorado license plates

From 2000-2015, CO license plates followed the pattern 000-AAA.
From 2015-2018, CO license plates followed the pattern AAA-000.
From 2018-present, CO license plates follow the pattern AAA-A00.







Questions.

1. How many possible plates for 2000-2015?

Exercise! Colorado license plates

From 2000-2015, CO license plates followed the pattern 000-AAA.
From 2015-2018, CO license plates followed the pattern AAA-000.
From 2018-present, CO license plates follow the pattern AAA-A00.







Questions.

1. How many possible plates for 2000-2015?

   10

Exercise! Colorado license plates

From 2000-2015, CO license plates followed the pattern 000-AAA.
From 2015-2018, CO license plates followed the pattern AAA-000.
From 2018-present, CO license plates follow the pattern AAA-A00.







Questions.

1. How many possible plates for 2000-2015?

   10×10

Exercise! Colorado license plates

From 2000-2015, CO license plates followed the pattern 000-AAA.
From 2015-2018, CO license plates followed the pattern AAA-000.
From 2018-present, CO license plates follow the pattern AAA-A00.







Questions.

1. How many possible plates for 2000-2015?

   $10 \times 10 \times 10$

Exercise! Colorado license plates

From 2000-2015, CO license plates followed the pattern 000-AAA.
From 2015-2018, CO license plates followed the pattern AAA-000.
From 2018-present, CO license plates follow the pattern AAA-A00.







Questions.

1. How many possible plates for 2000-2015?

   $10 \times 10 \times 10 \times 26$

Exercise! Colorado license plates

From 2000-2015, CO license plates followed the pattern 000-AAA.
From 2015-2018, CO license plates followed the pattern AAA-000.
From 2018-present, CO license plates follow the pattern AAA-A00.







Questions.

1. How many possible plates for 2000-2015?

   $10 \times 10 \times 10 \times 26 \times 26$

Exercise! Colorado license plates

From 2000-2015, CO license plates followed the pattern 000-AAA.
From 2015-2018, CO license plates followed the pattern AAA-000.
From 2018-present, CO license plates follow the pattern AAA-A00.







Questions.

1. How many possible plates for 2000-2015?

   $10 \times 10 \times 10 \times 26 \times 26 \times 26$

Exercise! Colorado license plates

From 2000-2015, CO license plates followed the pattern 000-AAA.
From 2015-2018, CO license plates followed the pattern AAA-000.
From 2018-present, CO license plates follow the pattern AAA-A00.







Questions.

1. How many possible plates for 2000-2015?

   $10 \times 10 \times 10 \times 26 \times 26 \times 26 = 10^3 \cdot 26^3$

Exercise! Colorado license plates

From 2000-2015, CO license plates followed the pattern 000-AAA.
From 2015-2018, CO license plates followed the pattern AAA-000.
From 2018-present, CO license plates follow the pattern AAA-A00.







Questions.

1. How many possible plates for 2000-2015?

   $10 \times 10 \times 10 \times 26 \times 26 \times 26 = 10^3 \cdot 26^3 = 17,576,000.$

Exercise! Colorado license plates

From 2000-2015, CO license plates followed the pattern 000-AAA.
From 2015-2018, CO license plates followed the pattern AAA-000.
From 2018-present, CO license plates follow the pattern AAA-A00.







Questions.

1. How many possible plates for 2000-2015?

   $10 \times 10 \times 10 \times 26 \times 26 \times 26 = 10^3 \cdot 26^3 = 17,576,000.$

2. How many possible plates for 2000-2018?

Exercise! Colorado license plates

From 2000-2015, CO license plates followed the pattern 000-AAA.
From 2015-2018, CO license plates followed the pattern AAA-000.
From 2018-present, CO license plates follow the pattern AAA-A00.







Questions.

1. How many possible plates for 2000-2015?

   $10 \times 10 \times 10 \times 26 \times 26 \times 26 = 10^3 \cdot 26^3 = 17,576,000.$

2. How many possible plates for 2000-2018?

Exercise! Colorado license plates

From 2000-2015, CO license plates followed the pattern 000-AAA.
From 2015-2018, CO license plates followed the pattern AAA-000.
From 2018-present, CO license plates follow the pattern AAA-A00.







Questions.

1. How many possible plates for 2000-2015?

   $10 \times 10 \times 10 \times 26 \times 26 \times 26 = 10^3 \cdot 26^3 = 17,576,000.$

2. How many possible plates for 2000-2018?

   $10^3 \cdot 26^3$

Exercise! Colorado license plates

From 2000-2015, CO license plates followed the pattern 000-AAA.
From 2015-2018, CO license plates followed the pattern AAA-000.
From 2018-present, CO license plates follow the pattern AAA-A00.







Questions.

1. How many possible plates for 2000-2015?

   $10 \times 10 \times 10 \times 26 \times 26 \times 26 = 10^3 \cdot 26^3 = 17,576,000.$

2. How many possible plates for 2000-2018?

   $10^3 \cdot 26^3 + 26^3 \cdot 10^3$

Exercise! Colorado license plates

From 2000-2015, CO license plates followed the pattern 000-AAA.
From 2015-2018, CO license plates followed the pattern AAA-000.
From 2018-present, CO license plates follow the pattern AAA-A00.







Questions.

1. How many possible plates for 2000-2015?

   $10 \times 10 \times 10 \times 26 \times 26 \times 26 = 10^3 \cdot 26^3 = 17,576,000.$

2. How many possible plates for 2000-2018?

   $10^3 \cdot 26^3 + 26^3 \cdot 10^3 = 2 \cdot 10^3 \cdot 26^3$

Exercise! Colorado license plates

From 2000-2015, CO license plates followed the pattern 000-AAA.
From 2015-2018, CO license plates followed the pattern AAA-000.
From 2018-present, CO license plates follow the pattern AAA-A00.







Questions.

1. How many possible plates for 2000-2015?

   $10 \times 10 \times 10 \times 26 \times 26 \times 26 = 10^3 \cdot 26^3 = 17,576,000.$

2. How many possible plates for 2000-2018?

   $10^3 \cdot 26^3 + 26^3 \cdot 10^3 = 2 \cdot 10^3 \cdot 26^3 = 35,152,000.$

Exercise! Colorado license plates

From 2000-2015, CO license plates followed the pattern 000-AAA.
From 2015-2018, CO license plates followed the pattern AAA-000.
From 2018-present, CO license plates follow the pattern AAA-A00.







Questions.

1. How many possible plates for 2000-2015?

   $10 \times 10 \times 10 \times 26 \times 26 \times 26 = 10^3 \cdot 26^3 = 17,576,000.$

2. How many possible plates for 2000-2018?

   $10^3 \cdot 26^3 + 26^3 \cdot 10^3 = 2 \cdot 10^3 \cdot 26^3 = 35,152,000.$

3. How many possible plates for 2000-present?

Exercise! Colorado license plates

From 2000-2015, CO license plates followed the pattern 000-AAA.
From 2015-2018, CO license plates followed the pattern AAA-000.
From 2018-present, CO license plates follow the pattern AAA-A00.







Questions.

1. How many possible plates for 2000-2015?

   $10 \times 10 \times 10 \times 26 \times 26 \times 26 = 10^3 \cdot 26^3 = 17,576,000.$

2. How many possible plates for 2000-2018?

   $10^3 \cdot 26^3 + 26^3 \cdot 10^3 = 2 \cdot 10^3 \cdot 26^3 = 35,152,000.$

3. How many possible plates for 2000-present?

Exercise! Colorado license plates

From 2000-2015, CO license plates followed the pattern 000-AAA.
From 2015-2018, CO license plates followed the pattern AAA-000.
From 2018-present, CO license plates follow the pattern AAA-A00.







Questions.

1. How many possible plates for 2000-2015?
   $10 \times 10 \times 10 \times 26 \times 26 \times 26 = 10^3 \cdot 26^3 = 17,576,000.$
2. How many possible plates for 2000-2018?
   $10^3 \cdot 26^3 + 26^3 \cdot 10^3 = 2 \cdot 10^3 \cdot 26^3 = 35,152,000.$
3. How many possible plates for 2000-present?
   $10^3 \cdot 26^3$

Exercise! Colorado license plates

From 2000-2015, CO license plates followed the pattern 000-AAA.
From 2015-2018, CO license plates followed the pattern AAA-000.
From 2018-present, CO license plates follow the pattern AAA-A00.







Questions.

1. How many possible plates for 2000-2015?

   $10 \times 10 \times 10 \times 26 \times 26 \times 26 = 10^3 \cdot 26^3 = 17,576,000.$

2. How many possible plates for 2000-2018?

   $10^3 \cdot 26^3 + 26^3 \cdot 10^3 = 2 \cdot 10^3 \cdot 26^3 = 35,152,000.$

3. How many possible plates for 2000-present?

   $10^3 \cdot 26^3 + 26^3 \cdot 10^3$

Exercise! Colorado license plates

From 2000-2015, CO license plates followed the pattern 000-AAA.
From 2015-2018, CO license plates followed the pattern AAA-000.
From 2018-present, CO license plates follow the pattern AAA-A00.







Questions.

1. How many possible plates for 2000-2015?

   $10 \times 10 \times 10 \times 26 \times 26 \times 26 = 10^3 \cdot 26^3 = 17,576,000.$

2. How many possible plates for 2000-2018?

   $10^3 \cdot 26^3 + 26^3 \cdot 10^3 = 2 \cdot 10^3 \cdot 26^3 = 35,152,000.$

3. How many possible plates for 2000-present?

   $10^3 \cdot 26^3 + 26^3 \cdot 10^3 + 26^4 \cdot 10^2$

Exercise! Colorado license plates

From 2000-2015, CO license plates followed the pattern 000-AAA.
From 2015-2018, CO license plates followed the pattern AAA-000.
From 2018-present, CO license plates follow the pattern AAA-A00.







Questions.

1. How many possible plates for 2000-2015?
   $10 \times 10 \times 10 \times 26 \times 26 \times 26 = 10^3 \cdot 26^3 = 17,576,000.$
2. How many possible plates for 2000-2018?
   $10^3 \cdot 26^3 + 26^3 \cdot 10^3 = 2 \cdot 10^3 \cdot 26^3 = 35,152,000.$
3. How many possible plates for 2000-present?
   $10^3 \cdot 26^3 + 26^3 \cdot 10^3 + 26^4 \cdot 10^2 = 80,849,600.$

Exercise! Colorado license plates

From 2000-2015, CO license plates followed the pattern 000-AAA.
From 2015-2018, CO license plates followed the pattern AAA-000.
From 2018-present, CO license plates follow the pattern AAA-A00.







Questions.

1. How many possible plates for 2000-2015?
   $10 \times 10 \times 10 \times 26 \times 26 \times 26 = 10^3 \cdot 26^3 = 17,576,000.$
2. How many possible plates for 2000-2018?
   $10^3 \cdot 26^3 + 26^3 \cdot 10^3 = 2 \cdot 10^3 \cdot 26^3 = 35,152,000.$
3. How many possible plates for 2000-present?
   $10^3 \cdot 26^3 + 26^3 \cdot 10^3 + 26^4 \cdot 10^2 = 80,849,600.$

A 2024 change in Colorado license plates!

A 2024 change in Colorado license plates!

In 2024, the letter “O” was removed from the fourth character position in the AAA-A00 format to reduce confusion with the number “0”.

A 2024 change in Colorado license plates!

In 2024, the letter “O” was removed from the fourth character position in the AAA-A00 format to reduce confusion with the number “0”.

Question.

A 2024 change in Colorado license plates!

In 2024, the letter “O” was removed from the fourth character position in the AAA-A00 format to reduce confusion with the number “0”.

Question.

1. How many possible plates for 2024-present?

A 2024 change in Colorado license plates!

In 2024, the letter “O” was removed from the fourth character position in the AAA-A00 format to reduce confusion with the number “0”.

Question.

1. How many possible plates for 2024-present?

A 2024 change in Colorado license plates!

In 2024, the letter "O" was removed from the fourth character position in the AAA-A00 format to reduce confusion with the number "0".

Question.

1. How many possible plates for 2024-present?

   $26^3 \cdot 25 \cdot 10^2$

A 2024 change in Colorado license plates!

In 2024, the letter "O" was removed from the fourth character position in the AAA-A00 format to reduce confusion with the number "0".

Question.

1. How many possible plates for 2024-present?

   $26^3 \cdot 25 \cdot 10^2 = 43,940,000.$

A 2024 change in Colorado license plates!

In 2024, the letter "O" was removed from the fourth character position in the AAA-A00 format to reduce confusion with the number "0".

Question.

1. How many possible plates for 2024-present?

   $26^3 \cdot 25 \cdot 10^2 = 43,940,000.$

Number of functions from k to n

Number of functions from k to n

Theorem.

Number of functions from k to n

Theorem. The number of functions $f : k \to n$ is n^k.

Number of functions from k to n

Theorem. The number of functions $f : k \to n$ is n^k.

Proof.

Number of functions from k to n

Theorem. The number of functions $f : k \to n$ is n^k.

Proof. Count descriptions of such functions.

Number of functions from k to n

Theorem. The number of functions $f : k \to n$ is n^k.

Proof. Count descriptions of such functions.

x	1	2	3	4	5	$\cdots$	k
$f(x)$	0	7	1	0	2	$\cdots$	3

Number of functions from k to n

Theorem. The number of functions $f : k \to n$ is n^k.

Proof. Count descriptions of such functions.

x	1	2	3	4	5	$\cdots$	k
$f(x)$	0	7	1	0	2	$\cdots$	3

There are n choices for $f(0)$, n (independent) choices for $f(1)$, etc.

Number of functions from k to n

Theorem. The number of functions $f : k \to n$ is n^k.

Proof. Count descriptions of such functions.

x	1	2	3	4	5	$\cdots$	k
$f(x)$	0	7	1	0	2	$\cdots$	3

There are n choices for $f(0)$, n (independent) choices for $f(1)$, etc. Hence the number of functions is $n \cdot n \cdots n = n^k$.

Number of functions from k to n

Theorem. The number of functions $f : k \to n$ is n^k.

Proof. Count descriptions of such functions.

x	1	2	3	4	5	$\cdots$	k
$f(x)$	0	7	1	0	2	$\cdots$	3

There are n choices for $f(0)$, n (independent) choices for $f(1)$, etc. Hence the number of functions is $n \cdot n \cdots n = n^k$. $\square$

Number of bijections from k to n

Number of bijections from k to n

Theorem.

Number of bijections from k to n

Theorem. The number of bijections from k to n is 0 if $k \neq n$; otherwise it is $n!$.

Number of bijections from k to n

Theorem. The number of bijections from k to n is 0 if $k \neq n$; otherwise it is $n!$.

Proof.

Number of bijections from k to n

Theorem. The number of bijections from k to n is 0 if $k \neq n$; otherwise it is $n!$.

Proof. Count descriptions of such functions. ...

Number of bijections from k to n

Theorem. The number of bijections from k to n is 0 if $k \neq n$; otherwise it is $n!$.

Proof. Count descriptions of such functions. . . .

Definition.

Number of bijections from k to n

Theorem. The number of bijections from k to n is 0 if $k \neq n$; otherwise it is $n!$.

Proof. Count descriptions of such functions. ...

Definition. A bijection $b\colon X \to X$ from a set X to itself is called a **permutation** of X.

Exercise! Create poetry with math!

Exercise! Create poetry with math!

Roses are red.

Exercise! Create poetry with math!

Roses are red.
Violets are blue.

Exercise! Create poetry with math!

Roses are red.
Violets are blue.
Sugar is sweet.

Exercise! Create poetry with math!

Roses are red.
Violets are blue.
Sugar is sweet.
And so are you.

Exercise! Create poetry with math!

Roses are red.
Violets are blue.
Sugar is sweet.
And so are you.

Questions.

Exercise! Create poetry with math!

Roses are red.
Violets are blue.
Sugar is sweet.
And so are you.

Questions.

1. How many ways can we permute the lines of this poem to create a new poem?

Exercise! Create poetry with math!

Roses are red.
Violets are blue.
Sugar is sweet.
And so are you.

Questions.

1. How many ways can we permute the lines of this poem to create a new poem?

Exercise! Create poetry with math!

Roses are red.
Violets are blue.
Sugar is sweet.
And so are you.

Questions.

1. How many ways can we permute the lines of this poem to create a new poem? $4! = 24$

Exercise! Create poetry with math!

Roses are red.
Violets are blue.
Sugar is sweet.
And so are you.

Questions.

1. How many ways can we permute the lines of this poem to create a new poem? $4! = 24$
2. How many ways can we permute the lines to create a grammatically correct poem?

Exercise! Create poetry with math!

Roses are red.
Violets are blue.
Sugar is sweet.
And so are you.

Questions.

1. How many ways can we permute the lines of this poem to create a new poem? $4! = 24$
2. How many ways can we permute the lines to create a grammatically correct poem?

Exercise! Create poetry with math!

Roses are red.
Violets are blue.
Sugar is sweet.
And so are you.

Questions.

1. How many ways can we permute the lines of this poem to create a new poem? $4! = 24$

2. How many ways can we permute the lines to create a grammatically correct poem? (The problem is that a poem should not begin with "and", since "and" is a conjuction.)

Exercise! Create poetry with math!

Roses are red.
Violets are blue.
Sugar is sweet.
And so are you.

Questions.

1. How many ways can we permute the lines of this poem to create a new poem? $4! = 24$

2. How many ways can we permute the lines to create a grammatically correct poem? (The problem is that a poem should not begin with "and", since "and" is a conjuction.) $3 \cdot 3! = 18$

Exercise! Create poetry with math!

Roses are red.
Violets are blue.
Sugar is sweet.
And so are you.

Questions.

1. How many ways can we permute the lines of this poem to create a new poem? $4! = 24$
2. How many ways can we permute the lines to create a grammatically correct poem? (The problem is that a poem should not begin with "and", since "and" is a conjuction.) $3 \cdot 3! = 18$
3. How many ways can we permute the lines to create a grammatically correct poem that rhymes?

Exercise! Create poetry with math!

Roses are red.
Violets are blue.
Sugar is sweet.
And so are you.

Questions.

1. How many ways can we permute the lines of this poem to create a new poem? $4! = 24$
2. How many ways can we permute the lines to create a grammatically correct poem? (The problem is that a poem should not begin with "and", since "and" is a conjuction.) $3 \cdot 3! = 18$
3. How many ways can we permute the lines to create a grammatically correct poem that rhymes?

Exercise! Create poetry with math!

Roses are red.
Violets are blue.
Sugar is sweet.
And so are you.

Questions.

1. How many ways can we permute the lines of this poem to create a new poem? $4! = 24$
2. How many ways can we permute the lines to create a grammatically correct poem? (The problem is that a poem should not begin with "and", since "and" is a conjuction.) $3 \cdot 3! = 18$
3. How many ways can we permute the lines to create a grammatically correct poem that rhymes? $2! \cdot 2! = 4$

Exercise! Create poetry with math!

Roses are red.
Violets are blue.
Sugar is sweet.
And so are you.

Questions.

1. How many ways can we permute the lines of this poem to create a new poem? $4! = 24$
2. How many ways can we permute the lines to create a grammatically correct poem? (The problem is that a poem should not begin with "and", since "and" is a conjuction.) $3 \cdot 3! = 18$
3. How many ways can we permute the lines to create a grammatically correct poem that rhymes? $2! \cdot 2! = 4$

Number of injections from k to n

Number of injections from k to n

Theorem.

Number of injections from k to n

Theorem. The number of injections from k to n is

Number of injections from k to n

Theorem. The number of injections from k to n is

$$(n)_k = \underbrace{n \cdot (n-1) \cdots (n-k+1)}_{k \text{ factors}} = \frac{n!}{(n-k)!}.$$

Number of injections from k to n

Theorem. The number of injections from k to n is

$$(n)_k = \underbrace{n \cdot (n-1) \cdots (n-k+1)}_{k \text{ factors}} = \frac{n!}{(n-k)!}.$$

$(n)_k$ is called a "falling factorial".

Number of injections from k to n

Theorem. The number of injections from k to n is

$$(n)_k = \underbrace{n \cdot (n-1) \cdots (n-k+1)}_{k \text{ factors}} = \frac{n!}{(n-k)!}.$$

$(n)_k$ is called a "falling factorial".

Proof.

Number of injections from k to n

Theorem. The number of injections from k to n is

$$(n)_k = \underbrace{n \cdot (n-1) \cdots (n-k+1)}_{k \text{ factors}} = \frac{n!}{(n-k)!}.$$

$(n)_k$ is called a "falling factorial".

Proof. Count descriptions of such functions. . . .

Double counting (or over counting)

Double counting (or over counting)

Exercises.

Double counting (or over counting)

Exercises.

1. How many cows do I have?

Double counting (or over counting)

Exercises.

1. How many cows do I have?

Double counting (or over counting)

Exercises.

1. How many cows do I have?

   Count the legs and divide by 4.

Double counting (or over counting)

Exercises.

1. How many cows do I have?

   Count the legs and divide by 4.

$$\#\text{cows} \quad = \quad \#\text{legs}/4$$

Double counting (or over counting)

Exercises.

1. How many cows do I have?

   Count the legs and divide by 4.

$$\#\text{cows} \quad = \quad \#\text{legs}/4$$

   Let X be a finite set, let E be a uniform equivalence relation on X, and let X/E be the set of E-classes.

$$|X/E| \quad = \quad |X|/(\text{common size of } E\text{-classes}).$$

Double counting (or over counting)

Exercises.

1. How many cows do I have?

   Count the legs and divide by 4.

   $$\#\text{cows} \quad = \quad \#\text{legs}/4$$

   Let X be a finite set, let E be a uniform equivalence relation on X, and let X/E be the set of E-classes.

   $$|X/E| \quad = \quad |X|/(\text{common size of } E\text{-classes}).$$

2. Show that the number of 2-element subsets of n is $n(n-1)/2$.

Double counting (or over counting)

Exercises.

1. How many cows do I have?

   Count the legs and divide by 4.

   $$\#\text{cows} \quad = \quad \#\text{legs}/4$$

   Let X be a finite set, let E be a uniform equivalence relation on X, and let X/E be the set of E-classes.

   $$|X/E| \quad = \quad |X|/(\text{common size of } E\text{-classes}).$$

2. Show that the number of 2-element subsets of n is $n(n-1)/2$.

Double counting (or over counting)

Exercises.

1. How many cows do I have?

   Count the legs and divide by 4.

   $$\#\text{cows} \quad = \quad \#\text{legs}/4$$

   Let X be a finite set, let E be a uniform equivalence relation on X, and let X/E be the set of E-classes.

   $$|X/E| \quad = \quad |X|/(\text{common size of } E\text{-classes}).$$

2. Show that the number of 2-element subsets of n is $n(n-1)/2$.

3. Show, more generally, that the number of k-element subsets of n is $(n)_k/k! = \binom{n}{k}$.

Double counting (or over counting)

Exercises.

1. How many cows do I have?

   Count the legs and divide by 4.

   $$\#\text{cows} \quad = \quad \#\text{legs}/4$$

   Let X be a finite set, let E be a uniform equivalence relation on X, and let X/E be the set of E-classes.

   $$|X/E| \quad = \quad |X|/(\text{common size of } E\text{-classes}).$$

2. Show that the number of 2-element subsets of n is $n(n-1)/2$.

3. Show, more generally, that the number of k-element subsets of n is $(n)_k/k! = \binom{n}{k}$.

Double counting (or over counting)

Exercises.

1. How many cows do I have?

   Count the legs and divide by 4.

   $$\#\text{cows} \quad = \quad \#\text{legs}/4$$

   Let X be a finite set, let E be a uniform equivalence relation on X, and let X/E be the set of E-classes.

   $$|X/E| \quad = \quad |X|/(\text{common size of } E\text{-classes}).$$

2. Show that the number of 2-element subsets of n is $n(n-1)/2$.

3. Show, more generally, that the number of k-element subsets of n is $(n)_k/k! = \binom{n}{k}$.