

Adjoint and Limits

Homological Algebra

March 17, 2010

Thm. Right adjoints preserve limits; left adjoints preserve colimits.

Adjoint on limits


Thm. Right adjoints preserve limits; left adjoints preserve colimits.

Proof. Assume $\varphi: \text{Hom}_{\mathcal{C}}(F(Y), X) \rightarrow \text{Hom}_{\mathcal{D}}(Y, G(X))$ is a natural isomorphism and $D: J \rightarrow \mathcal{C}$ is a diagram with $\varprojlim D = (L, \lambda)$.

Adjoint on limits

Thm. Right adjoints preserve limits; left adjoints preserve colimits.


Proof. Assume $\varphi: \text{Hom}_{\mathcal{C}}(F(Y), X) \rightarrow \text{Hom}_{\mathcal{D}}(Y, G(X))$ is a natural isomorphism and $D: J \rightarrow \mathcal{C}$ is a diagram with $\varprojlim D = (L, \lambda)$.



Adjoint on limits

Thm. Right adjoints preserve limits; left adjoints preserve colimits.


Proof. Assume $\varphi: \text{Hom}_{\mathcal{C}}(F(Y), X) \rightarrow \text{Hom}_{\mathcal{D}}(Y, G(X))$ is a natural isomorphism and $D: J \rightarrow \mathcal{C}$ is a diagram with $\varprojlim D = (L, \lambda)$.



Adjoint on limits

Thm. Right adjoints preserve limits; left adjoints preserve colimits.


Proof. Assume $\varphi: \text{Hom}_{\mathcal{C}}(F(Y), X) \rightarrow \text{Hom}_{\mathcal{D}}(Y, G(X))$ is a natural isomorphism and $D: J \rightarrow \mathcal{C}$ is a diagram with $\varprojlim D = (L, \lambda)$.



Adjoint on limits

Thm. Right adjoints preserve limits; left adjoints preserve colimits.


Proof. Assume $\varphi: \text{Hom}_{\mathcal{C}}(F(Y), X) \rightarrow \text{Hom}_{\mathcal{D}}(Y, G(X))$ is a natural isomorphism and $D: J \rightarrow \mathcal{C}$ is a diagram with $\varprojlim D = (L, \lambda)$.



Adjoint functors on limits

Thm. Right adjoints preserve limits; left adjoints preserve colimits.


Proof. Assume $\varphi: \text{Hom}_{\mathcal{C}}(F(Y), X) \rightarrow \text{Hom}_{\mathcal{D}}(Y, G(X))$ is a natural isomorphism and $D: J \rightarrow \mathcal{C}$ is a diagram with $\varprojlim D = (L, \lambda)$.



Adjoint on limits

Thm. Right adjoints preserve limits; left adjoints preserve colimits.


Proof. Assume $\varphi: \text{Hom}_{\mathcal{C}}(F(Y), X) \rightarrow \text{Hom}_{\mathcal{D}}(Y, G(X))$ is a natural isomorphism and $D: J \rightarrow \mathcal{C}$ is a diagram with $\varprojlim D = (L, \lambda)$.



Adjoint on limits

Thm. Right adjoints preserve limits; left adjoints preserve colimits.


Proof. Assume $\varphi: \text{Hom}_{\mathcal{C}}(F(Y), X) \rightarrow \text{Hom}_{\mathcal{D}}(Y, G(X))$ is a natural isomorphism and $D: J \rightarrow \mathcal{C}$ is a diagram with $\varprojlim D = (L, \lambda)$.



Adjoint on limits

Thm. Right adjoints preserve limits; left adjoints preserve colimits.


Proof. Assume $\varphi: \text{Hom}_{\mathcal{C}}(F(Y), X) \rightarrow \text{Hom}_{\mathcal{D}}(Y, G(X))$ is a natural isomorphism and $D: J \rightarrow \mathcal{C}$ is a diagram with $\varprojlim D = (L, \lambda)$.



Adjoint on limits

Thm. Right adjoints preserve limits; left adjoints preserve colimits.


Proof. Assume $\varphi: \text{Hom}_{\mathcal{C}}(F(Y), X) \rightarrow \text{Hom}_{\mathcal{D}}(Y, G(X))$ is a natural isomorphism and $D: J \rightarrow \mathcal{C}$ is a diagram with $\varprojlim D = (L, \lambda)$.



Adjoint on limits

Thm. Right adjoints preserve limits; left adjoints preserve colimits.

Proof. Assume $\varphi: \text{Hom}_{\mathcal{C}}(F(Y), X) \rightarrow \text{Hom}_{\mathcal{D}}(Y, G(X))$ is a natural isomorphism and $D: J \rightarrow \mathcal{C}$ is a diagram with $\varprojlim D = (L, \lambda)$.



Proof continued

We choose maps from $F(c)$ to $D(J)$ to be those in $\varphi^{-1}(\gamma)$

Proof continued

We choose maps from $F(c)$ to $D(J)$ to be those in $\varphi^{-1}(\gamma)$

To show:

Proof continued

We choose maps from $F(c)$ to $D(J)$ to be those in $\varphi^{-1}(\gamma)$

To show:

(1) $(F(c), \varphi^{-1}(\gamma))$ is a cone to D .

Proof continued

We choose maps from $F(c)$ to $D(J)$ to be those in $\varphi^{-1}(\gamma)$

To show:

- (1) $(F(c), \varphi^{-1}(\gamma))$ is a cone to D .
- (2) $\varphi(\theta)$ is a cone morphism.

Proof continued

We choose maps from $F(c)$ to $D(J)$ to be those in $\varphi^{-1}(\gamma)$

To show:


- (1) $(F(c), \varphi^{-1}(\gamma))$ is a cone to D .
- (2) $\varphi(\theta)$ is a cone morphism.
- (3) $\varphi(\theta)$ is the unique cone morphism from (c, γ) to $(G(L), G(\lambda))$.

Proof continued

We choose maps from $F(c)$ to $D(J)$ to be those in $\varphi^{-1}(\gamma)$

To show:

- (1) $(F(c), \varphi^{-1}(\gamma))$ is a cone to D .
- (2) $\varphi(\theta)$ is a cone morphism.
- (3) $\varphi(\theta)$ is the unique cone morphism from (c, γ) to $(G(L), G(\lambda))$.




Proof continued

We choose maps from $F(c)$ to $D(J)$ to be those in $\varphi^{-1}(\gamma)$

To show:

- (1) $(F(c), \varphi^{-1}(\gamma))$ is a cone to D .
- (2) $\varphi(\theta)$ is a cone morphism.
- (3) $\varphi(\theta)$ is the unique cone morphism from (c, γ) to $(G(L), G(\lambda))$.




Proof continued

We choose maps from $F(c)$ to $D(J)$ to be those in $\varphi^{-1}(\gamma)$

To show:

- (1) $(F(c), \varphi^{-1}(\gamma))$ is a cone to D .
- (2) $\varphi(\theta)$ is a cone morphism.
- (3) $\varphi(\theta)$ is the unique cone morphism from (c, γ) to $(G(L), G(\lambda))$.




Proof continued

We choose maps from $F(c)$ to $D(J)$ to be those in $\varphi^{-1}(\gamma)$


To show:

- (1) $(F(c), \varphi^{-1}(\gamma))$ is a cone to D .
- (2) $\varphi(\theta)$ is a cone morphism.
- (3) $\varphi(\theta)$ is the unique cone morphism from (c, γ) to $(G(L), G(\lambda))$.




(1) $(F(c), \varphi^{-1}(\gamma))$ is a cone to D


(1) $(F(c), \varphi^{-1}(\gamma))$ is a cone to D



(1) $(F(c), \varphi^{-1}(\gamma))$ is a cone to D




(1) $(F(c), \varphi^{-1}(\gamma))$ is a cone to D



Must show: $h = G(f) \circ g$ implies $\varphi_{c,y}^{-1}(h) = f \circ \varphi_{c,x}^{-1}(g)$.


(1) $(F(c), \varphi^{-1}(\gamma))$ is a cone to D



Must show: $h = G(f) \circ g$ implies $\varphi_{c,y}^{-1}(h) = f \circ \varphi_{c,x}^{-1}(g)$.

Recall: $\varphi_{c,y}^{-1}(h) = \varepsilon_y \circ F(h)$ where $\varepsilon: FG \rightarrow I$ is the counit of the adjunction.

(1) $(F(c), \varphi^{-1}(\gamma))$ is a cone to D




Must show: $h = G(f) \circ g$ implies $\varphi_{c,y}^{-1}(h) = f \circ \varphi_{c,x}^{-1}(g)$.

Recall: $\varphi_{c,y}^{-1}(h) = \varepsilon_y \circ F(h)$ where $\varepsilon: FG \rightarrow I$ is the counit of the adjunction.


$$\begin{aligned}
 \varphi_{c,y}^{-1}(h) &= \varepsilon_y \circ F(h) \\
 &= \varepsilon_y \circ F(G(f) \circ g) \\
 &= \varepsilon_y \circ F(G(f)) \circ F(g) \\
 &= f \circ \varepsilon_x \circ F(g) \\
 &= f \circ \varphi_{c,x}^{-1}(g)
 \end{aligned}$$

(2) $\varphi(\theta)$ is a cone morphism


(2) $\varphi(\theta)$ is a cone morphism




(2) $\varphi(\theta)$ is a cone morphism




(2) $\varphi(\theta)$ is a cone morphism




(2) $\varphi(\theta)$ is a cone morphism




(2) $\varphi(\theta)$ is a cone morphism



(2) $\varphi(\theta)$ is a cone morphism




(2) $\varphi(\theta)$ is a cone morphism



Must show: $\ell \circ \theta = \varphi_{c,x}^{-1}(g)$ implies $G(\ell) \circ \varphi_{c,L}(\theta) = g$.

(2) $\varphi(\theta)$ is a cone morphism



Must show: $\ell \circ \theta = \varphi_{c,x}^{-1}(g)$ implies $G(\ell) \circ \varphi_{c,L}(\theta) = g$.

$$\begin{aligned}
 G(\ell) \circ \varphi_{c,L}(\theta) &= G(\ell) \circ G(\theta) \circ \eta_c \\
 &= G(\ell \circ \theta) \circ \eta_c \\
 &= G(\varphi_{c,x}^{-1}(g)) \circ \eta_c \\
 &= \varphi_{c,x}(\varphi_{c,x}^{-1}(g)) \\
 &= g
 \end{aligned}$$

(3) $\varphi(\theta)$ is unique

(3) $\varphi(\theta)$ is unique

The previous argument showed that if $\theta: F(c) \rightarrow L$ is a cone map, then $\varphi(\theta): c \rightarrow G(L)$ is a cone map.

(3) $\varphi(\theta)$ is unique


The previous argument showed that if $\theta: F(c) \rightarrow L$ is a cone map, then $\varphi(\theta): c \rightarrow G(L)$ is a cone map.

The reverse is true, too.

(3) $\varphi(\theta)$ is unique

The previous argument showed that if $\theta: F(c) \rightarrow L$ is a cone map, then $\varphi(\theta): c \rightarrow G(L)$ is a cone map.


The reverse is true, too.



(3) $\varphi(\theta)$ is unique

The previous argument showed that if $\theta: F(c) \rightarrow L$ is a cone map, then $\varphi(\theta): c \rightarrow G(L)$ is a cone map.

The reverse is true, too.




$$\begin{aligned}\varphi_{c,x}(\ell \circ \varphi_{c,L}^{-1}(\alpha)) &= G(\ell \circ \varphi_{c,L}^{-1}(\alpha)) \circ \eta_x = G(\ell) \circ G(\varphi_{c,L}^{-1}(\alpha)) \circ \eta_x \\ &= G(\ell) \circ \varphi_{c,L}(\varphi_{c,L}^{-1}(\alpha)) = G(\ell) \circ \alpha = g.\end{aligned}$$

(3) $\varphi(\theta)$ is unique

The previous argument showed that if $\theta: F(c) \rightarrow L$ is a cone map, then $\varphi(\theta): c \rightarrow G(L)$ is a cone map.

The reverse is true, too.



$$\begin{aligned}\varphi_{c,x}(\ell \circ \varphi_{c,L}^{-1}(\alpha)) &= G(\ell \circ \varphi_{c,L}^{-1}(\alpha)) \circ \eta_x = G(\ell) \circ G(\varphi_{c,L}^{-1}(\alpha)) \circ \eta_x \\ &= G(\ell) \circ \varphi_{c,L}(\varphi_{c,L}^{-1}(\alpha)) = G(\ell) \circ \alpha = g.\end{aligned}$$

Thus cone maps correspond under φ , and the uniqueness of our earlier θ implies the uniqueness of $\varphi(\theta)$.

Consequences and remarks

Consequences and remarks

(1) Direct sum of modules is a colimit, so it is preserved by left adjoints like $_ \otimes M$.

Consequences and remarks

- (1) Direct sum of modules is a colimit, so it is preserved by left adjoints like $_ \otimes M$.
- (2) Directed union is a colimit, so it is preserved by $_ \otimes M$.

Consequences and remarks

- (1) Direct sum of modules is a colimit, so it is preserved by left adjoints like $_ \otimes M$.
- (2) Directed union is a colimit, so it is preserved by $_ \otimes M$.
- (3) The theorem can be used to prove some functors don't have adjoints. E.g., the functor $A \mapsto A \oplus \mathbb{Z}$ has neither a right or left adjoint.

Consequences and remarks

- (1) Direct sum of modules is a colimit, so it is preserved by left adjoints like $_ \otimes M$.
- (2) Directed union is a colimit, so it is preserved by $_ \otimes M$.
- (3) The theorem can be used to prove some functors don't have adjoints. E.g., the functor $A \mapsto A \oplus \mathbb{Z}$ has neither a right or left adjoint.
- (4) Limits over a fixed category J (when they exist) yield a functor $\varprojlim: \mathcal{A}^J \rightarrow \mathcal{A}$. Colimits over J yield a functor $\text{colim} = \varinjlim: \mathcal{A}^J \rightarrow \mathcal{A}$. These are right and left adjoints to the diagonal functor $\Delta: \mathcal{A} \rightarrow \mathcal{A}^J$. E.g.,

$$\text{Hom}_{\mathcal{A}^J}(\Delta c, D) \cong \text{Hom}_{\mathcal{A}}(c, \varprojlim D).$$

(Lefthand side = cones to D ; righthand side = maps to the limit.)

Consequences and remarks

- (1) Direct sum of modules is a colimit, so it is preserved by left adjoints like $_ \otimes M$.
- (2) Directed union is a colimit, so it is preserved by $_ \otimes M$.
- (3) The theorem can be used to prove some functors don't have adjoints. E.g., the functor $A \mapsto A \oplus \mathbb{Z}$ has neither a right or left adjoint.
- (4) Limits over a fixed category J (when they exist) yield a functor $\varprojlim: \mathcal{A}^J \rightarrow \mathcal{A}$. Colimits over J yield a functor $\varinjlim = \varprojlim: \mathcal{A}^J \rightarrow \mathcal{A}$. These are right and left adjoints to the diagonal functor $\Delta: \mathcal{A} \rightarrow \mathcal{A}^J$. E.g.,

$$\mathrm{Hom}_{\mathcal{A}^J}(\Delta c, D) \cong \mathrm{Hom}_{\mathcal{A}}(c, \varprojlim D).$$

(Lefthand side = cones to D ; righthand side = maps to the limit.)

- (5) In particular, the limit functor is left exact and the colimit functor is right exact. The situation where such functors are fully exact is important, since homology commutes with exact functors.