

Map Coloring

Discrete Math

Math 2001

Spring 2011

The Four Color Conjecture

The Four Color Conjecture

Conjecture (Guthrie, 1852)


Any map can be colored with four colors so that no two adjacent regions have the same color.

The Four Color Conjecture

Conjecture (Guthrie, 1852)

Any map can be colored with four colors so that no two adjacent regions have the same color.

Graph-theoretical formulation




The Four Color Conjecture

Conjecture (Guthrie, 1852)

Any map can be colored with four colors so that no two adjacent regions have the same color.

Graph-theoretical formulation





The Four Color Conjecture

Conjecture (Guthrie, 1852)

Any map can be colored with four colors so that no two adjacent regions have the same color.

Graph-theoretical formulation





The Four Color Conjecture

Conjecture (Guthrie, 1852)

Any map can be colored with four colors so that no two adjacent regions have the same color.

Graph-theoretical formulation



If a graph represents a planar map, then it is possible to color its vertices so that no two adjacent vertices have the same color.

History

History

- Proofs were announced for the four color conjecture by Kempe (1879) and Tait (1880).

History

- Proofs were announced for the four color conjecture by Kempe (1879) and Tait (1880).
- Kempe's proof was shown to be false by Heawood (1890).

History

- Proofs were announced for the four color conjecture by Kempe (1879) and Tait (1880).
- Kempe's proof was shown to be false by Heawood (1890). Tait's proof was shown to be false by Petersen (1891).

History

- Proofs were announced for the four color conjecture by Kempe (1879) and Tait (1880).
- Kempe's proof was shown to be false by Heawood (1890). Tait's proof was shown to be false by Petersen (1891).
- Heawood used ideas from Kempe's erroneous proof to show that any planar graph has a proper coloring using at most five colors.

History

- Proofs were announced for the four color conjecture by Kempe (1879) and Tait (1880).
- Kempe's proof was shown to be false by Heawood (1890). Tait's proof was shown to be false by Petersen (1891).
- Heawood used ideas from Kempe's erroneous proof to show that any planar graph has a proper coloring using at most five colors.
- The conjecture was finally proved by Appel and Haken (1977) using substantial computer assistance.

History

- Proofs were announced for the four color conjecture by Kempe (1879) and Tait (1880).
- Kempe's proof was shown to be false by Heawood (1890). Tait's proof was shown to be false by Petersen (1891).
- Heawood used ideas from Kempe's erroneous proof to show that any planar graph has a proper coloring using at most five colors.
- The conjecture was finally proved by Appel and Haken (1977) using substantial computer assistance.

Map coloring research may be split into two (mostly unrelated) areas:

History

- Proofs were announced for the four color conjecture by Kempe (1879) and Tait (1880).
- Kempe's proof was shown to be false by Heawood (1890). Tait's proof was shown to be false by Petersen (1891).
- Heawood used ideas from Kempe's erroneous proof to show that any planar graph has a proper coloring using at most five colors.
- The conjecture was finally proved by Appel and Haken (1977) using substantial computer assistance.

Map coloring research may be split into two (mostly unrelated) areas:

1. Graph coloring.

History

- Proofs were announced for the four color conjecture by Kempe (1879) and Tait (1880).
- Kempe's proof was shown to be false by Heawood (1890). Tait's proof was shown to be false by Petersen (1891).
- Heawood used ideas from Kempe's erroneous proof to show that any planar graph has a proper coloring using at most five colors.
- The conjecture was finally proved by Appel and Haken (1977) using substantial computer assistance.

Map coloring research may be split into two (mostly unrelated) areas:

1. Graph coloring.
2. Drawing graphs on surfaces.

Definition

If $G = (V, E)$ is a graph and C is a color set, then a ***proper coloring*** of G is a function $f: V \rightarrow C$ such that $\forall u, v (E(u, v) \rightarrow (f(u) \neq f(v)))$.

Graph Coloring

Definition

If $G = (V, E)$ is a graph and C is a color set, then a **proper coloring** of G is a function $f: V \rightarrow C$ such that $\forall u, v (E(u, v) \rightarrow (f(u) \neq f(v)))$.

Definition

The **chromatic number** of G is the least number of colors needed in a proper coloring of G . It is denoted $\chi(G)$.

Graph Coloring

Definition

If $G = (V, E)$ is a graph and C is a color set, then a **proper coloring** of G is a function $f: V \rightarrow C$ such that $\forall u, v (E(u, v) \rightarrow (f(u) \neq f(v)))$.

Definition

The **chromatic number** of G is the least number of colors needed in a proper coloring of G . It is denoted $\chi(G)$.

Observations

Graph Coloring

Definition

If $G = (V, E)$ is a graph and C is a color set, then a **proper coloring** of G is a function $f: V \rightarrow C$ such that $\forall u, v (E(u, v) \rightarrow (f(u) \neq f(v)))$.

Definition

The **chromatic number** of G is the least number of colors needed in a proper coloring of G . It is denoted $\chi(G)$.

Observations

① $\chi(K_n) = n$.

Definition

If $G = (V, E)$ is a graph and C is a color set, then a **proper coloring** of G is a function $f: V \rightarrow C$ such that $\forall u, v (E(u, v) \rightarrow (f(u) \neq f(v)))$.

Definition

The **chromatic number** of G is the least number of colors needed in a proper coloring of G . It is denoted $\chi(G)$.

Observations

- ① $\chi(K_n) = n$.
- ② In fact, $\omega(G) \leq \chi(G)$.

Definition

If $G = (V, E)$ is a graph and C is a color set, then a **proper coloring** of G is a function $f: V \rightarrow C$ such that $\forall u, v (E(u, v) \rightarrow (f(u) \neq f(v)))$.

Definition

The **chromatic number** of G is the least number of colors needed in a proper coloring of G . It is denoted $\chi(G)$.

Observations

- ① $\chi(K_n) = n$.
- ② In fact, $\omega(G) \leq \chi(G)$.

Graph Coloring

Definition

If $G = (V, E)$ is a graph and C is a color set, then a **proper coloring** of G is a function $f: V \rightarrow C$ such that $\forall u, v (E(u, v) \rightarrow (f(u) \neq f(v)))$.

Definition

The **chromatic number** of G is the least number of colors needed in a proper coloring of G . It is denoted $\chi(G)$.

Observations

- 1 $\chi(K_n) = n$.
- 2 In fact, $\omega(G) \leq \chi(G)$. (Use $H \leq G \implies \chi(H) \leq \chi(G)$.)

Graph Coloring

Definition

If $G = (V, E)$ is a graph and C is a color set, then a **proper coloring** of G is a function $f: V \rightarrow C$ such that $\forall u, v (E(u, v) \rightarrow (f(u) \neq f(v)))$.

Definition

The **chromatic number** of G is the least number of colors needed in a proper coloring of G . It is denoted $\chi(G)$.

Observations

- 1 $\chi(K_n) = n$.
- 2 In fact, $\omega(G) \leq \chi(G)$. (Use $H \leq G \implies \chi(H) \leq \chi(G)$.)
- 3 On the other hand, $\chi(G) \leq \Delta(G) + 1$, where $\Delta(G)$ denotes the largest degree of a vertex of G .

Graph coloring problems in disguise

Graph coloring problems in disguise

- How many tanks are required to safely house n species of fish if some species prey on others?

Graph coloring problems in disguise

- How many tanks are required to safely house n species of fish if some species prey on others?
- How many storerooms are required to store n canisters of chemicals if some pairs of chemicals are explosive in combination?

Graph coloring problems in disguise

- How many tanks are required to safely house n species of fish if some species prey on others?
- How many storerooms are required to store n canisters of chemicals if some pairs of chemicals are explosive in combination?
- How many frequencies are needed for n radio stations if two stations cannot use the same frequency when they are within 150 miles of each other?

Graph coloring problems in disguise

- How many tanks are required to safely house n species of fish if some species prey on others?
- How many storerooms are required to store n canisters of chemicals if some pairs of chemicals are explosive in combination?
- How many frequencies are needed for n radio stations if two stations cannot use the same frequency when they are within 150 miles of each other?
- How many final examination slots are needed for n courses if two courses that have a common student cannot be scheduled at the same time?

Graph coloring problems in disguise

- How many tanks are required to safely house n species of fish if some species prey on others?
- How many storerooms are required to store n canisters of chemicals if some pairs of chemicals are explosive in combination?
- How many frequencies are needed for n radio stations if two stations cannot use the same frequency when they are within 150 miles of each other?
- How many final examination slots are need for n courses if two courses that have a common student cannot be scheduled at the same time?
- How many tables are required to seat n guests if some pairs of guests hate each others' guts?

Graph coloring problems in disguise

- How many tanks are required to safely house n species of fish if some species prey on others?
- How many storerooms are required to store n canisters of chemicals if some pairs of chemicals are explosive in combination?
- How many frequencies are needed for n radio stations if two stations cannot use the same frequency when they are within 150 miles of each other?
- How many final examination slots are need for n courses if two courses that have a common student cannot be scheduled at the same time?
- How many tables are required to seat n guests if some pairs of guests hate each others' guts?
- Solve Sudoku.

Alternative terminology

Alternative terminology

G is 2-colorable $\longleftrightarrow G$ is *bipartite*

Alternative terminology

G is 2-colorable $\longleftrightarrow G$ is *bipartite*

G is 3-colorable $\longleftrightarrow G$ is *tripartite*

Alternative terminology

G is 2-colorable $\iff G$ is *bipartite*

G is 3-colorable $\iff G$ is *tripartite*

G is r -colorable $\iff G$ is *r -partite*

Alternative terminology

G is 2-colorable $\iff G$ is *bipartite*

G is 3-colorable $\iff G$ is *tripartite*

G is r -colorable $\iff G$ is *r -partite*

It is easy to determine quickly if a graph is bipartite (how?), but no one knows an easy way to decide which graphs are r -partite when $r > 2$.

Alternative terminology

G is 2-colorable $\iff G$ is *bipartite*

G is 3-colorable $\iff G$ is *tripartite*

G is r -colorable $\iff G$ is *r -partite*

It is easy to determine quickly if a graph is bipartite (how?), but no one knows an easy way to decide which graphs are r -partite when $r > 2$.

Definition

A *complete r -partite graph*, K_{n_1, \dots, n_r} , is a graph (V, E) where

Alternative terminology

G is 2-colorable $\longleftrightarrow G$ is *bipartite*

G is 3-colorable $\longleftrightarrow G$ is *tripartite*

G is r -colorable $\longleftrightarrow G$ is *r -partite*

It is easy to determine quickly if a graph is bipartite (how?), but no one knows an easy way to decide which graphs are r -partite when $r > 2$.

Definition

A *complete r -partite graph*, K_{n_1, \dots, n_r} , is a graph (V, E) where

- 1 V is a union $V_1 \cup V_2 \cup \dots \cup V_r$ of disjoint sets,

Alternative terminology

G is 2-colorable $\longleftrightarrow G$ is ***bipartite***

G is 3-colorable $\longleftrightarrow G$ is ***tripartite***

G is r -colorable $\longleftrightarrow G$ is ***r -partite***

It is easy to determine quickly if a graph is bipartite (how?), but no one knows an easy way to decide which graphs are r -partite when $r > 2$.

Definition

A ***complete r -partite graph***, K_{n_1, \dots, n_r} , is a graph (V, E) where

- 1 V is a union $V_1 \cup V_2 \cup \dots \cup V_r$ of disjoint sets,
- 2 every vertex of V_i is adjacent to every vertex of V_j if $i \neq j$, and

Alternative terminology

G is 2-colorable $\longleftrightarrow G$ is **bipartite**

G is 3-colorable $\longleftrightarrow G$ is **tripartite**

G is r -colorable $\longleftrightarrow G$ is **r -partite**

It is easy to determine quickly if a graph is bipartite (how?), but no one knows an easy way to decide which graphs are r -partite when $r > 2$.

Definition

A **complete r -partite graph**, K_{n_1, \dots, n_r} , is a graph (V, E) where

- 1 V is a union $V_1 \cup V_2 \cup \dots \cup V_r$ of disjoint sets,
- 2 every vertex of V_i is adjacent to every vertex of V_j if $i \neq j$, and
- 3 there are no other adjacencies.

Alternative terminology

G is 2-colorable $\longleftrightarrow G$ is *bipartite*

G is 3-colorable $\longleftrightarrow G$ is *tripartite*

G is r -colorable $\longleftrightarrow G$ is *r -partite*

It is easy to determine quickly if a graph is bipartite (how?), but no one knows an easy way to decide which graphs are r -partite when $r > 2$.

Definition

A **complete r -partite graph**, K_{n_1, \dots, n_r} , is a graph (V, E) where

- 1 V is a union $V_1 \cup V_2 \cup \dots \cup V_r$ of disjoint sets,
- 2 every vertex of V_i is adjacent to every vertex of V_j if $i \neq j$, and
- 3 there are no other adjacencies.

Alternative terminology

G is 2-colorable $\iff G$ is **bipartite**

G is 3-colorable $\iff G$ is **tripartite**

G is r -colorable $\iff G$ is **r -partite**

It is easy to determine quickly if a graph is bipartite (how?), but no one knows an easy way to decide which graphs are r -partite when $r > 2$.

Definition

A **complete r -partite graph**, K_{n_1, \dots, n_r} , is a graph (V, E) where

- 1 V is a union $V_1 \cup V_2 \cup \dots \cup V_r$ of disjoint sets,
- 2 every vertex of V_i is adjacent to every vertex of V_j if $i \neq j$, and
- 3 there are no other adjacencies.

This conflicts slightly with the notation K_n .

Drawing Graphs on Surfaces

Drawing Graphs on Surfaces

Every graph that represents a planar map is 4-colorable.

Drawing Graphs on Surfaces

Every graph that represents a planar map is 4-colorable.

But not every graph is 4-colorable (e.g. K_n for $n > 4$).

Drawing Graphs on Surfaces

Every graph that represents a planar map is 4-colorable.

But not every graph is 4-colorable (e.g. K_n for $n > 4$).

The conclusion is that not all graphs represent planar maps.

Drawing Graphs on Surfaces

Every graph that represents a planar map is 4-colorable.

But not every graph is 4-colorable (e.g. K_n for $n > 4$).

The conclusion is that not all graphs represent planar maps. Which do?

Drawing Graphs on Surfaces

Every graph that represents a planar map is 4-colorable.

But not every graph is 4-colorable (e.g. K_n for $n > 4$).

The conclusion is that not all graphs represent planar maps. Which do?

Definition

A graph is *planar* if it can be drawn in the plane without edges crossing.

Drawing Graphs on Surfaces

Every graph that represents a planar map is 4-colorable.

But not every graph is 4-colorable (e.g. K_n for $n > 4$).

The conclusion is that not all graphs represent planar maps. Which do?

Definition

A graph is *planar* if it can be drawn in the plane without edges crossing.

Drawing Graphs on Surfaces

Every graph that represents a planar map is 4-colorable.

But not every graph is 4-colorable (e.g. K_n for $n > 4$).

The conclusion is that not all graphs represent planar maps. Which do?

Definition

A graph is *planar* if it can be drawn in the plane without edges crossing.

These are exactly the graphs that can be drawn on the *sphere* without edges crossing.

Drawing Graphs on Surfaces

Every graph that represents a planar map is 4-colorable.

But not every graph is 4-colorable (e.g. K_n for $n > 4$).

The conclusion is that not all graphs represent planar maps. Which do?

Definition

A graph is *planar* if it can be drawn in the plane without edges crossing.

These are exactly the graphs that can be drawn on the *sphere* without edges crossing.

The connected planar graphs may be thought of as the graphs that represent dissections of the sphere into polygons.

Drawing Graphs on Surfaces

Every graph that represents a planar map is 4-colorable.

But not every graph is 4-colorable (e.g. K_n for $n > 4$).

The conclusion is that not all graphs represent planar maps. Which do?

Definition

A graph is ***planar*** if it can be drawn in the plane without edges crossing.





These are exactly the graphs that can be drawn on the *sphere* without edges crossing.

The connected planar graphs may be thought of as the graphs that represent dissections of the sphere into polygons.





This suggests the idea that one might be able to distinguish between surfaces (like the sphere versus the torus) by determining which graphs represent dissections of the surfaces into polygons.

Numerical Invariants of Planar Drawings

Numerical Invariants of Planar Drawings







Numerical Invariants of Planar Drawings






(v, e, r)

Numerical Invariants of Planar Drawings







$$(v, e, r) = (4, 6, 4),$$

Numerical Invariants of Planar Drawings







$$(v, e, r) = (4, 6, 4), (8, 12, 6),$$

Numerical Invariants of Planar Drawings







$$(v, e, r) = (4, 6, 4), (8, 12, 6), (6, 12, 8),$$

Numerical Invariants of Planar Drawings







$$(v, e, r) = (4, 6, 4), (8, 12, 6), (6, 12, 8), (20, 30, 12),$$

Numerical Invariants of Planar Drawings



$(v, e, r) = (4, 6, 4), (8, 12, 6), (6, 12, 8), (20, 30, 12), (12, 30, 20)$

Numerical Invariants of Planar Drawings







$$(v, e, r) = (4, 6, 4), (8, 12, 6), (6, 12, 8), (20, 30, 12), (12, 30, 20)$$

Euler's Formula

In any planar drawing of a connected graph, $v - e + r = 2$.

Numerical Invariants of Planar Drawings



$$(v, e, r) = (4, 6, 4), (8, 12, 6), (6, 12, 8), (20, 30, 12), (12, 30, 20)$$

Euler's Formula





In any planar drawing of a connected graph, $v - e + r = 2$.

Proof.

Induction on e .



Numerical Invariants of Planar Drawings



$$(v, e, r) = (4, 6, 4), (8, 12, 6), (6, 12, 8), (20, 30, 12), (12, 30, 20)$$

Euler's Formula





In any planar drawing of a connected graph, $v - e + r = 2$.

Proof.

Induction on e .



Numerical Invariants of Planar Drawings



$$(v, e, r) = (4, 6, 4), (8, 12, 6), (6, 12, 8), (20, 30, 12), (12, 30, 20)$$

Euler's Formula

In any planar drawing of a connected graph, $v - e + r = 2$.

Proof.

Induction on e . □

The proof actually shows that for *any* surface S , the number $v - e + r$ is the same for all graphs that represent dissections of S into polygons. This number may be different for different surfaces. It is called the ***Euler characteristic*** of S , $\chi(S)$.

Eliminating the “ r ” from Euler’s Formula.

Eliminating the “ r ” from Euler’s Formula.

Theorem

Let $G = (V, E)$ be a connected planar graph with $|V| = v$ and $|E| = e$.

Eliminating the “ r ” from Euler’s Formula.

Theorem

Let $G = (V, E)$ be a connected planar graph with $|V| = v$ and $|E| = e$.

① $e \leq 3v - 6$.

Eliminating the “ r ” from Euler’s Formula.

Theorem

Let $G = (V, E)$ be a connected planar graph with $|V| = v$ and $|E| = e$.

- ① $e \leq 3v - 6$.
- ② If G is bipartite, then $e \leq 2v - 4$.

Eliminating the “ r ” from Euler’s Formula.

Theorem

Let $G = (V, E)$ be a connected planar graph with $|V| = v$ and $|E| = e$.

- ① $e \leq 3v - 6$.
- ② If G is bipartite, then $e \leq 2v - 4$.

Corollary

K_5 and $K_{3,3}$ are not planar.

Eliminating the “ r ” from Euler’s Formula.

Theorem

Let $G = (V, E)$ be a connected planar graph with $|V| = v$ and $|E| = e$.

- ① $e \leq 3v - 6$.
- ② *If G is bipartite, then $e \leq 2v - 4$.*

Corollary

K_5 and $K_{3,3}$ are not planar.

Kuratowski’s Theorem

A graph G is planar iff it has no subdivision of K_5 or $K_{3,3}$ as a subgraph.