

## Some uncountable sets

# Cantor's Theorem

**Thm.**

# Cantor's Theorem

**Thm.** There is no surjective function  $f: A \rightarrow \mathcal{P}(A)$ .

# Cantor's Theorem

**Thm.** There is no surjective function  $f: A \rightarrow \mathcal{P}(A)$ .

In fact, if  $f: A \rightarrow \mathcal{P}(A)$  is any function, then the Russell-type set

# Cantor's Theorem

**Thm.** There is no surjective function  $f: A \rightarrow \mathcal{P}(A)$ .

In fact, if  $f: A \rightarrow \mathcal{P}(A)$  is any function, then the Russell-type set

$$C = \{x \in A \mid x \notin f(x)\}$$

# Cantor's Theorem

**Thm.** There is no surjective function  $f: A \rightarrow \mathcal{P}(A)$ .

In fact, if  $f: A \rightarrow \mathcal{P}(A)$  is any function, then the Russell-type set

$$C = \{x \in A \mid x \notin f(x)\}$$

is not in the image of  $f$ .

# Cantor's Theorem

**Thm.** There is no surjective function  $f: A \rightarrow \mathcal{P}(A)$ .

In fact, if  $f: A \rightarrow \mathcal{P}(A)$  is any function, then the Russell-type set

$$C = \{x \in A \mid x \notin f(x)\}$$

is not in the image of  $f$ .

*Proof.*

# Cantor's Theorem

**Thm.** There is no surjective function  $f: A \rightarrow \mathcal{P}(A)$ .

In fact, if  $f: A \rightarrow \mathcal{P}(A)$  is any function, then the Russell-type set

$$C = \{x \in A \mid x \notin f(x)\}$$

is not in the image of  $f$ .

*Proof.* Assume that  $x \in A$  and  $f(x) = C$ .



# Cantor's Theorem

**Thm.** There is no surjective function  $f: A \rightarrow \mathcal{P}(A)$ .

In fact, if  $f: A \rightarrow \mathcal{P}(A)$  is any function, then the Russell-type set

$$C = \{x \in A \mid x \notin f(x)\}$$

is not in the image of  $f$ .

*Proof.* Assume that  $x \in A$  and  $f(x) = C$ . Then  $x \in f(x)$  iff  $x \notin f(x)$ ,

# Cantor's Theorem

**Thm.** There is no surjective function  $f: A \rightarrow \mathcal{P}(A)$ .

In fact, if  $f: A \rightarrow \mathcal{P}(A)$  is any function, then the Russell-type set

$$C = \{x \in A \mid x \notin f(x)\}$$

is not in the image of  $f$ .

*Proof.* Assume that  $x \in A$  and  $f(x) = C$ . Then  $x \in f(x)$  iff  $x \notin f(x)$ , a contradiction.

# Cantor's Theorem

**Thm.** There is no surjective function  $f: A \rightarrow \mathcal{P}(A)$ .

In fact, if  $f: A \rightarrow \mathcal{P}(A)$  is any function, then the Russell-type set

$$C = \{x \in A \mid x \notin f(x)\}$$

is not in the image of  $f$ .

*Proof.* Assume that  $x \in A$  and  $f(x) = C$ . Then  $x \in f(x)$  iff  $x \notin f(x)$ , a contradiction.  $\square$

# Cantor's Theorem

**Thm.** There is no surjective function  $f: A \rightarrow \mathcal{P}(A)$ .

In fact, if  $f: A \rightarrow \mathcal{P}(A)$  is any function, then the Russell-type set

$$C = \{x \in A \mid x \notin f(x)\}$$

is not in the image of  $f$ .

*Proof.* Assume that  $x \in A$  and  $f(x) = C$ . Then  $x \in f(x)$  iff  $x \notin f(x)$ , a contradiction.  $\square$

**Cantor's Theorem.**

# Cantor's Theorem

**Thm.** There is no surjective function  $f: A \rightarrow \mathcal{P}(A)$ .

In fact, if  $f: A \rightarrow \mathcal{P}(A)$  is any function, then the Russell-type set

$$C = \{x \in A \mid x \notin f(x)\}$$

is not in the image of  $f$ .

*Proof.* Assume that  $x \in A$  and  $f(x) = C$ . Then  $x \in f(x)$  iff  $x \notin f(x)$ , a contradiction.  $\square$

**Cantor's Theorem.** For any set  $A$ ,  $|A| < |\mathcal{P}(A)|$ .

# Cantor's Theorem

**Thm.** There is no surjective function  $f: A \rightarrow \mathcal{P}(A)$ .

In fact, if  $f: A \rightarrow \mathcal{P}(A)$  is any function, then the Russell-type set

$$C = \{x \in A \mid x \notin f(x)\}$$

is not in the image of  $f$ .

*Proof.* Assume that  $x \in A$  and  $f(x) = C$ . Then  $x \in f(x)$  iff  $x \notin f(x)$ , a contradiction.  $\square$

**Cantor's Theorem.** For any set  $A$ ,  $|A| < |\mathcal{P}(A)|$ . Hence, if  $A$  is infinite, then  $\mathcal{P}(A)$  is uncountable.

# Cantor's Theorem

**Thm.** There is no surjective function  $f: A \rightarrow \mathcal{P}(A)$ .

In fact, if  $f: A \rightarrow \mathcal{P}(A)$  is any function, then the Russell-type set

$$C = \{x \in A \mid x \notin f(x)\}$$

is not in the image of  $f$ .

*Proof.* Assume that  $x \in A$  and  $f(x) = C$ . Then  $x \in f(x)$  iff  $x \notin f(x)$ , a contradiction.  $\square$

**Cantor's Theorem.** For any set  $A$ ,  $|A| < |\mathcal{P}(A)|$ . Hence, if  $A$  is infinite, then  $\mathcal{P}(A)$  is uncountable.

*Proof.*

# Cantor's Theorem

**Thm.** There is no surjective function  $f: A \rightarrow \mathcal{P}(A)$ .

In fact, if  $f: A \rightarrow \mathcal{P}(A)$  is any function, then the Russell-type set

$$C = \{x \in A \mid x \notin f(x)\}$$

is not in the image of  $f$ .

*Proof.* Assume that  $x \in A$  and  $f(x) = C$ . Then  $x \in f(x)$  iff  $x \notin f(x)$ , a contradiction.  $\square$

**Cantor's Theorem.** For any set  $A$ ,  $|A| < |\mathcal{P}(A)|$ . Hence, if  $A$  is infinite, then  $\mathcal{P}(A)$  is uncountable.

*Proof.* By HW3, Problem 1,  $F: A \rightarrow \mathcal{P}(A): a \mapsto \{a\}$  is injective,



# Cantor's Theorem

**Thm.** There is no surjective function  $f: A \rightarrow \mathcal{P}(A)$ .

In fact, if  $f: A \rightarrow \mathcal{P}(A)$  is any function, then the Russell-type set

$$C = \{x \in A \mid x \notin f(x)\}$$

is not in the image of  $f$ .

*Proof.* Assume that  $x \in A$  and  $f(x) = C$ . Then  $x \in f(x)$  iff  $x \notin f(x)$ , a contradiction.  $\square$

**Cantor's Theorem.** For any set  $A$ ,  $|A| < |\mathcal{P}(A)|$ . Hence, if  $A$  is infinite, then  $\mathcal{P}(A)$  is uncountable.

*Proof.* By HW3, Problem 1,  $F: A \rightarrow \mathcal{P}(A): a \mapsto \{a\}$  is injective, so  $|A| \leq |\mathcal{P}(A)|$ .

# Cantor's Theorem

**Thm.** There is no surjective function  $f: A \rightarrow \mathcal{P}(A)$ .

In fact, if  $f: A \rightarrow \mathcal{P}(A)$  is any function, then the Russell-type set

$$C = \{x \in A \mid x \notin f(x)\}$$

is not in the image of  $f$ .

*Proof.* Assume that  $x \in A$  and  $f(x) = C$ . Then  $x \in f(x)$  iff  $x \notin f(x)$ , a contradiction.  $\square$

**Cantor's Theorem.** For any set  $A$ ,  $|A| < |\mathcal{P}(A)|$ . Hence, if  $A$  is infinite, then  $\mathcal{P}(A)$  is uncountable.

*Proof.* By HW3, Problem 1,  $F: A \rightarrow \mathcal{P}(A): a \mapsto \{a\}$  is injective, so  $|A| \leq |\mathcal{P}(A)|$ . The theorem implies that  $|A| \neq |\mathcal{P}(A)|$ , so  $|A| < |\mathcal{P}(A)|$ .

# Cantor's Theorem

**Thm.** There is no surjective function  $f: A \rightarrow \mathcal{P}(A)$ .

In fact, if  $f: A \rightarrow \mathcal{P}(A)$  is any function, then the Russell-type set

$$C = \{x \in A \mid x \notin f(x)\}$$

is not in the image of  $f$ .

*Proof.* Assume that  $x \in A$  and  $f(x) = C$ . Then  $x \in f(x)$  iff  $x \notin f(x)$ , a contradiction.  $\square$

**Cantor's Theorem.** For any set  $A$ ,  $|A| < |\mathcal{P}(A)|$ . Hence, if  $A$  is infinite, then  $\mathcal{P}(A)$  is uncountable.

*Proof.* By HW3, Problem 1,  $F: A \rightarrow \mathcal{P}(A): a \mapsto \{a\}$  is injective, so  $|A| \leq |\mathcal{P}(A)|$ . The theorem implies that  $|A| \neq |\mathcal{P}(A)|$ , so  $|A| < |\mathcal{P}(A)|$ .  $\square$

# $\mathbb{R}$ is uncountable

**Thm.**

# $\mathbb{R}$ is uncountable

**Thm.**  $|\mathbb{R}| = |(0, 1)| = |\mathcal{P}(\mathbb{N})|$ .

# $\mathbb{R}$ is uncountable

**Thm.**  $|\mathbb{R}| = |(0, 1)| = |\mathcal{P}(\mathbb{N})|$ .

Define the **characteristic function** of a subset  $S \subseteq A$  to be the function

# $\mathbb{R}$ is uncountable

**Thm.**  $|\mathbb{R}| = |(0, 1)| = |\mathcal{P}(\mathbb{N})|$ .

Define the **characteristic function** of a subset  $S \subseteq A$  to be the function

$$\underbrace{\chi_S}_{\text{chi sub } S} : A \rightarrow \{0, 1\} : a \mapsto \begin{cases} 1 & \text{if } a \in S \\ 0 & \text{else.} \end{cases}$$



**Thm.**  $|\mathbb{R}| = |(0, 1)| = |\mathcal{P}(\mathbb{N})|$ .

Define the **characteristic function** of a subset  $S \subseteq A$  to be the function

$$\underbrace{\chi_S}_{\text{chi sub } S} : A \rightarrow \{0, 1\} : a \mapsto \begin{cases} 1 & \text{if } a \in S \\ 0 & \text{else.} \end{cases}$$

That is,  $\chi_S$  is the function from  $A$  to  $\{0, 1\}$  whose **support** is  $S$ .

# $\mathbb{R}$ is uncountable

**Thm.**  $|\mathbb{R}| = |(0, 1)| = |\mathcal{P}(\mathbb{N})|$ .

Define the **characteristic function** of a subset  $S \subseteq A$  to be the function

$$\underbrace{\chi_S}_{\text{chi sub } S} : A \rightarrow \{0, 1\} : a \mapsto \begin{cases} 1 & \text{if } a \in S \\ 0 & \text{else.} \end{cases}$$

That is,  $\chi_S$  is the function from  $A$  to  $\{0, 1\}$  whose **support** is  $S$ .

Idea of proof: Try to build bijections (or almost-bijections) between subsets of  $\mathbb{N}$ , characteristic functions of subsets of  $\mathbb{N}$ , and base 2 expansions of real numbers from  $(0, 1)$ .

# Additional facts about infinite cardinality

# Additional facts about infinite cardinality

**Thm.** (ZFC)

## Additional facts about infinite cardinality

**Thm.** (ZFC) Let  $A$  and  $B$  be nonempty sets with at least one of them infinite.

$$|A \cup B| = |A \times B| = \max(|A|, |B|).$$