

## Solutions to HW 10.

1. This problem involves a deck of 52 distinct playing cards.
  - (a) In how many ways can a 13-card bridge hand be dealt from the deck?

We want an ordered sequence of 13 cards chosen from a 52-element set. There are  $13! \cdot \binom{52}{13} = (52)_{13} = \frac{52!}{39!}$  of these.

- (b) How many different 13-card bridge hands are there?

We want a 13-card subset of a 52-element set. There are  $\binom{52}{13} = \frac{52!}{13!39!}$  of these.

2. (a) How many paths are there from the point  $(0,0)$  of  $\mathbb{R}^2$  to the point  $(10,15)$  of  $\mathbb{R}^2$  if each path consists of a sequence of steps of length 1 moving in the direction of the positive  $x$ -axis or the positive  $y$ -axis?

We can describe a path by a list of instructions of the form “ $(x, x, y, x, y, y, \dots, x, y, y)$ ”, which is a string of 10  $x$ ’s and 15  $y$ ’s in some order. If the first two instructions are  $x$ , then we take our first two steps in the  $x$  direction; if our next instruction is  $y$ , we take our next step in the  $y$  direction. ETC.

The number of paths is equal to the number of lists of instructions, which is equal to the number of strings of length  $10 + 15 = 25$  consisting of  $x$ ’s and  $y$ ’s, which contain 10  $x$ ’s and 15  $y$ ’s. This number is  $\binom{25}{10} = \frac{25!}{10!15!}$ . (You have 25 instructions, and you “choose” 10 instructions to be  $x$ ’s and leave the remaining 15 to be  $y$ ’s.)

- (b) How many paths are there from the point  $(0,0,0)$  of  $\mathbb{R}^3$  to the point  $(10,15,20)$  of  $\mathbb{R}^3$  if each path consists of a sequence of steps of length 1 moving in the direction of the positive  $x$ -axis, the positive  $y$ -axis or the positive  $z$ -axis?

Using the same reasoning, we want to count strings of length  $10 + 15 + 20 = 45$  which have 10  $x$ ’s, 15  $y$ ’s, and 20  $z$ ’s. The number is  $\binom{45}{10,15,20} = \frac{45!}{10!15!20!}$ .

3. Let  $MC(n, k)$  be the number “ $n$ -multichoose- $k$ ”. Use a combinatorial argument to show that  $MC(n, 0) + MC(n, 1) + \dots + MC(n, k) = MC(n + 1, k)$ .

**Solution 1.** (A combinatorial argument.) Let  $D$  be the set of all distributions of  $k$  identical balls to  $n + 1$  distinct boxes with repetition allowed.  $|D| = MC(n + 1, k) = \binom{n+1+k-1}{k} = \binom{n+k}{k}$ .

Partition  $D$  into sets  $D_0, D_1, \dots, D_k$  where  $D_i \subseteq D$  is the number of distributions where  $i$  balls are distributed to the first  $n$  boxes, while the remaining  $k - i$  balls are distributed to the last box. Since we can distribute  $i$  identical balls to the first  $n$  boxes in  $MC(n, i)$  ways, and we have no choice but to put the remaining  $k - i$  balls into the last box, we have  $|D_i| = MC(n, i)$ . Since we have a partition,

$$MC(n + 1, k) = |D| = |D_0| + |D_1| + \dots + |D_k| = MC(n, 0) + MC(n, 1) + \dots + MC(n, k).$$

**Solution 2.** (Not a combinatorial argument, but OK.)

$$\begin{aligned}
MC(n, 0) + MC(n, 1) + \cdots + MC(n, k) &= \binom{n-1}{0} + \binom{n}{1} + \binom{n+1}{2} + \cdots + \binom{n+k-1}{k} \\
&= [\binom{n}{0} + \binom{n}{1}] + \binom{n+1}{2} + \cdots + \binom{n+k-1}{k} \\
&= [\binom{n+1}{1}] + \binom{n+1}{2} + \cdots + \binom{n+k-1}{k} \\
&= [\binom{n+1}{1} + \binom{n+1}{2}] + \cdots + \binom{n+k-1}{k} \\
&\quad \vdots \\
&= \binom{n+k-1}{k-1} + \binom{n+k-1}{k} \\
&= \binom{n+k}{k} = MC(n+1, k).
\end{aligned}$$