


Truth versus Provability



Semantic versus Syntactic Consequence

Semantic versus Syntactic Consequence

Let Σ be a set of sentences

Semantic versus Syntactic Consequence

Let Σ be a set of sentences (like the axioms for set theory, or like the empty set of sentences).

Semantic versus Syntactic Consequence

Let Σ be a set of sentences (like the axioms for set theory, or like the empty set of sentences).

Let P be a single sentence.

Semantic versus Syntactic Consequence

Let Σ be a set of sentences (like the axioms for set theory, or like the empty set of sentences).

Let P be a single sentence. Is P a consequence of Σ ?

Semantic versus Syntactic Consequence

Let Σ be a set of sentences (like the axioms for set theory, or like the empty set of sentences).

Let P be a single sentence. Is P a consequence of Σ ? (Does P “follow” from Σ ?)

Semantic versus Syntactic Consequence

Let Σ be a set of sentences (like the axioms for set theory, or like the empty set of sentences).

Let P be a single sentence. Is P a consequence of Σ ? (Does P “follow” from Σ ?)

Definition.

Semantic versus Syntactic Consequence

Let Σ be a set of sentences (like the axioms for set theory, or like the empty set of sentences).

Let P be a single sentence. Is P a consequence of Σ ? (Does P “follow” from Σ ?)

Definition. P is a **semantic consequence** of Σ if every structure satisfying Σ must satisfy P .

Semantic versus Syntactic Consequence

Let Σ be a set of sentences (like the axioms for set theory, or like the empty set of sentences).

Let P be a single sentence. Is P a consequence of Σ ? (Does P “follow” from Σ ?)

Definition. P is a **semantic consequence** of Σ if every structure satisfying Σ must satisfy P . (Write $\Sigma \models P$.)

Semantic versus Syntactic Consequence

Let Σ be a set of sentences (like the axioms for set theory, or like the empty set of sentences).

Let P be a single sentence. Is P a consequence of Σ ? (Does P “follow” from Σ ?)

Definition. P is a **semantic consequence** of Σ if every structure satisfying Σ must satisfy P . (Write $\Sigma \models P$.)

P is a **syntactic consequence** of Σ if there is a “proof of P from Σ ”

Semantic versus Syntactic Consequence

Let Σ be a set of sentences (like the axioms for set theory, or like the empty set of sentences).

Let P be a single sentence. Is P a consequence of Σ ? (Does P “follow” from Σ ?)

Definition. P is a **semantic consequence** of Σ if every structure satisfying Σ must satisfy P . (Write $\Sigma \models P$.)

P is a **syntactic consequence** of Σ if there is a “proof of P from Σ ” (that is, a proof which uses the statements in Σ as hypotheses, or as additional axioms).

Semantic versus Syntactic Consequence

Let Σ be a set of sentences (like the axioms for set theory, or like the empty set of sentences).

Let P be a single sentence. Is P a consequence of Σ ? (Does P “follow” from Σ ?)

Definition. P is a **semantic consequence** of Σ if every structure satisfying Σ must satisfy P . (Write $\Sigma \models P$.)

P is a **syntactic consequence** of Σ if there is a “proof of P from Σ ” (that is, a proof which uses the statements in Σ as hypotheses, or as additional axioms). (Write $\Sigma \vdash P$.)

Semantic versus Syntactic Consequence

Let Σ be a set of sentences (like the axioms for set theory, or like the empty set of sentences).

Let P be a single sentence. Is P a consequence of Σ ? (Does P “follow” from Σ ?)

Definition. P is a **semantic consequence** of Σ if every structure satisfying Σ must satisfy P . (Write $\Sigma \models P$.)

P is a **syntactic consequence** of Σ if there is a “proof of P from Σ ” (that is, a proof which uses the statements in Σ as hypotheses, or as additional axioms). (Write $\Sigma \vdash P$.)

When $\Sigma = \emptyset$, we write “ $\models P$ ” for “ P is true” and “ $\vdash P$ ” for “ P is provable”.

Semantic versus Syntactic Consequence

Let Σ be a set of sentences (like the axioms for set theory, or like the empty set of sentences).

Let P be a single sentence. Is P a consequence of Σ ? (Does P “follow” from Σ ?)

Definition. P is a **semantic consequence** of Σ if every structure satisfying Σ must satisfy P . (Write $\Sigma \models P$.)

P is a **syntactic consequence** of Σ if there is a “proof of P from Σ ” (that is, a proof which uses the statements in Σ as hypotheses, or as additional axioms). (Write $\Sigma \vdash P$.)

When $\Sigma = \emptyset$, we write “ $\models P$ ” for “ P is true” and “ $\vdash P$ ” for “ P is provable”.

We have already discussed how to check whether a statement P is true in a structure

Semantic versus Syntactic Consequence

Let Σ be a set of sentences (like the axioms for set theory, or like the empty set of sentences).

Let P be a single sentence. Is P a consequence of Σ ? (Does P “follow” from Σ ?)

Definition. P is a **semantic consequence** of Σ if every structure satisfying Σ must satisfy P . (Write $\Sigma \models P$.)

P is a **syntactic consequence** of Σ if there is a “proof of P from Σ ” (that is, a proof which uses the statements in Σ as hypotheses, or as additional axioms). (Write $\Sigma \vdash P$.)

When $\Sigma = \emptyset$, we write “ $\models P$ ” for “ P is true” and “ $\vdash P$ ” for “ P is provable”.

We have already discussed how to check whether a statement P is true in a structure (check the tables of the structural elements! play quantifier games!).

Semantic versus Syntactic Consequence

Let Σ be a set of sentences (like the axioms for set theory, or like the empty set of sentences).

Let P be a single sentence. Is P a consequence of Σ ? (Does P “follow” from Σ ?)

Definition. P is a **semantic consequence** of Σ if every structure satisfying Σ must satisfy P . (Write $\Sigma \models P$.)

P is a **syntactic consequence** of Σ if there is a “proof of P from Σ ” (that is, a proof which uses the statements in Σ as hypotheses, or as additional axioms). (Write $\Sigma \vdash P$.)

When $\Sigma = \emptyset$, we write “ $\models P$ ” for “ P is true” and “ $\vdash P$ ” for “ P is provable”.

We have already discussed how to check whether a statement P is true in a structure (check the tables of the structural elements! play quantifier games!). Today we will discuss provability.

A **proof** of a **theorem** T is finite sequence of statements

$$S_1, S_2, \dots, S_n = T$$

A **proof** of a **theorem** T is finite sequence of statements

$$S_1, S_2, \dots, S_n = T$$

which ends at T and which has the property that each statement in the list follows from earlier statements in the list by some (accepted) law of deduction.

A **proof** of a **theorem** T is finite sequence of statements

$$S_1, S_2, \dots, S_n = T$$

which ends at T and which has the property that each statement in the list follows from earlier statements in the list by some (accepted) law of deduction.

First question.

A **proof** of a **theorem** T is finite sequence of statements

$$S_1, S_2, \dots, S_n = T$$

which ends at T and which has the property that each statement in the list follows from earlier statements in the list by some (accepted) law of deduction.

First question. How is this possible?

A **proof** of a **theorem** T is finite sequence of statements

$$S_1, S_2, \dots, S_n = T$$

which ends at T and which has the property that each statement in the list follows from earlier statements in the list by some (accepted) law of deduction.

First question. How is this possible? How do proofs get started?

A **proof** of a **theorem** T is finite sequence of statements

$$S_1, S_2, \dots, S_n = T$$

which ends at T and which has the property that each statement in the list follows from earlier statements in the list by some (accepted) law of deduction.

First question. How is this possible? How do proofs get started? (What does it mean for S_1 to follow from earlier statements if there are no earlier statements?)

A **proof** of a **theorem** T is finite sequence of statements

$$S_1, S_2, \dots, S_n = T$$

which ends at T and which has the property that each statement in the list follows from earlier statements in the list by some (accepted) law of deduction.

First question. How is this possible? How do proofs get started? (What does it mean for S_1 to follow from earlier statements if there are no earlier statements?)

Answer.

A **proof** of a **theorem** T is finite sequence of statements

$$S_1, S_2, \dots, S_n = T$$

which ends at T and which has the property that each statement in the list follows from earlier statements in the list by some (accepted) law of deduction.

First question. How is this possible? How do proofs get started? (What does it mean for S_1 to follow from earlier statements if there are no earlier statements?)

Answer. Axioms are statements that follow from the empty collection of preceding statements!

A **proof** of a **theorem** T is finite sequence of statements

$$S_1, S_2, \dots, S_n = T$$

which ends at T and which has the property that each statement in the list follows from earlier statements in the list by some (accepted) law of deduction.

First question. How is this possible? How do proofs get started? (What does it mean for S_1 to follow from earlier statements if there are no earlier statements?)

Answer. Axioms are statements that follow from the empty collection of preceding statements!

A **proof** of a **theorem** T is finite sequence of statements

$$S_1, S_2, \dots, S_n = T$$

which ends at T and which has the property that each statement in the list follows from earlier statements in the list by some (accepted) law of deduction.

First question. How is this possible? How do proofs get started? (What does it mean for S_1 to follow from earlier statements if there are no earlier statements?)

Answer. Axioms are statements that follow from the empty collection of preceding statements!

So, a “proof system” typically specifies its axioms and also the accepted rules of deduction.

What does a law of deduction look like?

What does a law of deduction look like?

Examples.

What does a law of deduction look like?

Examples.

- 1 (Modus Ponens)

What does a law of deduction look like?

Examples.

- 1 (Modus Ponens)

What does a law of deduction look like?

Examples.

- 1 (Modus Ponens) = “The mode that affirms”.

What does a law of deduction look like?

Examples.

- 1 (Modus Ponens) = “The mode that affirms”.

$$\frac{(P \rightarrow Q), P}{Q}$$

What does a law of deduction look like?

Examples.

- ① (Modus Ponens) = “The mode that affirms”.

$$\frac{(P \rightarrow Q), P}{Q}$$

So if your proof has the structure

What does a law of deduction look like?

Examples.

- ❶ (Modus Ponens) = “The mode that affirms”.

$$\frac{(P \rightarrow Q), P}{Q}$$

So if your proof has the structure $S_1, S_2, \dots, P, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step by appending $S_{k+1} = Q$ to this proof.

What does a law of deduction look like?

Examples.

- ① (Modus Ponens) = “The mode that affirms”.

$$\frac{(P \rightarrow Q), P}{Q}$$

So if your proof has the structure $S_1, S_2, \dots, P, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step by appending $S_{k+1} = Q$ to this proof.

- ② (Modus Tollens)

What does a law of deduction look like?

Examples.

- ① (Modus Ponens) = “The mode that affirms”.

$$\frac{(P \rightarrow Q), P}{Q}$$

So if your proof has the structure $S_1, S_2, \dots, P, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step by appending $S_{k+1} = Q$ to this proof.

- ② (Modus Tollens)

What does a law of deduction look like?

Examples.

- ① (Modus Ponens) = “The mode that affirms”.

$$\frac{(P \rightarrow Q), P}{Q}$$

So if your proof has the structure $S_1, S_2, \dots, P, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step by appending $S_{k+1} = Q$ to this proof.

- ② (Modus Tollens) = “The mode that denies”.

What does a law of deduction look like?

Examples.

- ❶ (Modus Ponens) = “The mode that affirms”.

$$\frac{(P \rightarrow Q), P}{Q}$$

So if your proof has the structure $S_1, S_2, \dots, P, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step by appending $S_{k+1} = Q$ to this proof.

- ❷ (Modus Tollens) = “The mode that denies”.

$$\frac{(P \rightarrow Q), \neg Q}{\neg P}$$

What does a law of deduction look like?

Examples.

- ❶ (Modus Ponens) = “The mode that affirms”.

$$\frac{(P \rightarrow Q), P}{Q}$$

So if your proof has the structure $S_1, S_2, \dots, P, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step by appending $S_{k+1} = Q$ to this proof.

- ❷ (Modus Tollens) = “The mode that denies”.

$$\frac{(P \rightarrow Q), \neg Q}{\neg P}$$

So if your proof has the structure

What does a law of deduction look like?

Examples.

- ① (Modus Ponens) = “The mode that affirms”.

$$\frac{(P \rightarrow Q), P}{Q}$$

So if your proof has the structure $S_1, S_2, \dots, P, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step by appending $S_{k+1} = Q$ to this proof.

- ② (Modus Tollens) = “The mode that denies”.

$$\frac{(P \rightarrow Q), \neg Q}{\neg P}$$

So if your proof has the structure $S_1, S_2, \dots, \neg Q, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step to get $S_1, S_2, \dots, \neg Q, \dots, P \rightarrow Q, \dots, S_k, \neg P$.

What does a law of deduction look like?

Examples.

- ❶ (Modus Ponens) = “The mode that affirms”.

$$\frac{(P \rightarrow Q), P}{Q}$$

So if your proof has the structure $S_1, S_2, \dots, P, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step by appending $S_{k+1} = Q$ to this proof.

- ❷ (Modus Tollens) = “The mode that denies”.

$$\frac{(P \rightarrow Q), \neg Q}{\neg P}$$

So if your proof has the structure $S_1, S_2, \dots, \neg Q, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step to get $S_1, S_2, \dots, \neg Q, \dots, P \rightarrow Q, \dots, S_k, \neg P$.

- ❸ (Axioms) $\frac{}{A}$.

What does a law of deduction look like?

Examples.

- ❶ (Modus Ponens) = “The mode that affirms”.

$$\frac{(P \rightarrow Q), P}{Q}$$

So if your proof has the structure $S_1, S_2, \dots, P, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step by appending $S_{k+1} = Q$ to this proof.

- ❷ (Modus Tollens) = “The mode that denies”.

$$\frac{(P \rightarrow Q), \neg Q}{\neg P}$$

So if your proof has the structure $S_1, S_2, \dots, \neg Q, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step to get $S_1, S_2, \dots, \neg Q, \dots, P \rightarrow Q, \dots, S_k, \neg P$.

- ❸ (Axioms) $\frac{}{A}$.

What does a law of deduction look like?

Examples.

- ❶ (Modus Ponens) = “The mode that affirms”.

$$\frac{(P \rightarrow Q), P}{Q}$$

So if your proof has the structure $S_1, S_2, \dots, P, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step by appending $S_{k+1} = Q$ to this proof.

- ❷ (Modus Tollens) = “The mode that denies”.

$$\frac{(P \rightarrow Q), \neg Q}{\neg P}$$

So if your proof has the structure $S_1, S_2, \dots, \neg Q, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step to get $S_1, S_2, \dots, \neg Q, \dots, P \rightarrow Q, \dots, S_k, \neg P$.

- ❸ (Axioms) $\frac{}{A}$.

- ❹ (Hypothetical syllogism)

What does a law of deduction look like?

Examples.

- ❶ (Modus Ponens) = “The mode that affirms”.

$$\frac{(P \rightarrow Q), P}{Q}$$

So if your proof has the structure $S_1, S_2, \dots, P, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step by appending $S_{k+1} = Q$ to this proof.

- ❷ (Modus Tollens) = “The mode that denies”.

$$\frac{(P \rightarrow Q), \neg Q}{\neg P}$$

So if your proof has the structure $S_1, S_2, \dots, \neg Q, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step to get $S_1, S_2, \dots, \neg Q, \dots, P \rightarrow Q, \dots, S_k, \neg P$.

- ❸ (Axioms) $\frac{}{A}$.

- ❹ (Hypothetical syllogism)

What does a law of deduction look like?

Examples.

- ① (Modus Ponens) = “The mode that affirms”.

$$\frac{(P \rightarrow Q), P}{Q}$$

So if your proof has the structure $S_1, S_2, \dots, P, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step by appending $S_{k+1} = Q$ to this proof.

- ② (Modus Tollens) = “The mode that denies”.

$$\frac{(P \rightarrow Q), \neg Q}{\neg P}$$

So if your proof has the structure $S_1, S_2, \dots, \neg Q, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step to get $S_1, S_2, \dots, \neg Q, \dots, P \rightarrow Q, \dots, S_k, \neg P$.

- ③ (Axioms) $\frac{}{A}$.

- ④ (Hypothetical syllogism) $\frac{(P \rightarrow Q), (Q \rightarrow R)}{(P \rightarrow R)}$

What does a law of deduction look like?

Examples.

- ❶ (Modus Ponens) = “The mode that affirms”.

$$\frac{(P \rightarrow Q), P}{Q}$$

So if your proof has the structure $S_1, S_2, \dots, P, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step by appending $S_{k+1} = Q$ to this proof.

- ❷ (Modus Tollens) = “The mode that denies”.

$$\frac{(P \rightarrow Q), \neg Q}{\neg P}$$

So if your proof has the structure $S_1, S_2, \dots, \neg Q, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step to get $S_1, S_2, \dots, \neg Q, \dots, P \rightarrow Q, \dots, S_k, \neg P$.

- ❸ (Axioms) $\frac{}{A}$.

- ❹ (Hypothetical syllogism) $\frac{(P \rightarrow Q), (Q \rightarrow R)}{(P \rightarrow R)}$

- ❺ (Disjunctive syllogism)

What does a law of deduction look like?

Examples.

- ❶ (Modus Ponens) = “The mode that affirms”.

$$\frac{(P \rightarrow Q), P}{Q}$$

So if your proof has the structure $S_1, S_2, \dots, P, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step by appending $S_{k+1} = Q$ to this proof.

- ❷ (Modus Tollens) = “The mode that denies”.

$$\frac{(P \rightarrow Q), \neg Q}{\neg P}$$

So if your proof has the structure $S_1, S_2, \dots, \neg Q, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step to get $S_1, S_2, \dots, \neg Q, \dots, P \rightarrow Q, \dots, S_k, \neg P$.

- ❸ (Axioms) $\frac{}{A}$.

- ❹ (Hypothetical syllogism) $\frac{(P \rightarrow Q), (Q \rightarrow R)}{(P \rightarrow R)}$

- ❺ (Disjunctive syllogism)

What does a law of deduction look like?

Examples.

- ❶ (Modus Ponens) = “The mode that affirms”.

$$\frac{(P \rightarrow Q), P}{Q}$$

So if your proof has the structure $S_1, S_2, \dots, P, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step by appending $S_{k+1} = Q$ to this proof.

- ❷ (Modus Tollens) = “The mode that denies”.

$$\frac{(P \rightarrow Q), \neg Q}{\neg P}$$

So if your proof has the structure $S_1, S_2, \dots, \neg Q, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step to get $S_1, S_2, \dots, \neg Q, \dots, P \rightarrow Q, \dots, S_k, \neg P$.

- ❸ (Axioms) $\frac{}{A}$.

- ❹ (Hypothetical syllogism) $\frac{(P \rightarrow Q), (Q \rightarrow R)}{(P \rightarrow R)}$

- ❺ (Disjunctive syllogism) $\frac{(P \vee Q), \neg P}{Q}$

What does a law of deduction look like?

Examples.

- ❶ (Modus Ponens) = “The mode that affirms”.

$$\frac{(P \rightarrow Q), P}{Q}$$

So if your proof has the structure $S_1, S_2, \dots, P, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step by appending $S_{k+1} = Q$ to this proof.

- ❷ (Modus Tollens) = “The mode that denies”.

$$\frac{(P \rightarrow Q), \neg Q}{\neg P}$$

So if your proof has the structure $S_1, S_2, \dots, \neg Q, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step to get $S_1, S_2, \dots, \neg Q, \dots, P \rightarrow Q, \dots, S_k, \neg P$.

- ❸ (Axioms) $\frac{}{A}$.

- ❹ (Hypothetical syllogism) $\frac{(P \rightarrow Q), (Q \rightarrow R)}{(P \rightarrow R)}$

- ❺ (Disjunctive syllogism) $\frac{(P \vee Q), \neg P}{Q}$

- ❻ (Case analysis)

What does a law of deduction look like?

Examples.

- ❶ (Modus Ponens) = “The mode that affirms”.

$$\frac{(P \rightarrow Q), P}{Q}$$

So if your proof has the structure $S_1, S_2, \dots, P, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step by appending $S_{k+1} = Q$ to this proof.

- ❷ (Modus Tollens) = “The mode that denies”.

$$\frac{(P \rightarrow Q), \neg Q}{\neg P}$$

So if your proof has the structure $S_1, S_2, \dots, \neg Q, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step to get $S_1, S_2, \dots, \neg Q, \dots, P \rightarrow Q, \dots, S_k, \neg P$.

- ❸ (Axioms) $\frac{}{A}$.

- ❹ (Hypothetical syllogism) $\frac{(P \rightarrow Q), (Q \rightarrow R)}{(P \rightarrow R)}$

- ❺ (Disjunctive syllogism) $\frac{(P \vee Q), \neg P}{Q}$

- ❻ (Case analysis)

What does a law of deduction look like?

Examples.

- ❶ (Modus Ponens) = “The mode that affirms”.

$$\frac{(P \rightarrow Q), P}{Q}$$

So if your proof has the structure $S_1, S_2, \dots, P, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step by appending $S_{k+1} = Q$ to this proof.

- ❷ (Modus Tollens) = “The mode that denies”.

$$\frac{(P \rightarrow Q), \neg Q}{\neg P}$$

So if your proof has the structure $S_1, S_2, \dots, \neg Q, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step to get $S_1, S_2, \dots, \neg Q, \dots, P \rightarrow Q, \dots, S_k, \neg P$.

- ❸ (Axioms) $\frac{}{A}$.

- ❹ (Hypothetical syllogism) $\frac{(P \rightarrow Q), (Q \rightarrow R)}{(P \rightarrow R)}$

- ❺ (Disjunctive syllogism) $\frac{(P \vee Q), \neg P}{Q}$

- ❻ (Case analysis) $\frac{(P \rightarrow R), (Q \rightarrow R), (P \vee Q)}{R}$

What does a law of deduction look like?

Examples.

- ❶ (Modus Ponens) = “The mode that affirms”.

$$\frac{(P \rightarrow Q), P}{Q}$$

So if your proof has the structure $S_1, S_2, \dots, P, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step by appending $S_{k+1} = Q$ to this proof.

- ❷ (Modus Tollens) = “The mode that denies”.

$$\frac{(P \rightarrow Q), \neg Q}{\neg P}$$

So if your proof has the structure $S_1, S_2, \dots, \neg Q, \dots, P \rightarrow Q, \dots, S_k$, then you may continue it one step to get $S_1, S_2, \dots, \neg Q, \dots, P \rightarrow Q, \dots, S_k, \neg P$.

- ❸ (Axioms) $\frac{}{A}$.

- ❹ (Hypothetical syllogism) $\frac{(P \rightarrow Q), (Q \rightarrow R)}{(P \rightarrow R)}$

- ❺ (Disjunctive syllogism) $\frac{(P \vee Q), \neg P}{Q}$

- ❻ (Case analysis) $\frac{(P \rightarrow R), (Q \rightarrow R), (P \vee Q)}{R}$

Silly proof systems

Silly proof systems

Consider a proof system with no axioms and no rules of deduction.

Silly proof systems

Consider a proof system with no axioms and no rules of deduction. This is a **sound** proof system, since you can't prove any false statements.

Silly proof systems

Consider a proof system with no axioms and no rules of deduction. This is a **sound** proof system, since you can't prove any false statements. (You also can't prove any true statements!)

Silly proof systems

Consider a proof system with no axioms and no rules of deduction. This is a **sound** proof system, since you can't prove any false statements. (You also can't prove any true statements!)

Consider a proof system in which every statement is taken to be an axiom.

Silly proof systems

Consider a proof system with no axioms and no rules of deduction. This is a **sound** proof system, since you can't prove any false statements. (You also can't prove any true statements!)

Consider a proof system in which every statement is taken to be an axiom. This is an **unsound** proof system, since you do prove false statements.

Silly proof systems

Consider a proof system with no axioms and no rules of deduction. This is a **sound** proof system, since you can't prove any false statements. (You also can't prove any true statements!)

Consider a proof system in which every statement is taken to be an axiom. This is an **unsound** proof system, since you do prove false statements. (In fact, any statement T , even a contradictory statement, has a proof of length 1, namely " T ".

Silly proof systems

Consider a proof system with no axioms and no rules of deduction. This is a **sound** proof system, since you can't prove any false statements. (You also can't prove any true statements!)

Consider a proof system in which every statement is taken to be an axiom. This is an **unsound** proof system, since you do prove false statements. (In fact, any statement T , even a contradictory statement, has a proof of length 1, namely " T ".

Definitions.

Silly proof systems

Consider a proof system with no axioms and no rules of deduction. This is a **sound** proof system, since you can't prove any false statements. (You also can't prove any true statements!)

Consider a proof system in which every statement is taken to be an axiom. This is an **unsound** proof system, since you do prove false statements. (In fact, any statement T , even a contradictory statement, has a proof of length 1, namely " T ".

Definitions. A proof system is **sound** if it does not prove any false statements.

Silly proof systems

Consider a proof system with no axioms and no rules of deduction. This is a **sound** proof system, since you can't prove any false statements. (You also can't prove any true statements!)

Consider a proof system in which every statement is taken to be an axiom. This is an **unsound** proof system, since you do prove false statements. (In fact, any statement T , even a contradictory statement, has a proof of length 1, namely " T ".

Definitions. A proof system is **sound** if it does not prove any false statements. A proof system is **complete** if it proves every true statement.

Silly proof systems

Consider a proof system with no axioms and no rules of deduction. This is a **sound** proof system, since you can't prove any false statements. (You also can't prove any true statements!)

Consider a proof system in which every statement is taken to be an axiom. This is an **unsound** proof system, since you do prove false statements. (In fact, any statement T , even a contradictory statement, has a proof of length 1, namely " T ".

Definitions. A proof system is **sound** if it does not prove any false statements. A proof system is **complete** if it proves every true statement.

Gödel's Completeness Theorem

Gödel's Completeness Theorem

In 1929, Kurt Gödel introduced a proof system for first-order logic.

Gödel's Completeness Theorem

In 1929, Kurt Gödel introduced a proof system for first-order logic. It has finitely many types of axioms and finitely many rules of deduction.

Gödel's Completeness Theorem

In 1929, Kurt Gödel introduced a proof system for first-order logic. It has finitely many types of axioms and finitely many rules of deduction.

Gödel's Completeness Theorem. If $\Sigma \cup \{P\}$ consists of first-order sentences, then $\Sigma \models P$ if and only if $\Sigma \vdash P$.

Gödel's Completeness Theorem

In 1929, Kurt Gödel introduced a proof system for first-order logic. It has finitely many types of axioms and finitely many rules of deduction.

Gödel's Completeness Theorem. If $\Sigma \cup \{P\}$ consists of first-order sentences, then $\Sigma \models P$ if and only if $\Sigma \vdash P$.

“First-order sentences” are well-formed sentences that have finite length

Gödel's Completeness Theorem

In 1929, Kurt Gödel introduced a proof system for first-order logic. It has finitely many types of axioms and finitely many rules of deduction.

Gödel's Completeness Theorem. If $\Sigma \cup \{P\}$ consists of first-order sentences, then $\Sigma \models P$ if and only if $\Sigma \vdash P$.

“First-order sentences” are well-formed sentences that have finite length and are expressible with quantifiers $(\forall x)$ and $(\exists y)$ “of the first-order”.

Gödel's Completeness Theorem

In 1929, Kurt Gödel introduced a proof system for first-order logic. It has finitely many types of axioms and finitely many rules of deduction.

Gödel's Completeness Theorem. If $\Sigma \cup \{P\}$ consists of first-order sentences, then $\Sigma \models P$ if and only if $\Sigma \vdash P$.

“First-order sentences” are well-formed sentences that have finite length and are expressible with quantifiers $(\forall x)$ and $(\exists y)$ “of the first-order”. First-order quantifiers apply to individual elements of a structure, but not “higher-order” entities like subsets, functions, relations, sets of sets, etc.

Gödel's Completeness Theorem

In 1929, Kurt Gödel introduced a proof system for first-order logic. It has finitely many types of axioms and finitely many rules of deduction.

Gödel's Completeness Theorem. If $\Sigma \cup \{P\}$ consists of first-order sentences, then $\Sigma \models P$ if and only if $\Sigma \vdash P$.

“First-order sentences” are well-formed sentences that have finite length and are expressible with quantifiers $(\forall x)$ and $(\exists y)$ “of the first-order”.

First-order quantifiers apply to individual elements of a structure, but not “higher-order” entities like subsets, functions, relations, sets of sets, etc.

This result should be interpreted to mean that,

Gödel's Completeness Theorem

In 1929, Kurt Gödel introduced a proof system for first-order logic. It has finitely many types of axioms and finitely many rules of deduction.

Gödel's Completeness Theorem. If $\Sigma \cup \{P\}$ consists of first-order sentences, then $\Sigma \models P$ if and only if $\Sigma \vdash P$.

“First-order sentences” are well-formed sentences that have finite length and are expressible with quantifiers $(\forall x)$ and $(\exists y)$ “of the first-order”.

First-order quantifiers apply to individual elements of a structure, but not “higher-order” entities like subsets, functions, relations, sets of sets, etc.

This result should be interpreted to mean that, at the first-order level,

Gödel's Completeness Theorem

In 1929, Kurt Gödel introduced a proof system for first-order logic. It has finitely many types of axioms and finitely many rules of deduction.

Gödel's Completeness Theorem. If $\Sigma \cup \{P\}$ consists of first-order sentences, then $\Sigma \models P$ if and only if $\Sigma \vdash P$.

“First-order sentences” are well-formed sentences that have finite length and are expressible with quantifiers $(\forall x)$ and $(\exists y)$ “of the first-order”.

First-order quantifiers apply to individual elements of a structure, but not “higher-order” entities like subsets, functions, relations, sets of sets, etc.

This result should be interpreted to mean that, at the first-order level, if a statement is provable, then it is true, and if it is true then it is provable.

Gödel's Completeness Theorem

In 1929, Kurt Gödel introduced a proof system for first-order logic. It has finitely many types of axioms and finitely many rules of deduction.

Gödel's Completeness Theorem. If $\Sigma \cup \{P\}$ consists of first-order sentences, then $\Sigma \models P$ if and only if $\Sigma \vdash P$.

“First-order sentences” are well-formed sentences that have finite length and are expressible with quantifiers $(\forall x)$ and $(\exists y)$ “of the first-order”.

First-order quantifiers apply to individual elements of a structure, but not “higher-order” entities like subsets, functions, relations, sets of sets, etc.

This result should be interpreted to mean that, at the first-order level, if a statement is provable, then it is true, and if it is true then it is provable.

Another way to think about this is: at the first-order level, every statement has a proof or a counterexample.

An almost counter-example to Gödel's Theorem

An almost counter-example to Gödel's Theorem

Theorem.

An almost counter-example to Gödel's Theorem

Theorem. If X is infinite, then X is infinite.

An almost counter-example to Gödel's Theorem

Theorem. If X is infinite, then X is infinite.

Recall that “ X is infinite” means that X is not finite,

An almost counter-example to Gödel's Theorem

Theorem. If X is infinite, then X is infinite.

Recall that “ X is infinite” means that X is not finite, which is expressible with the set $\Sigma = \{P_0, P_1, P_2, \dots\}$ of sentences, where P_n is “ $|X| \neq n$ ”,

An almost counter-example to Gödel's Theorem

Theorem. If X is infinite, then X is infinite.

Recall that “ X is infinite” means that X is not finite, which is expressible with the set $\Sigma = \{P_0, P_1, P_2, \dots\}$ of sentences, where P_n is “ $|X| \neq n$ ”, or “There is no bijection $f: n \rightarrow X$ ”.

An almost counter-example to Gödel's Theorem

Theorem. If X is infinite, then X is infinite.

Recall that “ X is infinite” means that X is not finite, which is expressible with the set $\Sigma = \{P_0, P_1, P_2, \dots\}$ of sentences, where P_n is “ $|X| \neq n$ ”, or “There is no bijection $f: n \rightarrow X$ ”. We might also express this with a single sentence Q which says

$$(\forall n)(\forall f)(\neg(f: n \rightarrow X \text{ is a bijection})).$$

An almost counter-example to Gödel's Theorem

Theorem. If X is infinite, then X is infinite.

Recall that “ X is infinite” means that X is not finite, which is expressible with the set $\Sigma = \{P_0, P_1, P_2, \dots\}$ of sentences, where P_n is “ $|X| \neq n$ ”, or “There is no bijection $f: n \rightarrow X$ ”. We might also express this with a single sentence Q which says

$$(\forall n)(\forall f)(\neg(f: n \rightarrow X \text{ is a bijection})).$$

Then $\Sigma \models Q$, but $\Sigma \not\vdash Q$ for any proof system requiring finite-length proofs.

The structure of a theorem statement

The structure of a theorem statement

Most common structure:

The structure of a theorem statement

Most common structure:

Theorem.

The structure of a theorem statement

Most common structure:

Theorem. $H \rightarrow C$.

The structure of a theorem statement

Most common structure:

Theorem. $H \rightarrow C$. (Hypothesis implies conclusion.)

The structure of a theorem statement

Most common structure:

Theorem. $H \rightarrow C$. (Hypothesis implies conclusion.)

There might be more than one hypothesis.

The structure of a theorem statement

Most common structure:

Theorem. $H \rightarrow C$. (Hypothesis implies conclusion.)

There might be more than one hypothesis.

Theorem.

The structure of a theorem statement

Most common structure:

Theorem. $H \rightarrow C$. (Hypothesis implies conclusion.)

There might be more than one hypothesis.

Theorem. $(H_1 \wedge H_2) \rightarrow C$.

The structure of a theorem statement

Most common structure:

Theorem. $H \rightarrow C$. (Hypothesis implies conclusion.)

There might be more than one hypothesis.

Theorem. $(H_1 \wedge H_2) \rightarrow C$. (Or even $(H_1 \wedge \cdots \wedge H_k) \rightarrow C$.)

The structure of a theorem statement

Most common structure:

Theorem. $H \rightarrow C$. (Hypothesis implies conclusion.)

There might be more than one hypothesis.

Theorem. $(H_1 \wedge H_2) \rightarrow C$. (Or even $(H_1 \wedge \cdots \wedge H_k) \rightarrow C$.)

The hypotheses might be “hidden” or “understood”.

The structure of a theorem statement

Most common structure:

Theorem. $H \rightarrow C$. (Hypothesis implies conclusion.)

There might be more than one hypothesis.

Theorem. $(H_1 \wedge H_2) \rightarrow C$. (Or even $(H_1 \wedge \cdots \wedge H_k) \rightarrow C$.)

The hypotheses might be “hidden” or “understood”.

Theorem.

The structure of a theorem statement

Most common structure:

Theorem. $H \rightarrow C$. (Hypothesis implies conclusion.)

There might be more than one hypothesis.

Theorem. $(H_1 \wedge H_2) \rightarrow C$. (Or even $(H_1 \wedge \cdots \wedge H_k) \rightarrow C$.)

The hypotheses might be “hidden” or “understood”.

Theorem. C .

The structure of a theorem statement

Most common structure:

Theorem. $H \rightarrow C$. (Hypothesis implies conclusion.)

There might be more than one hypothesis.

Theorem. $(H_1 \wedge H_2) \rightarrow C$. (Or even $(H_1 \wedge \dots \wedge H_k) \rightarrow C$.)

The hypotheses might be “hidden” or “understood”.

Theorem. C .

(That is, C is a consequence of some unspecified set of axioms and previously-established theorems.)

The structure of a theorem statement

Most common structure:

Theorem. $H \rightarrow C$. (Hypothesis implies conclusion.)

There might be more than one hypothesis.

Theorem. $(H_1 \wedge H_2) \rightarrow C$. (Or even $(H_1 \wedge \dots \wedge H_k) \rightarrow C$.)

The hypotheses might be “hidden” or “understood”.

Theorem. C .

(That is, C is a consequence of some unspecified set of axioms and previously-established theorems.)

Examples.

The structure of a theorem statement

Most common structure:

Theorem. $H \rightarrow C$. (Hypothesis implies conclusion.)

There might be more than one hypothesis.

Theorem. $(H_1 \wedge H_2) \rightarrow C$. (Or even $(H_1 \wedge \cdots \wedge H_k) \rightarrow C$.)

The hypotheses might be “hidden” or “understood”.

Theorem. C .

(That is, C is a consequence of some unspecified set of axioms and previously-established theorems.)

Examples.

❶ **Theorem.**

The structure of a theorem statement

Most common structure:

Theorem. $H \rightarrow C$. (Hypothesis implies conclusion.)

There might be more than one hypothesis.

Theorem. $(H_1 \wedge H_2) \rightarrow C$. (Or even $(H_1 \wedge \dots \wedge H_k) \rightarrow C$.)

The hypotheses might be “hidden” or “understood”.

Theorem. C .

(That is, C is a consequence of some unspecified set of axioms and previously-established theorems.)

Examples.

❶ **Theorem.**

The structure of a theorem statement

Most common structure:

Theorem. $H \rightarrow C$. (Hypothesis implies conclusion.)

There might be more than one hypothesis.

Theorem. $(H_1 \wedge H_2) \rightarrow C$. (Or even $(H_1 \wedge \cdots \wedge H_k) \rightarrow C$.)

The hypotheses might be “hidden” or “understood”.

Theorem. C .

(That is, C is a consequence of some unspecified set of axioms and previously-established theorems.)

Examples.

① **Theorem.** If $0 < x < 1$, then $x^2 < x$.

The structure of a theorem statement

Most common structure:

Theorem. $H \rightarrow C$. (Hypothesis implies conclusion.)

There might be more than one hypothesis.

Theorem. $(H_1 \wedge H_2) \rightarrow C$. (Or even $(H_1 \wedge \cdots \wedge H_k) \rightarrow C$.)

The hypotheses might be “hidden” or “understood”.

Theorem. C .

(That is, C is a consequence of some unspecified set of axioms and previously-established theorems.)

Examples.

① **Theorem.** If $0 < x < 1$, then $x^2 < x$.

② **Theorem.**

The structure of a theorem statement

Most common structure:

Theorem. $H \rightarrow C$. (Hypothesis implies conclusion.)

There might be more than one hypothesis.

Theorem. $(H_1 \wedge H_2) \rightarrow C$. (Or even $(H_1 \wedge \cdots \wedge H_k) \rightarrow C$.)

The hypotheses might be “hidden” or “understood”.

Theorem. C .

(That is, C is a consequence of some unspecified set of axioms and previously-established theorems.)

Examples.

① **Theorem.** If $0 < x < 1$, then $x^2 < x$.

② **Theorem.**

The structure of a theorem statement

Most common structure:

Theorem. $H \rightarrow C$. (Hypothesis implies conclusion.)

There might be more than one hypothesis.

Theorem. $(H_1 \wedge H_2) \rightarrow C$. (Or even $(H_1 \wedge \cdots \wedge H_k) \rightarrow C$.)

The hypotheses might be “hidden” or “understood”.

Theorem. C .

(That is, C is a consequence of some unspecified set of axioms and previously-established theorems.)

Examples.

① **Theorem.** If $0 < x < 1$, then $x^2 < x$.

② **Theorem.** If g is a surjective function, f is a surjective function, and g and f are composable, then $g \circ f$ is a surjective function.

The structure of a theorem statement

Most common structure:

Theorem. $H \rightarrow C$. (Hypothesis implies conclusion.)

There might be more than one hypothesis.

Theorem. $(H_1 \wedge H_2) \rightarrow C$. (Or even $(H_1 \wedge \cdots \wedge H_k) \rightarrow C$.)

The hypotheses might be “hidden” or “understood”.

Theorem. C .

(That is, C is a consequence of some unspecified set of axioms and previously-established theorems.)

Examples.

① **Theorem.** If $0 < x < 1$, then $x^2 < x$.

② **Theorem.** If g is a surjective function, f is a surjective function, and g and f are composable, then $g \circ f$ is a surjective function.

③ **Theorem.**

The structure of a theorem statement

Most common structure:

Theorem. $H \rightarrow C$. (Hypothesis implies conclusion.)

There might be more than one hypothesis.

Theorem. $(H_1 \wedge H_2) \rightarrow C$. (Or even $(H_1 \wedge \cdots \wedge H_k) \rightarrow C$.)

The hypotheses might be “hidden” or “understood”.

Theorem. C .

(That is, C is a consequence of some unspecified set of axioms and previously-established theorems.)

Examples.

① **Theorem.** If $0 < x < 1$, then $x^2 < x$.

② **Theorem.** If g is a surjective function, f is a surjective function, and g and f are composable, then $g \circ f$ is a surjective function.

③ **Theorem.**

The structure of a theorem statement

Most common structure:

Theorem. $H \rightarrow C$. (Hypothesis implies conclusion.)

There might be more than one hypothesis.

Theorem. $(H_1 \wedge H_2) \rightarrow C$. (Or even $(H_1 \wedge \cdots \wedge H_k) \rightarrow C$.)

The hypotheses might be “hidden” or “understood”.

Theorem. C .

(That is, C is a consequence of some unspecified set of axioms and previously-established theorems.)

Examples.

① **Theorem.** If $0 < x < 1$, then $x^2 < x$.

② **Theorem.** If g is a surjective function, f is a surjective function, and g and f are composable, then $g \circ f$ is a surjective function.

③ **Theorem.** There are infinitely many primes.

The structure of a theorem statement

Most common structure:

Theorem. $H \rightarrow C$. (Hypothesis implies conclusion.)

There might be more than one hypothesis.

Theorem. $(H_1 \wedge H_2) \rightarrow C$. (Or even $(H_1 \wedge \cdots \wedge H_k) \rightarrow C$.)

The hypotheses might be “hidden” or “understood”.

Theorem. C .

(That is, C is a consequence of some unspecified set of axioms and previously-established theorems.)

Examples.

① **Theorem.** If $0 < x < 1$, then $x^2 < x$.

② **Theorem.** If g is a surjective function, f is a surjective function, and g and f are composable, then $g \circ f$ is a surjective function.

③ **Theorem.** There are infinitely many primes.

Simple proof strategies

Simple proof strategies

(Hypotheses are treated as if they were axioms, and previously-proved theorems are also treated as if they were axioms.)

Simple proof strategies

(Hypotheses are treated as if they were axioms, and previously-proved theorems are also treated as if they were axioms.)

Theorem.

Simple proof strategies

(Hypotheses are treated as if they were axioms, and previously-proved theorems are also treated as if they were axioms.)

Theorem. $H \rightarrow C$.

Simple proof strategies

(Hypotheses are treated as if they were axioms, and previously-proved theorems are also treated as if they were axioms.)

Theorem. $H \rightarrow C$.

Proof structure #1.

$H = S_1, S_2, \dots, S_k = C$.

Simple proof strategies

(Hypotheses are treated as if they were axioms, and previously-proved theorems are also treated as if they were axioms.)

Theorem. $H \rightarrow C$.

Proof structure #1.

$H = S_1, S_2, \dots, S_k = C. \square$

Simple proof strategies

(Hypotheses are treated as if they were axioms, and previously-proved theorems are also treated as if they were axioms.)

Theorem. $H \rightarrow C$.

Proof structure #1.

$H = S_1, S_2, \dots, S_k = C. \square$ (Direct proof.)

Simple proof strategies

(Hypotheses are treated as if they were axioms, and previously-proved theorems are also treated as if they were axioms.)

Theorem. $H \rightarrow C$.

Proof structure #1.

$H = S_1, S_2, \dots, S_k = C. \square$ (Direct proof.)

Proof structure #2.

$\neg C, S_2, \dots, \neg H.$

Simple proof strategies

(Hypotheses are treated as if they were axioms, and previously-proved theorems are also treated as if they were axioms.)

Theorem. $H \rightarrow C$.

Proof structure #1.

$H = S_1, S_2, \dots, S_k = C. \square$ (Direct proof.)

Proof structure #2.

$\neg C, S_2, \dots, \neg H. \square$

Simple proof strategies

(Hypotheses are treated as if they were axioms, and previously-proved theorems are also treated as if they were axioms.)

Theorem. $H \rightarrow C$.

Proof structure #1.

$H = S_1, S_2, \dots, S_k = C. \square$ (Direct proof.)

Proof structure #2.

$\neg C, S_2, \dots, \neg H. \square$ (Proof by contraposition,

Simple proof strategies

(Hypotheses are treated as if they were axioms, and previously-proved theorems are also treated as if they were axioms.)

Theorem. $H \rightarrow C$.

Proof structure #1.

$H = S_1, S_2, \dots, S_k = C. \square$ (Direct proof.)

Proof structure #2.

$\neg C, S_2, \dots, \neg H. \square$ (Proof by contraposition, or direct proof of the **contrapositive** statement $(\neg C) \rightarrow (\neg H)$.)

Simple proof strategies

(Hypotheses are treated as if they were axioms, and previously-proved theorems are also treated as if they were axioms.)

Theorem. $H \rightarrow C$.

Proof structure #1.

$H = S_1, S_2, \dots, S_k = C. \square$ (Direct proof.)

Proof structure #2.

$\neg C, S_2, \dots, \neg H. \square$ (Proof by contraposition, or direct proof of the **contrapositive** statement $(\neg C) \rightarrow (\neg H)$.)

Proof structure #3.

$H, \neg C, S_3, \dots, \perp.$

Simple proof strategies

(Hypotheses are treated as if they were axioms, and previously-proved theorems are also treated as if they were axioms.)

Theorem. $H \rightarrow C$.

Proof structure #1.

$H = S_1, S_2, \dots, S_k = C. \square$ (Direct proof.)

Proof structure #2.

$\neg C, S_2, \dots, \neg H. \square$ (Proof by contraposition, or direct proof of the **contrapositive** statement $(\neg C) \rightarrow (\neg H)$.)

Proof structure #3.

$H, \neg C, S_3, \dots, \perp. \square$

Simple proof strategies

(Hypotheses are treated as if they were axioms, and previously-proved theorems are also treated as if they were axioms.)

Theorem. $H \rightarrow C$.

Proof structure #1.

$H = S_1, S_2, \dots, S_k = C. \square$ (Direct proof.)

Proof structure #2.

$\neg C, S_2, \dots, \neg H. \square$ (Proof by contraposition, or direct proof of the **contrapositive** statement $(\neg C) \rightarrow (\neg H)$.)

Proof structure #3.

$H, \neg C, S_3, \dots, \perp. \square$ (Proof by contradiction.)

Examples!

Examples!

Let's try these three forms of proof on a theorem concerning \mathbb{R} .

Examples!

Let's try these three forms of proof on a theorem concerning \mathbb{R} .

Theorem.

Examples!

Let's try these three forms of proof on a theorem concerning \mathbb{R} .

Theorem. If $0 < x < 1$, then $x^2 < x$.

Examples!

Let's try these three forms of proof on a theorem concerning \mathbb{R} .

Theorem. If $0 < x < 1$, then $x^2 < x$.

Proof structure #1.

Examples!

Let's try these three forms of proof on a theorem concerning \mathbb{R} .

Theorem. If $0 < x < 1$, then $x^2 < x$.

Proof structure #1.

Assume that $0 < x < 1$.

Examples!

Let's try these three forms of proof on a theorem concerning \mathbb{R} .

Theorem. If $0 < x < 1$, then $x^2 < x$.

Proof structure #1.

Assume that $0 < x < 1$. Since $0 < x$, multiplication by x preserves inequalities.

Examples!

Let's try these three forms of proof on a theorem concerning \mathbb{R} .

Theorem. If $0 < x < 1$, then $x^2 < x$.

Proof structure #1.

Assume that $0 < x < 1$. Since $0 < x$, multiplication by x preserves inequalities. Multiply $x < 1$ by x to obtain $x^2 < x$. \square

Examples!

Let's try these three forms of proof on a theorem concerning \mathbb{R} .

Theorem. If $0 < x < 1$, then $x^2 < x$.

Proof structure #1.

Assume that $0 < x < 1$. Since $0 < x$, multiplication by x preserves inequalities. Multiply $x < 1$ by x to obtain $x^2 < x$. \square

Proof structure #2.

Examples!

Let's try these three forms of proof on a theorem concerning \mathbb{R} .

Theorem. If $0 < x < 1$, then $x^2 < x$.

Proof structure #1.

Assume that $0 < x < 1$. Since $0 < x$, multiplication by x preserves inequalities. Multiply $x < 1$ by x to obtain $x^2 < x$. \square

Proof structure #2.

Assume that $x^2 \not< x$.

Examples!

Let's try these three forms of proof on a theorem concerning \mathbb{R} .

Theorem. If $0 < x < 1$, then $x^2 < x$.

Proof structure #1.

Assume that $0 < x < 1$. Since $0 < x$, multiplication by x preserves inequalities. Multiply $x < 1$ by x to obtain $x^2 < x$. \square

Proof structure #2.

Assume that $x^2 \not< x$. Then $x \leq x^2$.

Examples!

Let's try these three forms of proof on a theorem concerning \mathbb{R} .

Theorem. If $0 < x < 1$, then $x^2 < x$.

Proof structure #1.

Assume that $0 < x < 1$. Since $0 < x$, multiplication by x preserves inequalities. Multiply $x < 1$ by x to obtain $x^2 < x$. \square

Proof structure #2.

Assume that $x^2 \not< x$. Then $x \leq x^2$. Hence $0 \leq x^2 - x = x(x - 1)$.

Examples!

Let's try these three forms of proof on a theorem concerning \mathbb{R} .

Theorem. If $0 < x < 1$, then $x^2 < x$.

Proof structure #1.

Assume that $0 < x < 1$. Since $0 < x$, multiplication by x preserves inequalities. Multiply $x < 1$ by x to obtain $x^2 < x$. \square

Proof structure #2.

Assume that $x^2 \not< x$. Then $x \leq x^2$. Hence $0 \leq x^2 - x = x(x - 1)$. Hence $0 \leq x, x - 1$ or $x, x - 1 \leq 0$.

Examples!

Let's try these three forms of proof on a theorem concerning \mathbb{R} .

Theorem. If $0 < x < 1$, then $x^2 < x$.

Proof structure #1.

Assume that $0 < x < 1$. Since $0 < x$, multiplication by x preserves inequalities. Multiply $x < 1$ by x to obtain $x^2 < x$. \square

Proof structure #2.

Assume that $x^2 \not< x$. Then $x \leq x^2$. Hence $0 \leq x^2 - x = x(x - 1)$. Hence $0 \leq x, x - 1$ or $x, x - 1 \leq 0$. The first leads to $0 \leq x - 1$, or $1 \leq x$,

Examples!

Let's try these three forms of proof on a theorem concerning \mathbb{R} .

Theorem. If $0 < x < 1$, then $x^2 < x$.

Proof structure #1.

Assume that $0 < x < 1$. Since $0 < x$, multiplication by x preserves inequalities. Multiply $x < 1$ by x to obtain $x^2 < x$. \square

Proof structure #2.

Assume that $x^2 \not< x$. Then $x \leq x^2$. Hence $0 \leq x^2 - x = x(x - 1)$. Hence $0 \leq x, x - 1$ or $x, x - 1 \leq 0$. The first leads to $0 \leq x - 1$, or $1 \leq x$, while the second leads to $x \leq 0$.

Examples!

Let's try these three forms of proof on a theorem concerning \mathbb{R} .

Theorem. If $0 < x < 1$, then $x^2 < x$.

Proof structure #1.

Assume that $0 < x < 1$. Since $0 < x$, multiplication by x preserves inequalities. Multiply $x < 1$ by x to obtain $x^2 < x$. \square

Proof structure #2.

Assume that $x^2 \not< x$. Then $x \leq x^2$. Hence $0 \leq x^2 - x = x(x - 1)$. Hence $0 \leq x, x - 1$ or $x, x - 1 \leq 0$. The first leads to $0 \leq x - 1$, or $1 \leq x$, while the second leads to $x \leq 0$. Either way, $0 < x < 1$ fails.

Examples!

Let's try these three forms of proof on a theorem concerning \mathbb{R} .

Theorem. If $0 < x < 1$, then $x^2 < x$.

Proof structure #1.

Assume that $0 < x < 1$. Since $0 < x$, multiplication by x preserves inequalities. Multiply $x < 1$ by x to obtain $x^2 < x$. \square

Proof structure #2.

Assume that $x^2 \not< x$. Then $x \leq x^2$. Hence $0 \leq x^2 - x = x(x - 1)$. Hence $0 \leq x, x - 1$ or $x, x - 1 \leq 0$. The first leads to $0 \leq x - 1$, or $1 \leq x$, while the second leads to $x \leq 0$. Either way, $0 < x < 1$ fails.

Proof structure #3.

Examples!

Let's try these three forms of proof on a theorem concerning \mathbb{R} .

Theorem. If $0 < x < 1$, then $x^2 < x$.

Proof structure #1.

Assume that $0 < x < 1$. Since $0 < x$, multiplication by x preserves inequalities. Multiply $x < 1$ by x to obtain $x^2 < x$. \square

Proof structure #2.

Assume that $x^2 \not< x$. Then $x \leq x^2$. Hence $0 \leq x^2 - x = x(x - 1)$. Hence $0 \leq x, x - 1$ or $x, x - 1 \leq 0$. The first leads to $0 \leq x - 1$, or $1 \leq x$, while the second leads to $x \leq 0$. Either way, $0 < x < 1$ fails.

Proof structure #3.

Assume that $0 < x < 1$ and $x^2 \not< x$.

Examples!

Let's try these three forms of proof on a theorem concerning \mathbb{R} .

Theorem. If $0 < x < 1$, then $x^2 < x$.

Proof structure #1.

Assume that $0 < x < 1$. Since $0 < x$, multiplication by x preserves inequalities. Multiply $x < 1$ by x to obtain $x^2 < x$. \square

Proof structure #2.

Assume that $x^2 \not< x$. Then $x \leq x^2$. Hence $0 \leq x^2 - x = x(x - 1)$. Hence $0 \leq x, x - 1$ or $x, x - 1 \leq 0$. The first leads to $0 \leq x - 1$, or $1 \leq x$, while the second leads to $x \leq 0$. Either way, $0 < x < 1$ fails.

Proof structure #3.

Assume that $0 < x < 1$ and $x^2 \not< x$. The first leads to $x > 0$ and $x - 1 < 0$, hence $x(x - 1) < 0$.

Examples!

Let's try these three forms of proof on a theorem concerning \mathbb{R} .

Theorem. If $0 < x < 1$, then $x^2 < x$.

Proof structure #1.

Assume that $0 < x < 1$. Since $0 < x$, multiplication by x preserves inequalities. Multiply $x < 1$ by x to obtain $x^2 < x$. \square

Proof structure #2.

Assume that $x^2 \not< x$. Then $x \leq x^2$. Hence $0 \leq x^2 - x = x(x - 1)$. Hence $0 \leq x, x - 1$ or $x, x - 1 \leq 0$. The first leads to $0 \leq x - 1$, or $1 \leq x$, while the second leads to $x \leq 0$. Either way, $0 < x < 1$ fails.

Proof structure #3.

Assume that $0 < x < 1$ and $x^2 \not< x$. The first leads to $x > 0$ and $x - 1 < 0$, hence $x(x - 1) < 0$. The second leads to $x^2 - x \not< 0$.

Examples!

Let's try these three forms of proof on a theorem concerning \mathbb{R} .

Theorem. If $0 < x < 1$, then $x^2 < x$.

Proof structure #1.

Assume that $0 < x < 1$. Since $0 < x$, multiplication by x preserves inequalities. Multiply $x < 1$ by x to obtain $x^2 < x$. \square

Proof structure #2.

Assume that $x^2 \not< x$. Then $x \leq x^2$. Hence $0 \leq x^2 - x = x(x - 1)$. Hence $0 \leq x, x - 1$ or $x, x - 1 \leq 0$. The first leads to $0 \leq x - 1$, or $1 \leq x$, while the second leads to $x \leq 0$. Either way, $0 < x < 1$ fails.

Proof structure #3.

Assume that $0 < x < 1$ and $x^2 \not< x$. The first leads to $x > 0$ and $x - 1 < 0$, hence $x(x - 1) < 0$. The second leads to $x^2 - x \not< 0$. These two statements contradict one another.

Examples!

Let's try these three forms of proof on a theorem concerning \mathbb{R} .

Theorem. If $0 < x < 1$, then $x^2 < x$.

Proof structure #1.

Assume that $0 < x < 1$. Since $0 < x$, multiplication by x preserves inequalities. Multiply $x < 1$ by x to obtain $x^2 < x$. \square

Proof structure #2.

Assume that $x^2 \not< x$. Then $x \leq x^2$. Hence $0 \leq x^2 - x = x(x - 1)$. Hence $0 \leq x, x - 1$ or $x, x - 1 \leq 0$. The first leads to $0 \leq x - 1$, or $1 \leq x$, while the second leads to $x \leq 0$. Either way, $0 < x < 1$ fails.

Proof structure #3.

Assume that $0 < x < 1$ and $x^2 \not< x$. The first leads to $x > 0$ and $x - 1 < 0$, hence $x(x - 1) < 0$. The second leads to $x^2 - x \not< 0$. These two statements contradict one another. \square

Proofs involving quantifiers

Proofs involving quantifiers

We can use quantifier games to construct a proof strategy for theorems involving quantifiers.

Proofs involving quantifiers

We can use quantifier games to construct a proof strategy for theorems involving quantifiers. Consider the following theorem, which is true for any nonempty structure and for any formula $\varphi(x)$.

Theorem.

Proofs involving quantifiers

We can use quantifier games to construct a proof strategy for theorems involving quantifiers. Consider the following theorem, which is true for any nonempty structure and for any formula $\varphi(x)$.

Theorem. If $(\forall x)\varphi(x)$, then $(\exists x)\varphi(x)$.

Proofs involving quantifiers

We can use quantifier games to construct a proof strategy for theorems involving quantifiers. Consider the following theorem, which is true for any nonempty structure and for any formula $\varphi(x)$.

Theorem. If $(\forall x)\varphi(x)$, then $(\exists x)\varphi(x)$.

Proof.

Proofs involving quantifiers

We can use quantifier games to construct a proof strategy for theorems involving quantifiers. Consider the following theorem, which is true for any nonempty structure and for any formula $\varphi(x)$.

Theorem. If $(\forall x)\varphi(x)$, then $(\exists x)\varphi(x)$.

Proof. We will give a winning strategy for \exists in the game determined by the sentence “ $(\exists x)\varphi(x)$ ”.

Proofs involving quantifiers

We can use quantifier games to construct a proof strategy for theorems involving quantifiers. Consider the following theorem, which is true for any nonempty structure and for any formula $\varphi(x)$.

Theorem. If $(\forall x)\varphi(x)$, then $(\exists x)\varphi(x)$.

Proof. We will give a winning strategy for \exists in the game determined by the sentence “ $(\exists x)\varphi(x)$ ”. In order to access the information in the hypothesis, we first play a “side game” using the strategy for \exists in the game determined by the sentence “ $(\forall x)\varphi(x)$ ”.

Proofs involving quantifiers

We can use quantifier games to construct a proof strategy for theorems involving quantifiers. Consider the following theorem, which is true for any nonempty structure and for any formula $\varphi(x)$.

Theorem. If $(\forall x)\varphi(x)$, then $(\exists x)\varphi(x)$.

Proof. We will give a winning strategy for \exists in the game determined by the sentence “ $(\exists x)\varphi(x)$ ”. In order to access the information in the hypothesis, we first play a “side game” using the strategy for \exists in the game determined by the sentence “ $(\forall x)\varphi(x)$ ”.

- \forall chooses some $x = r$. Loses. This means that $\varphi(r)$ holds.

Proofs involving quantifiers

We can use quantifier games to construct a proof strategy for theorems involving quantifiers. Consider the following theorem, which is true for any nonempty structure and for any formula $\varphi(x)$.

Theorem. If $(\forall x)\varphi(x)$, then $(\exists x)\varphi(x)$.

Proof. We will give a winning strategy for \exists in the game determined by the sentence “ $(\exists x)\varphi(x)$ ”. In order to access the information in the hypothesis, we first play a “side game” using the strategy for \exists in the game determined by the sentence “ $(\forall x)\varphi(x)$ ”.

- \forall chooses some $x = r$. Loses. This means that $\varphi(r)$ holds.

Proofs involving quantifiers

We can use quantifier games to construct a proof strategy for theorems involving quantifiers. Consider the following theorem, which is true for any nonempty structure and for any formula $\varphi(x)$.

Theorem. If $(\forall x)\varphi(x)$, then $(\exists x)\varphi(x)$.

Proof. We will give a winning strategy for \exists in the game determined by the sentence “ $(\exists x)\varphi(x)$ ”. In order to access the information in the hypothesis, we first play a “side game” using the strategy for \exists in the game determined by the sentence “ $(\forall x)\varphi(x)$ ”.

- \forall chooses some $x = r$. Loses. This means that $\varphi(r)$ holds.

Now we are in a position to provide a winning strategy for \exists in the game determined by the sentence “ $(\exists x)\varphi(x)$ ”.

Proofs involving quantifiers

We can use quantifier games to construct a proof strategy for theorems involving quantifiers. Consider the following theorem, which is true for any nonempty structure and for any formula $\varphi(x)$.

Theorem. If $(\forall x)\varphi(x)$, then $(\exists x)\varphi(x)$.

Proof. We will give a winning strategy for \exists in the game determined by the sentence “ $(\exists x)\varphi(x)$ ”. In order to access the information in the hypothesis, we first play a “side game” using the strategy for \exists in the game determined by the sentence “ $(\forall x)\varphi(x)$ ”.

- \forall chooses some $x = r$. Loses. This means that $\varphi(r)$ holds.

Now we are in a position to provide a winning strategy for \exists in the game determined by the sentence “ $(\exists x)\varphi(x)$ ”.

- \exists chooses $x = r$. Wins, because the side game guarantees that $\varphi(r)$ holds.

Proofs involving quantifiers, 2

Proofs involving quantifiers, 2

The previous proof can be written more informally as:

Theorem.

Proofs involving quantifiers, 2

The previous proof can be written more informally as:

Theorem. If $(\forall x)\varphi(x)$, then $(\exists x)\varphi(x)$.

Proofs involving quantifiers, 2

The previous proof can be written more informally as:

Theorem. If $(\forall x)\varphi(x)$, then $(\exists x)\varphi(x)$.

Proof.

Proofs involving quantifiers, 2

The previous proof can be written more informally as:

Theorem. If $(\forall x)\varphi(x)$, then $(\exists x)\varphi(x)$.

Proof. Choose any r .

Proofs involving quantifiers, 2

The previous proof can be written more informally as:

Theorem. If $(\forall x)\varphi(x)$, then $(\exists x)\varphi(x)$.

Proof. Choose any r . According to the hypothesis, $\varphi(r)$ holds.

Proofs involving quantifiers, 2

The previous proof can be written more informally as:

Theorem. If $(\forall x)\varphi(x)$, then $(\exists x)\varphi(x)$.

Proof. Choose any r . According to the hypothesis, $\varphi(r)$ holds. Therefore, $x = r$ is a witness to the fact that $(\exists x)\varphi(x)$ holds.

Proofs involving quantifiers, 2

The previous proof can be written more informally as:

Theorem. If $(\forall x)\varphi(x)$, then $(\exists x)\varphi(x)$.

Proof. Choose any r . According to the hypothesis, $\varphi(r)$ holds. Therefore, $x = r$ is a witness to the fact that $(\exists x)\varphi(x)$ holds. \square