

Solutions to HW 9.

1. This problem involves a deck of 52 distinct playing cards.

- (a) In how many ways can a 13-card bridge hand be dealt from the deck?

We want an ordered sequence of 13 cards chosen from a 52-element set. There are $13! \cdot \binom{52}{13} = (52)_{13} = \frac{52!}{39!}$ of these.

- (b) How many different 13-card bridge hands are there?

We want a 13-card subset of a 52-element set. There are $\binom{52}{13} = \frac{52!}{13!39!}$ of these.

2. (a) How many paths are there from the point $(0, 0)$ of \mathbb{R}^2 to the point $(10, 15)$ of \mathbb{R}^2 if each path consists of a sequence of steps of length 1 moving in the direction of the positive x -axis or the positive y -axis?

We can describe a path by a list of the form “ $(x, x, y, x, y, y, \dots, x, y, y)$ ”, which is a string of 10 x ’s and 15 y ’s in some order. If the first two instructions are x , then we take our first two steps in the x direction; if our next instruction is y , we take our next step in the y direction. ETC.

The number of paths is equal to the number of lists of instructions, which is equal to the number of strings of length $10 + 15 = 25$ consisting of x ’s and y ’s, which contain 10 x ’s and 15 y ’s. This number is $\binom{25}{10} = \frac{25!}{10!15!}$. (You have 25 instructions, and you “choose” 10 instructions to be x ’s and leave the remaining 15 to be y ’s.)

- (b) How many paths are there from the point $(0, 0, 0)$ of \mathbb{R}^3 to the point $(10, 15, 20)$ of \mathbb{R}^3 if each path consists of a sequence of steps of length 1 moving in the direction of the positive x -axis, the positive y -axis or the positive z -axis?

Using the same reasoning, we want to count strings of length $10 + 15 + 20 = 45$ which have 10 x ’s, 15 y ’s, and 20 z ’s. The number is $\binom{45}{10,15,20} = \frac{45!}{10!15!20!}$.

3. Let $MC(n, k)$ be the number “ n -multichoose- k ”. Use a combinatorial argument to show that $MC(n, 0) + MC(n, 1) + \dots + MC(n, k) = MC(n + 1, k)$.

Let D be the set of all distributions of k identical balls to $n + 1$ distinct boxes with repetition allowed. $|D| = MC(n + 1, k) = \binom{n+1+k-1}{k} = \binom{n+k}{k}$.

Partition D into sets D_0, D_1, \dots, D_k where $D_i \subseteq D$ is the number of distributions where i balls are distributed to the first n boxes, while the remaining $k - i$ balls are distributed to the last box. Since we can distribute i identical balls to the first n boxes in $MC(n, i)$ ways, and we have no choice but to put the remaining $k - i$ balls into the last box, we have $|D_i| = MC(n, i)$. Since we have a partition,

$$MC(n + 1, k) = |D| = |D_0| + |D_1| + \dots + |D_k| = MC(n, 0) + MC(n, 1) + \dots + MC(n, k).$$