

# Homomorphisms

# The concept

# The concept

In mathematics, we often use “structure-preserving” functions to compare objects.

# The concept

In mathematics, we often use “structure-preserving” functions to compare objects.

**Definition.**

# The concept

In mathematics, we often use “structure-preserving” functions to compare objects.

**Definition.** (Homomorphism)

# The concept

In mathematics, we often use “structure-preserving” functions to compare objects.

**Definition.** (Homomorphism)

If  $\mathbb{A} = \langle A; F^{\mathbb{A}} \rangle$  and  $\mathbb{B} = \langle B; F^{\mathbb{B}} \rangle$  are algebras of the same signature,

# The concept

In mathematics, we often use “structure-preserving” functions to compare objects.

**Definition.** (Homomorphism)

If  $\mathbb{A} = \langle A; F^{\mathbb{A}} \rangle$  and  $\mathbb{B} = \langle B; F^{\mathbb{B}} \rangle$  are algebras of the same signature, then a **homomorphism** from  $\mathbb{A}$  to  $\mathbb{B}$  is a (set-) function  $h: A \rightarrow B$ ,

# The concept

In mathematics, we often use “structure-preserving” functions to compare objects.

**Definition.** (Homomorphism)

If  $\mathbb{A} = \langle A; F^{\mathbb{A}} \rangle$  and  $\mathbb{B} = \langle B; F^{\mathbb{B}} \rangle$  are algebras of the same signature, then a **homomorphism** from  $\mathbb{A}$  to  $\mathbb{B}$  is a (set-) function  $h: A \rightarrow B$ , from the universe of  $\mathbb{A}$  to the universe of  $\mathbb{B}$  which **preserves the operations**

# The concept

In mathematics, we often use “structure-preserving” functions to compare objects.

**Definition.** (Homomorphism)

If  $\mathbb{A} = \langle A; F^{\mathbb{A}} \rangle$  and  $\mathbb{B} = \langle B; F^{\mathbb{B}} \rangle$  are algebras of the same signature, then a **homomorphism** from  $\mathbb{A}$  to  $\mathbb{B}$  is a (set-) function  $h: A \rightarrow B$ , from the universe of  $\mathbb{A}$  to the universe of  $\mathbb{B}$  which **preserves the operations** in the sense that for every  $n$ , every  $n$ -ary  $f \in F$ , and every  $a_1, \dots, a_n \in A$  we have

$$h(f^{\mathbb{A}}(a_1, \dots, a_n)) = f^{\mathbb{B}}(h(a_1), \dots, h(a_n)).$$

# The concept

In mathematics, we often use “structure-preserving” functions to compare objects.

**Definition.** (Homomorphism)

If  $\mathbb{A} = \langle A; F^{\mathbb{A}} \rangle$  and  $\mathbb{B} = \langle B; F^{\mathbb{B}} \rangle$  are algebras of the same signature, then a **homomorphism** from  $\mathbb{A}$  to  $\mathbb{B}$  is a (set-) function  $h: A \rightarrow B$ , from the universe of  $\mathbb{A}$  to the universe of  $\mathbb{B}$  which **preserves the operations** in the sense that for every  $n$ , every  $n$ -ary  $f \in F$ , and every  $a_1, \dots, a_n \in A$  we have

$$h(f^{\mathbb{A}}(a_1, \dots, a_n)) = f^{\mathbb{B}}(h(a_1), \dots, h(a_n)).$$

If our algebras are groups, this reduces to three conditions:

# The concept

In mathematics, we often use “structure-preserving” functions to compare objects.

**Definition.** (Homomorphism)

If  $\mathbb{A} = \langle A; F^{\mathbb{A}} \rangle$  and  $\mathbb{B} = \langle B; F^{\mathbb{B}} \rangle$  are algebras of the same signature, then a **homomorphism** from  $\mathbb{A}$  to  $\mathbb{B}$  is a (set-) function  $h: A \rightarrow B$ , from the universe of  $\mathbb{A}$  to the universe of  $\mathbb{B}$  which **preserves the operations** in the sense that for every  $n$ , every  $n$ -ary  $f \in F$ , and every  $a_1, \dots, a_n \in A$  we have

$$h(f^{\mathbb{A}}(a_1, \dots, a_n)) = f^{\mathbb{B}}(h(a_1), \dots, h(a_n)).$$

If our algebras are groups, this reduces to three conditions:

①  $h(1) = 1$

# The concept

In mathematics, we often use “structure-preserving” functions to compare objects.

**Definition.** (Homomorphism)

If  $\mathbb{A} = \langle A; F^{\mathbb{A}} \rangle$  and  $\mathbb{B} = \langle B; F^{\mathbb{B}} \rangle$  are algebras of the same signature, then a **homomorphism** from  $\mathbb{A}$  to  $\mathbb{B}$  is a (set-) function  $h: A \rightarrow B$ , from the universe of  $\mathbb{A}$  to the universe of  $\mathbb{B}$  which **preserves the operations** in the sense that for every  $n$ , every  $n$ -ary  $f \in F$ , and every  $a_1, \dots, a_n \in A$  we have

$$h(f^{\mathbb{A}}(a_1, \dots, a_n)) = f^{\mathbb{B}}(h(a_1), \dots, h(a_n)).$$

If our algebras are groups, this reduces to three conditions:

- 1  $h(1) = 1$
- 2  $h(x^{-1}) = h(x)^{-1}$

# The concept

In mathematics, we often use “structure-preserving” functions to compare objects.

**Definition.** (Homomorphism)

If  $\mathbb{A} = \langle A; F^{\mathbb{A}} \rangle$  and  $\mathbb{B} = \langle B; F^{\mathbb{B}} \rangle$  are algebras of the same signature, then a **homomorphism** from  $\mathbb{A}$  to  $\mathbb{B}$  is a (set-) function  $h: A \rightarrow B$ , from the universe of  $\mathbb{A}$  to the universe of  $\mathbb{B}$  which **preserves the operations** in the sense that for every  $n$ , every  $n$ -ary  $f \in F$ , and every  $a_1, \dots, a_n \in A$  we have

$$h(f^{\mathbb{A}}(a_1, \dots, a_n)) = f^{\mathbb{B}}(h(a_1), \dots, h(a_n)).$$

If our algebras are groups, this reduces to three conditions:

- 1  $h(1) = 1$
- 2  $h(x^{-1}) = h(x)^{-1}$
- 3  $h(x \cdot y) = h(x) \cdot h(y)$ .

# Some special properties of ALGEBRA homomorphisms

# Some special properties of ALGEBRA homomorphisms

Assume that  $h: \mathbb{A} \rightarrow \mathbb{B}$  is a homomorphism.

# Some special properties of ALGEBRA homomorphisms

Assume that  $h: \mathbb{A} \rightarrow \mathbb{B}$  is a homomorphism.

- 1 There is a unique algebra structure defined on the image of  $h$  for which  $h: \mathbb{A} \rightarrow \text{im}(h)$  is a homomorphism.

# Some special properties of ALGEBRA homomorphisms

Assume that  $h: \mathbb{A} \rightarrow \mathbb{B}$  is a homomorphism.

- 1 There is a unique algebra structure defined on the image of  $h$  for which  $h: \mathbb{A} \rightarrow \text{im}(h)$  is a homomorphism.
- 2 The function  $\iota: \text{im}(h) \rightarrow \mathbb{B}$  is a homomorphism.

# Some special properties of ALGEBRA homomorphisms

Assume that  $h: \mathbb{A} \rightarrow \mathbb{B}$  is a homomorphism.

- 1 There is a unique algebra structure defined on the image of  $h$  for which  $h: \mathbb{A} \rightarrow \text{im}(h)$  is a homomorphism.
- 2 The function  $\iota: \text{im}(h) \rightarrow \mathbb{B}$  is a homomorphism.
- 3 There is a unique algebra structure on the coimage of  $h$  that makes  $\nu: \mathbb{A} \rightarrow \text{coim}(h)$  a homomorphism.

# Some special properties of ALGEBRA homomorphisms

Assume that  $h: \mathbb{A} \rightarrow \mathbb{B}$  is a homomorphism.

- 1 There is a unique algebra structure defined on the image of  $h$  for which  $h: \mathbb{A} \rightarrow \text{im}(h)$  is a homomorphism.
- 2 The function  $\iota: \text{im}(h) \rightarrow \mathbb{B}$  is a homomorphism.
- 3 There is a unique algebra structure on the coimage of  $h$  that makes  $\nu: \mathbb{A} \rightarrow \text{coim}(h)$  a homomorphism.
- 4 If  $\text{im}(h)$  and  $\text{coim}(h)$  are given their appropriate algebra structures, then  $\bar{h}: \text{coim}(h) \rightarrow \text{im}(h)$  is an isomorphism.

# Some special properties of ALGEBRA homomorphisms

Assume that  $h: \mathbb{A} \rightarrow \mathbb{B}$  is a homomorphism.

- 1 There is a unique algebra structure defined on the image of  $h$  for which  $h: \mathbb{A} \rightarrow \text{im}(h)$  is a homomorphism.
- 2 The function  $\iota: \text{im}(h) \rightarrow \mathbb{B}$  is a homomorphism.
- 3 There is a unique algebra structure on the coimage of  $h$  that makes  $\nu: \mathbb{A} \rightarrow \text{coim}(h)$  a homomorphism.
- 4 If  $\text{im}(h)$  and  $\text{coim}(h)$  are given their appropriate algebra structures, then  $\bar{h}: \text{coim}(h) \rightarrow \text{im}(h)$  is an isomorphism.
- 5 The image of a subalgebra  $\mathbb{S} \leq \mathbb{A}$  under  $h$  is a subalgebra of  $\mathbb{B}$ .  
( $h(\mathbb{S}) \leq \mathbb{B}$ .)

# Some special properties of ALGEBRA homomorphisms

Assume that  $h: \mathbb{A} \rightarrow \mathbb{B}$  is a homomorphism.

- 1 There is a unique algebra structure defined on the image of  $h$  for which  $h: \mathbb{A} \rightarrow \text{im}(h)$  is a homomorphism.
- 2 The function  $\iota: \text{im}(h) \rightarrow \mathbb{B}$  is a homomorphism.
- 3 There is a unique algebra structure on the coimage of  $h$  that makes  $\nu: \mathbb{A} \rightarrow \text{coim}(h)$  a homomorphism.
- 4 If  $\text{im}(h)$  and  $\text{coim}(h)$  are given their appropriate algebra structures, then  $\bar{h}: \text{coim}(h) \rightarrow \text{im}(h)$  is an isomorphism.
- 5 The image of a subalgebra  $\mathbb{S} \leq \mathbb{A}$  under  $h$  is a subalgebra of  $\mathbb{B}$ .  
( $h(\mathbb{S}) \leq \mathbb{B}$ .)
- 6 The preimage of a subalgebra  $\mathbb{T} \leq \mathbb{B}$  under  $h$  is a subalgebra of  $\mathbb{A}$ .  
( $h^{-1}(\mathbb{T}) \leq \mathbb{A}$ .)

# Additional properties of GROUP homomorphisms

# Additional properties of GROUP homomorphisms

Assume that  $h: \mathbb{G} \rightarrow \mathbb{H}$  is a group homomorphism.

# Additional properties of GROUP homomorphisms

Assume that  $h: \mathbb{G} \rightarrow \mathbb{H}$  is a group homomorphism. Let  $K = h^{-1}(1_{\mathbb{B}})$  be the fiber over  $1_{\mathbb{B}}$ .

## Additional properties of GROUP homomorphisms

Assume that  $h: \mathbb{G} \rightarrow \mathbb{H}$  is a group homomorphism. Let  $K = h^{-1}(1_{\mathbb{B}})$  be the fiber over  $1_{\mathbb{B}}$ . I will call  $K$  the **Kernel** of  $h$  and write it  $\text{Ker}(h)$ .

# Additional properties of GROUP homomorphisms

Assume that  $h: \mathbb{G} \rightarrow \mathbb{H}$  is a group homomorphism. Let  $K = h^{-1}(1_{\mathbb{B}})$  be the fiber over  $1_{\mathbb{B}}$ . I will call  $K$  the **Kernel** of  $h$  and write it  $\text{Ker}(h)$ .  $\text{Ker}(h)$  is a single equivalence class of the equivalence relation

$$\text{ker}(h) = \{(x, y) \in A^2 \mid h(x) = h(y)\}.$$

That is,  $\text{Ker}(h) = 1_{\mathbb{A}}/\text{ker}(h)$ .

# Additional properties of GROUP homomorphisms

Assume that  $h: \mathbb{G} \rightarrow \mathbb{H}$  is a group homomorphism. Let  $K = h^{-1}(1_{\mathbb{B}})$  be the fiber over  $1_{\mathbb{B}}$ . I will call  $K$  the **Kernel** of  $h$  and write it  $\text{Ker}(h)$ .  $\text{Ker}(h)$  is a single equivalence class of the equivalence relation

$$\ker(h) = \{(x, y) \in A^2 \mid h(x) = h(y)\}.$$

That is,  $\text{Ker}(h) = 1_{\mathbb{A}}/\ker(h)$ .

- 1  $K$  is a subgroup of  $\mathbb{A}$ .

# Additional properties of GROUP homomorphisms

Assume that  $h: \mathbb{G} \rightarrow \mathbb{H}$  is a group homomorphism. Let  $K = h^{-1}(1_{\mathbb{B}})$  be the fiber over  $1_{\mathbb{B}}$ . I will call  $K$  the **Kernel** of  $h$  and write it  $\text{Ker}(h)$ .  $\text{Ker}(h)$  is a single equivalence class of the equivalence relation

$$\ker(h) = \{(x, y) \in A^2 \mid h(x) = h(y)\}.$$

That is,  $\text{Ker}(h) = 1_{\mathbb{A}}/\ker(h)$ .

- 1  $K$  is a subgroup of  $\mathbb{A}$ .

# Additional properties of GROUP homomorphisms

Assume that  $h: \mathbb{G} \rightarrow \mathbb{H}$  is a group homomorphism. Let  $K = h^{-1}(1_{\mathbb{B}})$  be the fiber over  $1_{\mathbb{B}}$ . I will call  $K$  the **Kernel** of  $h$  and write it  $\text{Ker}(h)$ .  $\text{Ker}(h)$  is a single equivalence class of the equivalence relation

$$\text{ker}(h) = \{(x, y) \in A^2 \mid h(x) = h(y)\}.$$

That is,  $\text{Ker}(h) = 1_{\mathbb{A}}/\text{ker}(h)$ .

- 1  $K$  is a subgroup of  $\mathbb{A}$ .
- 2 The fibers of  $h$  are the **left cosets** of  $K$ ,  $gK$ ,  $g \in G$ .

# Additional properties of GROUP homomorphisms

Assume that  $h: \mathbb{G} \rightarrow \mathbb{H}$  is a group homomorphism. Let  $K = h^{-1}(1_{\mathbb{B}})$  be the fiber over  $1_{\mathbb{B}}$ . I will call  $K$  the **Kernel** of  $h$  and write it  $\text{Ker}(h)$ .  $\text{Ker}(h)$  is a single equivalence class of the equivalence relation

$$\text{ker}(h) = \{(x, y) \in A^2 \mid h(x) = h(y)\}.$$

That is,  $\text{Ker}(h) = 1_{\mathbb{A}}/\text{ker}(h)$ .

- 1  $K$  is a subgroup of  $\mathbb{A}$ .
- 2 The fibers of  $h$  are the **left cosets** of  $K$ ,  $gK$ ,  $g \in G$ .

# Additional properties of GROUP homomorphisms

Assume that  $h: \mathbb{G} \rightarrow \mathbb{H}$  is a group homomorphism. Let  $K = h^{-1}(1_{\mathbb{B}})$  be the fiber over  $1_{\mathbb{B}}$ . I will call  $K$  the **Kernel** of  $h$  and write it  $\text{Ker}(h)$ .  $\text{Ker}(h)$  is a single equivalence class of the equivalence relation

$$\text{ker}(h) = \{(x, y) \in A^2 \mid h(x) = h(y)\}.$$

That is,  $\text{Ker}(h) = 1_{\mathbb{A}}/\text{ker}(h)$ .

- 1  $K$  is a subgroup of  $\mathbb{A}$ .
- 2 The fibers of  $h$  are the **left cosets** of  $K$ ,  $gK$ ,  $g \in G$ .
- 3 The fibers of  $h$  are also the **right cosets** of  $K$ ,  $Kg$ ,  $g \in G$ .

# Additional properties of GROUP homomorphisms

Assume that  $h: \mathbb{G} \rightarrow \mathbb{H}$  is a group homomorphism. Let  $K = h^{-1}(1_{\mathbb{B}})$  be the fiber over  $1_{\mathbb{B}}$ . I will call  $K$  the **Kernel** of  $h$  and write it  $\text{Ker}(h)$ .  $\text{Ker}(h)$  is a single equivalence class of the equivalence relation

$$\text{ker}(h) = \{(x, y) \in A^2 \mid h(x) = h(y)\}.$$

That is,  $\text{Ker}(h) = 1_{\mathbb{A}}/\text{ker}(h)$ .

- 1  $K$  is a subgroup of  $\mathbb{A}$ .
- 2 The fibers of  $h$  are the **left cosets** of  $K$ ,  $gK$ ,  $g \in G$ .
- 3 The fibers of  $h$  are also the **right cosets** of  $K$ ,  $Kg$ ,  $g \in G$ .

# Additional properties of GROUP homomorphisms

Assume that  $h: \mathbb{G} \rightarrow \mathbb{H}$  is a group homomorphism. Let  $K = h^{-1}(1_{\mathbb{B}})$  be the fiber over  $1_{\mathbb{B}}$ . I will call  $K$  the **Kernel** of  $h$  and write it  $\text{Ker}(h)$ .  $\text{Ker}(h)$  is a single equivalence class of the equivalence relation

$$\ker(h) = \{(x, y) \in A^2 \mid h(x) = h(y)\}.$$

That is,  $\text{Ker}(h) = 1_{\mathbb{A}}/\ker(h)$ .

- 1  $K$  is a subgroup of  $\mathbb{A}$ .
- 2 The fibers of  $h$  are the **left cosets** of  $K$ ,  $gK$ ,  $g \in G$ .
- 3 The fibers of  $h$  are also the **right cosets** of  $K$ ,  $Kg$ ,  $g \in G$ .
- 4 The coimage of  $h$  is the set of fibers of  $h$ ,  $G/K = \{gK \mid g \in G\}$ .

# Additional properties of GROUP homomorphisms

Assume that  $h: \mathbb{G} \rightarrow \mathbb{H}$  is a group homomorphism. Let  $K = h^{-1}(1_{\mathbb{B}})$  be the fiber over  $1_{\mathbb{B}}$ . I will call  $K$  the **Kernel** of  $h$  and write it  $\text{Ker}(h)$ .  $\text{Ker}(h)$  is a single equivalence class of the equivalence relation

$$\ker(h) = \{(x, y) \in A^2 \mid h(x) = h(y)\}.$$

That is,  $\text{Ker}(h) = 1_{\mathbb{A}}/\ker(h)$ .

- 1  $K$  is a subgroup of  $\mathbb{A}$ .
- 2 The fibers of  $h$  are the **left cosets** of  $K$ ,  $gK$ ,  $g \in G$ .
- 3 The fibers of  $h$  are also the **right cosets** of  $K$ ,  $Kg$ ,  $g \in G$ .
- 4 The coimage of  $h$  is the set of fibers of  $h$ ,  $G/K = \{gK \mid g \in G\}$ .

# Additional properties of GROUP homomorphisms

Assume that  $h: \mathbb{G} \rightarrow \mathbb{H}$  is a group homomorphism. Let  $K = h^{-1}(1_{\mathbb{B}})$  be the fiber over  $1_{\mathbb{B}}$ . I will call  $K$  the **Kernel** of  $h$  and write it  $\text{Ker}(h)$ .  $\text{Ker}(h)$  is a single equivalence class of the equivalence relation

$$\text{ker}(h) = \{(x, y) \in A^2 \mid h(x) = h(y)\}.$$

That is,  $\text{Ker}(h) = 1_{\mathbb{A}}/\text{ker}(h)$ .

- 1  $K$  is a subgroup of  $\mathbb{A}$ .
- 2 The fibers of  $h$  are the **left cosets** of  $K$ ,  $gK$ ,  $g \in G$ .
- 3 The fibers of  $h$  are also the **right cosets** of  $K$ ,  $Kg$ ,  $g \in G$ .
- 4 The coimage of  $h$  is the set of fibers of  $h$ ,  $G/K = \{gK \mid g \in G\}$ . The group structure on the coimage is:

# Additional properties of GROUP homomorphisms

Assume that  $h: \mathbb{G} \rightarrow \mathbb{H}$  is a group homomorphism. Let  $K = h^{-1}(1_{\mathbb{B}})$  be the fiber over  $1_{\mathbb{B}}$ . I will call  $K$  the **Kernel** of  $h$  and write it  $\text{Ker}(h)$ .  $\text{Ker}(h)$  is a single equivalence class of the equivalence relation

$$\text{ker}(h) = \{(x, y) \in A^2 \mid h(x) = h(y)\}.$$

That is,  $\text{Ker}(h) = 1_{\mathbb{A}}/\text{ker}(h)$ .

- 1  $K$  is a subgroup of  $\mathbb{A}$ .
- 2 The fibers of  $h$  are the **left cosets** of  $K$ ,  $gK$ ,  $g \in G$ .
- 3 The fibers of  $h$  are also the **right cosets** of  $K$ ,  $Kg$ ,  $g \in G$ .
- 4 The coimage of  $h$  is the set of fibers of  $h$ ,  $G/K = \{gK \mid g \in G\}$ . The group structure on the coimage is:
  - 1  $1_{G/K} = 1 \cdot K$ .

# Additional properties of GROUP homomorphisms

Assume that  $h: \mathbb{G} \rightarrow \mathbb{H}$  is a group homomorphism. Let  $K = h^{-1}(1_{\mathbb{B}})$  be the fiber over  $1_{\mathbb{B}}$ . I will call  $K$  the **Kernel** of  $h$  and write it  $\text{Ker}(h)$ .  $\text{Ker}(h)$  is a single equivalence class of the equivalence relation

$$\text{ker}(h) = \{(x, y) \in A^2 \mid h(x) = h(y)\}.$$

That is,  $\text{Ker}(h) = 1_{\mathbb{A}}/\text{ker}(h)$ .

- 1  $K$  is a subgroup of  $\mathbb{A}$ .
- 2 The fibers of  $h$  are the **left cosets** of  $K$ ,  $gK$ ,  $g \in G$ .
- 3 The fibers of  $h$  are also the **right cosets** of  $K$ ,  $Kg$ ,  $g \in G$ .
- 4 The coimage of  $h$  is the set of fibers of  $h$ ,  $G/K = \{gK \mid g \in G\}$ . The group structure on the coimage is:
  - 1  $1_{G/K} = 1 \cdot K$ .

# Additional properties of GROUP homomorphisms

Assume that  $h: \mathbb{G} \rightarrow \mathbb{H}$  is a group homomorphism. Let  $K = h^{-1}(1_{\mathbb{B}})$  be the fiber over  $1_{\mathbb{B}}$ . I will call  $K$  the **Kernel** of  $h$  and write it  $\text{Ker}(h)$ .  $\text{Ker}(h)$  is a single equivalence class of the equivalence relation

$$\text{ker}(h) = \{(x, y) \in A^2 \mid h(x) = h(y)\}.$$

That is,  $\text{Ker}(h) = 1_{\mathbb{A}}/\text{ker}(h)$ .

- ①  $K$  is a subgroup of  $\mathbb{A}$ .
- ② The fibers of  $h$  are the **left cosets** of  $K$ ,  $gK$ ,  $g \in G$ .
- ③ The fibers of  $h$  are also the **right cosets** of  $K$ ,  $Kg$ ,  $g \in G$ .
- ④ The coimage of  $h$  is the set of fibers of  $h$ ,  $G/K = \{gK \mid g \in G\}$ . The group structure on the coimage is:
  - ①  $1_{G/K} = 1 \cdot K$ .
  - ②  $(xK)^{-1} = x^{-1}K$ .

# Additional properties of GROUP homomorphisms

Assume that  $h: \mathbb{G} \rightarrow \mathbb{H}$  is a group homomorphism. Let  $K = h^{-1}(1_{\mathbb{B}})$  be the fiber over  $1_{\mathbb{B}}$ . I will call  $K$  the **Kernel** of  $h$  and write it  $\text{Ker}(h)$ .  $\text{Ker}(h)$  is a single equivalence class of the equivalence relation

$$\text{ker}(h) = \{(x, y) \in A^2 \mid h(x) = h(y)\}.$$

That is,  $\text{Ker}(h) = 1_{\mathbb{A}}/\text{ker}(h)$ .

- 1  $K$  is a subgroup of  $\mathbb{A}$ .
- 2 The fibers of  $h$  are the **left cosets** of  $K$ ,  $gK$ ,  $g \in G$ .
- 3 The fibers of  $h$  are also the **right cosets** of  $K$ ,  $Kg$ ,  $g \in G$ .
- 4 The coimage of  $h$  is the set of fibers of  $h$ ,  $G/K = \{gK \mid g \in G\}$ . The group structure on the coimage is:
  - 1  $1_{G/K} = 1 \cdot K$ .
  - 2  $(xK)^{-1} = x^{-1}K$ .

# Additional properties of GROUP homomorphisms

Assume that  $h: \mathbb{G} \rightarrow \mathbb{H}$  is a group homomorphism. Let  $K = h^{-1}(1_{\mathbb{B}})$  be the fiber over  $1_{\mathbb{B}}$ . I will call  $K$  the **Kernel** of  $h$  and write it  $\text{Ker}(h)$ .  $\text{Ker}(h)$  is a single equivalence class of the equivalence relation

$$\text{ker}(h) = \{(x, y) \in A^2 \mid h(x) = h(y)\}.$$

That is,  $\text{Ker}(h) = 1_{\mathbb{A}}/\text{ker}(h)$ .

- 1  $K$  is a subgroup of  $\mathbb{A}$ .
- 2 The fibers of  $h$  are the **left cosets** of  $K$ ,  $gK$ ,  $g \in G$ .
- 3 The fibers of  $h$  are also the **right cosets** of  $K$ ,  $Kg$ ,  $g \in G$ .
- 4 The coimage of  $h$  is the set of fibers of  $h$ ,  $G/K = \{gK \mid g \in G\}$ . The group structure on the coimage is:
  - 1  $1_{G/K} = 1 \cdot K$ .
  - 2  $(xK)^{-1} = x^{-1}K$ .
  - 3  $(xK)(yK) = (xy)K$ .