

The Paris-Harrington Theorem:

An Introduction to "Natural" Incompleteness

Everett Piper

`william.piper@colorado.edu`

Department of Mathematics
University of Colorado, Boulder (UCB)

Introduction

Theorem: (Gödel, 1931) There are true Π_1^0 sentences Ψ which are independent of the axioms of Peano Arithmetic (PA).

Introduction

Theorem: (Gödel, 1931) There are true Π_1^0 sentences Ψ which are independent of the axioms of Peano Arithmetic (PA).

- One such Ψ is the formal sentence expressing "PA is consistent".

Introduction

Theorem: (Gödel, 1931) There are true Π_1^0 sentences Ψ which are independent of the axioms of Peano Arithmetic (PA).

- One such Ψ is the formal sentence expressing "PA is consistent".
- Problem: Gödel's Ψ is arguably "unnatural";

Introduction

Theorem: (Gödel, 1931) There are true Π_1^0 sentences Ψ which are independent of the axioms of Peano Arithmetic (PA).

- One such Ψ is the formal sentence expressing "PA is consistent".
- Problem: Gödel's Ψ is arguably "unnatural";
Only refers to logicians' notions (provability and coding of syntax) so *metamathematical*.

Introduction

Theorem: (Gödel, 1931) There are true Π_1^0 sentences Ψ which are independent of the axioms of Peano Arithmetic (PA).

- One such Ψ is the formal sentence expressing "PA is consistent".
- Problem: Gödel's Ψ is arguably "unnatural";
Only refers to logicians' notions (provability and coding of syntax) so *metamathematical*.
No reference to classical or mainstream mathematical objects.

Introduction

Theorem: (Gödel, 1931) There are true Π_1^0 sentences Ψ which are independent of the axioms of Peano Arithmetic (PA).

- One such Ψ is the formal sentence expressing "PA is consistent".
- Problem: Gödel's Ψ is arguably "unnatural";
Only refers to logicians' notions (provability and coding of syntax) so *metamathematical*.
No reference to classical or mainstream mathematical objects.
- Motivates a search for independent sentences of a more "natural" flavor.

Introduction

Theorem: (Paris-Harrington, 1976) There is a "natural" Π_2^0 sentence Φ which is true in ω (the standard model), but also independent of the axioms of Peano Arithmetic.

Introduction

Theorem: (Paris-Harrington, 1976) There is a "natural" Π_2^0 sentence Φ which is true in ω (the standard model), but also independent of the axioms of Peano Arithmetic.

- Used a combination of ideas from large cardinal investigations....

Introduction

Theorem: (Paris-Harrington, 1976) There is a "natural" Π_2^0 sentence Φ which is true in ω (the standard model), but also independent of the axioms of Peano Arithmetic.

- Used a combination of ideas from large cardinal investigations....

Indiscernibles for various models (originated in research by Ehrenfeucht and Mostowski and also Jack Silver).

Introduction

Theorem: (Paris-Harrington, 1976) There is a "natural" Π_2^0 sentence Φ which is true in ω (the standard model), but also independent of the axioms of Peano Arithmetic.

- Used a combination of ideas from large cardinal investigations....

Indiscernibles for various models (originated in research by Ehrenfeucht and Mostowski and also Jack Silver).

Extensive use Ramsey's Theorem and the partition calculus.

Introduction

Theorem: (Paris-Harrington, 1976) There is a "natural" Π_2^0 sentence Φ which is true in ω (the standard model), but also independent of the axioms of Peano Arithmetic.

- Used a combination of ideas from large cardinal investigations....

Indiscernibles for various models (originated in research by Ehrenfeucht and Mostowski and also Jack Silver).

Extensive use Ramsey's Theorem and the partition calculus.

- Φ actually *equivalent* to 1-consistency of PA and transfinite induction through ϵ_0 .

Ramsey's Theorem

Theorem: (Ramsey, 1926) For any partition P of increasing e -tuples from ω into c colors, there exists some infinite $X \subseteq \omega$ such that P is constant on the increasing e -tuples from X .

Ramsey's Theorem

Theorem: (Ramsey, 1926) For any partition P of increasing e -tuples from ω into c colors, there exists some infinite $X \subseteq \omega$ such that P is constant on the increasing e -tuples from X .

- Such a set X is called *homogeneous* for the partition P .

Ramsey's Theorem

Theorem: (Ramsey, 1926) For any partition P of increasing e -tuples from ω into c colors, there exists some infinite $X \subseteq \omega$ such that P is constant on the increasing e -tuples from X .

- Such a set X is called *homogeneous* for the partition P .
- Useful notation:

$$\omega \rightarrow (\omega)_c^e$$

Ramsey's Theorem

Theorem: (Ramsey, 1926) For any partition P of increasing e -tuples from ω into c colors, there exists some infinite $X \subseteq \omega$ such that P is constant on the increasing e -tuples from X .

- Such a set X is called *homogeneous* for the partition P .
- Useful notation:

$$\omega \rightarrow (\omega)_c^e$$

- A finite ("miniature") version is provable using König's lemma on finitely branching, infinite trees:

$$\forall l, s, e, c \exists u ([l, u] \rightarrow (s)_c^e)$$

Basic Strategy

Start with a non-standard model $\mathcal{M} \models PA$ and candidate sentence Φ such that $\mathcal{M} \models \Phi$. Given a in \mathcal{M} ,

Basic Strategy

Start with a non-standard model $\mathcal{M} \models PA$ and candidate sentence Φ such that $\mathcal{M} \models \Phi$. Given a in \mathcal{M} ,

- Pick least b that satisfies Φ ;

Basic Strategy

Start with a non-standard model $\mathcal{M} \models PA$ and candidate sentence Φ such that $\mathcal{M} \models \Phi$. Given a in \mathcal{M} ,

- Pick least b that satisfies Φ ;
- Apply Ramsey's Theorem (or variant) to find large homogeneous subsets of \mathcal{M} ;

Basic Strategy

Start with a non-standard model $\mathcal{M} \models PA$ and candidate sentence Φ such that $\mathcal{M} \models \Phi$. Given a in \mathcal{M} ,

- Pick least b that satisfies Φ ;
- Apply Ramsey's Theorem (or variant) to find large homogeneous subsets of \mathcal{M} ;
- Use homogeneous sets \mathcal{H} as indiscernible elements for an initial segment \mathcal{I} of \mathcal{M} .

Basic Strategy

Start with a non-standard model $\mathcal{M} \models PA$ and candidate sentence Φ such that $\mathcal{M} \models \Phi$. Given a in \mathcal{M} ,

- Pick least b that satisfies Φ ;
- Apply Ramsey's Theorem (or variant) to find large homogeneous subsets of \mathcal{M} ;
- Use homogeneous sets \mathcal{H} as indiscernible elements for an initial segment \mathcal{I} of \mathcal{M} .
- If we can find \mathcal{H} between a and b , can get \mathcal{I} to contain a but not b (this will guarantee $\mathcal{I} \models \neg\Phi$).

Basic Strategy

Start with a non-standard model $\mathcal{M} \models PA$ and candidate sentence Φ such that $\mathcal{M} \models \Phi$. Given a in \mathcal{M} ,

- Pick least b that satisfies Φ ;
- Apply Ramsey's Theorem (or variant) to find large homogeneous subsets of \mathcal{M} ;
- Use homogeneous sets \mathcal{H} as indiscernible elements for an initial segment \mathcal{I} of \mathcal{M} .
- If we can't get \mathcal{H} between a and b , can get \mathcal{I} to contain a but not b (this will guarantee $\mathcal{I} \models \neg\Phi$).
- Show $\mathcal{I} \models PA$, and we're done!

Basic Strategy

Start with a non-standard model $\mathcal{M} \models PA$ and candidate sentence Φ such that $\mathcal{M} \models \Phi$. Given a in \mathcal{M} ,

- Pick least b that satisfies Φ ;
- Apply Ramsey's Theorem (or variant) to find large homogeneous subsets of \mathcal{M} ;
- Use homogeneous sets \mathcal{H} as indiscernible elements for an initial segment \mathcal{I} of \mathcal{M} .
- If we can't get \mathcal{H} between a and b , can get \mathcal{I} to contain a but not b (this will guarantee $\mathcal{I} \models \neg\Phi$).
- Show $\mathcal{I} \models PA$, and we're done!

Strategy suggests looking for Φ which are variants of Ramsey's Theorem.

Paris-Harrington Principle (PH)

Call a finite set $\mathcal{H} \subseteq \omega$ a *relatively large* set if $\min(\mathcal{H}) < |\mathcal{H}|$.

Paris-Harrington Principle (PH)

Call a finite set $\mathcal{H} \subseteq \omega$ a *relatively large* set if $\min(\mathcal{H}) < |\mathcal{H}|$.

- The set $\{3, 5, 26, 1767\}$ is relatively large.

Paris-Harrington Principle (PH)

Call a finite set $\mathcal{H} \subseteq \omega$ a *relatively large* set if $\min(\mathcal{H}) < |\mathcal{H}|$.

- The set $\{3, 5, 26, 1767\}$ is relatively large.
- The set $\{5, 26, 1767, 10^{10}\}$ is not relatively large.

Paris-Harrington Principle (PH)

Call a finite set $\mathcal{H} \subseteq \omega$ a *relatively large* set if $\min(\mathcal{H}) < |\mathcal{H}|$.

- The set $\{3, 5, 26, 1767\}$ is relatively large.
- The set $\{5, 26, 1767, 10^{10}\}$ is not relatively large.

Remark. Notions of largeness for finite sets of integers play prominent role in modern independence proofs.

Paris-Harrington Principle (PH)

Given natural numbers e, c, s, u we use the notation

$$[0, u] \rightarrow_* (s)_c^e$$

to mean that for every partition $P : [0, u]^e \rightarrow c$ there is a relatively large $H \subseteq [0, u]$ which is homogeneous for P of size at least s .

Paris-Harrington Principle (PH)

Given natural numbers e, c, s, u we use the notation

$$[0, u] \rightarrow_* (s)_c^e$$

to mean that for every partition $P : [0, u]^e \rightarrow c$ there is a relatively large $H \subseteq [0, u]$ which is homogeneous for P of size at least s .

● PH: $\forall e, c, s \exists u ([0, u] \rightarrow_* (s)_c^e)$ (Note this is Π_2^0).

Paris-Harrington Principle (PH)

Given natural numbers e, c, s, u we use the notation

$$[0, u] \rightarrow_* (s)_c^e$$

to mean that for every partition $P : [0, u]^e \rightarrow c$ there is a relatively large $H \subseteq [0, u]$ which is homogeneous for P of size at least s .

- PH: $\forall e, c, s \exists u ([0, u] \rightarrow_* (s)_c^e)$ (Note this is Π_2^0).
- Without the $*$ underneath the arrow, this is just the Finite Ramsey Theorem (which is provable in PA).

Paris-Harrington Principle (PH)

Given natural numbers e, c, s, u we use the notation

$$[0, u] \rightarrow_* (s)_c^e$$

to mean that for every partition $P : [0, u]^e \rightarrow c$ there is a relatively large $H \subseteq [0, u]$ which is homogeneous for P of size at least s .

- PH: $\forall e, c, s \exists u ([0, u] \rightarrow_* (s)_c^e)$ (Note this is Π_2^0).
- Without the $*$ underneath the arrow, this is just the Finite Ramsey Theorem (which is provable in PA).
- PH is a consequence of the Infinite Ramsey Theorem and the proof *cannot* be carried out in PA.

IRT implies PH

Theorem: The Infinite Ramsey Theorem implies the Paris-Harrington Principle.

IRT implies PH

Theorem: The Infinite Ramsey Theorem implies the Paris-Harrington Principle.

Proof. Fix e, c, s and suppose otherwise that no such interval $[0, u]$ exists satisfying PH. Call P a counterexample for $[0, u]$ if P partitions $[0, u]^e$ into c colors but no $H \subset [0, u]$ which is homogeneous for P is also relatively large.

IRT implies PH

Theorem: The Infinite Ramsey Theorem implies the Paris-Harrington Principle.

Proof. Fix e, c, s and suppose otherwise that no such interval $[0, u]$ exists satisfying PH. Call P a counterexample for $[0, u]$ if P partitions $[0, u]^e$ into c colors but no $H \subset [0, u]$ which is homogeneous for P is also relatively large.

Organize the collection of counterexamples into an infinite, finitely branching tree T . For P and P' counterexamples, put $P <_T P'$ if $u < u'$ and P is the restriction of P' to $[0, u]^e$.

IRT implies PH (cont'd)

Theorem: The Infinite Ramsey Theorem implies the Paris-Harrington Principle.

Proof. (cont'd)

IRT implies PH (cont'd)

Theorem: The Infinite Ramsey Theorem implies the Paris-Harrington Principle.

Proof. (cont'd)

Apply König's lemma to get a partition $P : [0, \omega)^e \rightarrow c$ where for every u , P restricted to $[0, u]^e$ is a counterexample.

IRT implies PH (cont'd)

Theorem: The Infinite Ramsey Theorem implies the Paris-Harrington Principle.

Proof. (cont'd)

Apply König's lemma to get a partition $P : [0, \omega)^e \rightarrow c$ where for every u , P restricted to $[0, u]^e$ is a counterexample.

By IRT there is an infinite $H \subseteq \omega$ homogeneous for P . But by choosing u large enough compared to s and $\min(H)$, we have $H \cap [0, u]$ is relatively large and homogeneous for P restricted to $[0, u]^e$, a contradiction. \diamond

IRT implies PH (cont'd)

Theorem: The Infinite Ramsey Theorem implies the Paris-Harrington Principle.

Proof. (cont'd)

Apply König's lemma to get a partition $P : [0, \omega)^e \rightarrow c$ where for every u , P restricted to $[0, u]^e$ is a counterexample.

By IRT there is an infinite $H \subseteq \omega$ homogeneous for P . But by choosing u large enough compared to s and $\min(H)$, we have $H \cap [0, u]$ is relatively large and homogeneous for P restricted to $[0, u]^e$, a contradiction. \diamond

- *Remark.* The proof indicates many “self-refining” variants of PH, i.e., there are homogeneous sets H with $\min(H) + s < |H|$, $(\min H)^2 < |H|$, etc.

IRT implies PH (cont'd)

Theorem: The Infinite Ramsey Theorem implies the Paris-Harrington Principle.

Proof. (cont'd)

Apply König's lemma to get a partition $P : [0, \omega)^e \rightarrow c$ where for every u , P restricted to $[0, u]^e$ is a counterexample.

By IRT there is an infinite $H \subseteq \omega$ homogeneous for P . But by choosing u large enough compared to s and $\min(H)$, we have $H \cap [0, u]$ is relatively large and homogeneous for P restricted to $[0, u]^e$, a contradiction. \diamond

- *Remark.* The proof indicates many “self-refining” variants of PH, i.e., there are homogeneous sets H with $\min(H) + s < |H|$, $(\min H)^2 < |H|$, etc.

Original Proof (Outline)

Main ingredients of Paris' original proof:

Original Proof (Outline)

Main ingredients of Paris' original proof:

- Combinatorics of Partitions.

Original Proof (Outline)

Main ingredients of Paris' original proof:

- Combinatorics of Partitions.
- PH implies \clubsuit .

Original Proof (Outline)

Main ingredients of Paris' original proof:

- Combinatorics of Partitions.
- PH implies \clubsuit .
- \clubsuit implies a certain theory T is consistent:

Original Proof (Outline)

Main ingredients of Paris' original proof:

- Combinatorics of Partitions.
- PH implies ♣.
- ♣ implies a certain theory T is consistent:
 1. Given finite $T_0 \subseteq T$, use ♣ to find a collection of indiscernibles and build a model of T_0 .

Original Proof (Outline)

Main ingredients of Paris' original proof:

- Combinatorics of Partitions.
- PH implies ♣.
- ♣ implies a certain theory T is consistent:
 1. Given finite $T_0 \subseteq T$, use ♣ to find a collection of indiscernibles and build a model of T_0 .
 2. Apply Compactness Theorem.

Original Proof (Outline)

Main ingredients of Paris' original proof:

- Combinatorics of Partitions.
- PH implies \clubsuit .
- \clubsuit implies a certain theory T is consistent:
 1. Given finite $T_0 \subseteq T$, use \clubsuit to find a collection of indiscernibles and build a model of T_0 .
 2. Apply Compactness Theorem.
- PA proves $Con(T) \rightarrow Con(PA)$.

Original Proof (Outline)

Main ingredients of Paris' original proof:

- Combinatorics of Partitions.
- PH implies \clubsuit .
- \clubsuit implies a certain theory T is consistent:
 1. Given finite $T_0 \subseteq T$, use \clubsuit to find a collection of indiscernibles and build a model of T_0 .
 2. Apply Compactness Theorem.
- PA proves $Con(T) \rightarrow Con(PA)$.
- *Remark.* Nowadays, combinatorics of original proof is bypassed by showing PH implies KM (another combinatoric principle) and then showing independence of KM from PA.

Some Details

A (long) series of lemmas on partition combinatorics:

Some Details

A (long) series of lemmas on partition combinatorics:

Lemma. Given $P_1 : [0, u]^e \rightarrow c_1$ and $P_2 : [0, u]^e \rightarrow c_2$, there is $P : [0, u]^e \rightarrow c_1 c_2$ where any $H \subseteq [0, u]$ is homogeneous for P iff also homogeneous for both P_1 and P_2 .

Some Details

A (long) series of lemmas on partition combinatorics:

Lemma. Given $P_1 : [0, u]^e \rightarrow c_1$ and $P_2 : [0, u]^e \rightarrow c_2$, there is $P : [0, u]^e \rightarrow c_1 c_2$ where any $H \subseteq [0, u]$ is homogeneous for P iff also homogeneous for both P_1 and P_2 .

Lemma. Given $P : [0, u]^e \rightarrow c$ there is $P' : [0, u]^{e+1} \rightarrow (1 + 2\sqrt{c})$ where any $H \subseteq [0, u]$ of size $> e + 1$ is homogeneous for P iff also homogeneous for P' .

More Combinatorics

Lemma. For every c there is $P : [0, u]^1 \rightarrow c + 1$ where any H homogeneous for P of size at least 2 guarantees $e \leq \min(H)$.

More Combinatorics

Lemma. For every c there is $P : [0, u]^1 \rightarrow c + 1$ where any H homogeneous for P of size at least 2 guarantees $e \leq \min(H)$.

Lemma. Given $P : [0, u]^e \rightarrow c$, any $H \subseteq [0, u]$ is homogeneous for P iff every subset of H of size $e + 1$ is homogeneous for P .

More Combinatorics

Lemma. For every c there is $P : [0, u]^1 \rightarrow c + 1$ where any H homogeneous for P of size at least 2 guarantees $e \leq \min(H)$.

Lemma. Given $P : [0, u]^e \rightarrow c$, any $H \subseteq [0, u]$ is homogeneous for P iff every subset of H of size $e + 1$ is homogeneous for P .

Lemma. For every c there is $P : [0, u]^1 \rightarrow c + 1$ where any H homogeneous for P of size at least 2 guarantees $e \leq \min(H)$.

More Combinatorics

Lemma. For every c there is $P : [0, u]^1 \rightarrow c + 1$ where any H homogeneous for P of size at least 2 guarantees $e \leq \min(H)$.

Lemma. Given $P : [0, u]^e \rightarrow c$, any $H \subseteq [0, u]$ is homogeneous for P iff every subset of H of size $e + 1$ is homogeneous for P .

Lemma. For every c there is $P : [0, u]^1 \rightarrow c + 1$ where any H homogeneous for P of size at least 2 guarantees $e \leq \min(H)$.

Corollary. Given a finite number of partitions P_1, \dots, P_i , can "glue" them all together into a single partition P where a set H of appropriate size is homogeneous for P iff also homogeneous for each P_i .

More Combinatorics

Lemma. For every c there is $P : [0, u]^1 \rightarrow c + 1$ where any H homogeneous for P of size at least 2 guarantees $e \leq \min(H)$.

Lemma. Given $P : [0, u]^e \rightarrow c$, any $H \subseteq [0, u]$ is homogeneous for P iff every subset of H of size $e + 1$ is homogeneous for P .

Lemma. For every c there is $P : [0, u]^1 \rightarrow c + 1$ where any H homogeneous for P of size at least 2 guarantees $e \leq \min(H)$.

Corollary. Given a finite number of partitions P_1, \dots, P_i , can "glue" them all together into a single partition P where a set H of appropriate size is homogeneous for P iff also homogeneous for each P_i .

Fast-Growing Functions

Definition: $f_0(x) = x + 2$, $f_{n+1}(x) = f_n^{[x]}(2)$ where the exponent $[x]$ on f denotes composition with itself x times.

Fast-Growing Functions

Definition: $f_0(x) = x + 2$, $f_{n+1}(x) = f_n^{[x]}(2)$ where the exponent $[x]$ on f denotes composition with itself x times.

Lemma. For each m there is $P : [0, u]^2 \rightarrow c$ (where c depends only on m) such that any H homogeneous for P and relatively large has the property that $x, y \in H$ where $x < y$ implies $f_m(x) < y$.

Fast-Growing Functions

Definition: $f_0(x) = x + 2$, $f_{n+1}(x) = f_n^{[x]}(2)$ where the exponent $[x]$ on f denotes composition with itself x times.

Lemma. For each m there is $P : [0, u]^2 \rightarrow c$ (where c depends only on m) such that any H homogeneous for P and relatively large has the property that $x, y \in H$ where $x < y$ implies $f_m(x) < y$.

Lemma. For any given m , $P : [0, u]^e \rightarrow c$ with $2 \leq e$ there is $P' : [0, u]^e \rightarrow c'$ (where c' depends only on m, e , and c) such that if there is a relatively large H of size $> e$ homogeneous for P' , there is an H' homogeneous for P with $\max(e + 1, f_m(\min(H'))) \leq |H'|$.

Fast-Growing Functions

Definition: $f_0(x) = x + 2$, $f_{n+1}(x) = f_n^{[x]}(2)$ where the exponent $[x]$ on f denotes composition with itself x times.

Lemma. For each m there is $P : [0, u]^2 \rightarrow c$ (where c depends only on m) such that any H homogeneous for P and relatively large has the property that $x, y \in H$ where $x < y$ implies $f_m(x) < y$.

Lemma. For any given m , $P : [0, u]^e \rightarrow c$ with $2 \leq e$ there is $P' : [0, u]^e \rightarrow c'$ (where c' depends only on m, e , and c) such that if there is a relatively large H of size $> e$ homogeneous for P' , there is an H' homogeneous for P with $\max(e + 1, f_m(\min(H'))) \leq |H'|$.

- *Corollary:* Relatively large homogeneous sets can be sufficiently “spread out”. (This is what we want!)

Even More Combinatorics!

Lemma. ♣: For all e, s, c there is a u such that for any family $\{P_\alpha : \alpha < 2^u\}$ of partitions $P_\alpha : [0, u]^e \rightarrow c$ there is an X of size at least s such that

Even More Combinatorics!

Lemma. ♣: For all e, s, c there is a u such that for any family $\{P_\alpha : \alpha < 2^u\}$ of partitions $P_\alpha : [0, u]^e \rightarrow c$ there is an X of size at least s such that

1. $a, b \in X$ and $a < b$ implies $a^2 < b$.

Even More Combinatorics!

Lemma. ♣: For all e, s, c there is a u such that for any family $\{P_\alpha : \alpha < 2^u\}$ of partitions $P_\alpha : [0, u]^e \rightarrow c$ there is an X of size at least s such that

1. $a, b \in X$ and $a < b$ implies $a^2 < b$.
2. $a \in X$ and $\alpha < 2^a$ implies $X - (a + 1)$ is homogeneous for P_α .

Even More Combinatorics!

Lemma. ♣: For all e, s, c there is a u such that for any family $\{P_\alpha : \alpha < 2^u\}$ of partitions $P_\alpha : [0, u]^e \rightarrow c$ there is an X of size at least s such that

1. $a, b \in X$ and $a < b$ implies $a^2 < b$.
2. $a \in X$ and $\alpha < 2^a$ implies $X - (a + 1)$ is homogeneous for P_α .

Lemma. PH implies ♣.

The Theory T

Let c_1, c_2, \dots be a countable sequence of new constant symbols and add them to the language of PA. Define a theory T in this new language with the following axioms:

The Theory T

Let c_1, c_2, \dots be a countable sequence of new constant symbols and add them to the language of PA. Define a theory T in this new language with the following axioms:

- (i) Usual recursive axioms for $+$, \times , $<$ and induction axioms for formulas with bounded quantifiers.

The Theory T

Let c_1, c_2, \dots be a countable sequence of new constant symbols and add them to the language of PA. Define a theory T in this new language with the following axioms:

- (i) Usual recursive axioms for $+$, \times , $<$ and induction axioms for formulas with bounded quantifiers.
- (ii) The axiom $c_i^2 < c_{i+1}$ for each $i \in \omega$.

The Theory T

Let c_1, c_2, \dots be a countable sequence of new constant symbols and add them to the language of PA. Define a theory T in this new language with the following axioms:

- (i) Usual recursive axioms for $+$, \times , $<$ and induction axioms for formulas with bounded quantifiers.
- (ii) The axiom $c_i^2 < c_{i+1}$ for each $i \in \omega$.
- (iii) For each finite subset $\bar{i} = i_1, \dots, i_r \subseteq \omega$, let $c(\bar{i}) = c_{i_1}, \dots, c_{i_r}$. For each $j < \min(\bar{k}, \bar{k}')$ and each Δ_0 formula $\psi(\bar{y}; \bar{z})$ where \bar{k}, \bar{k}' and \bar{z} all have the same length, we have the axiom

$$\forall \bar{y} < j [\psi(\bar{y}; c(\bar{k})) \leftrightarrow \psi(\bar{y}; c(\bar{k}'))].$$

The Theory T

Let c_1, c_2, \dots be a countable sequence of new constant symbols and add them to the language of PA. Define a theory T in this new language with the following axioms:

- (i) Usual recursive axioms for $+$, \times , $<$ and induction axioms for formulas with bounded quantifiers.
- (ii) The axiom $c_i^2 < c_{i+1}$ for each $i \in \omega$.
- (iii) For each finite subset $\bar{i} = i_1, \dots, i_r \subseteq \omega$, let $c(\bar{i}) = c_{i_1}, \dots, c_{i_r}$. For each $j < \min(\bar{k}, \bar{k}')$ and each Δ_0 formula $\psi(\bar{y}; \bar{z})$ where \bar{k}, \bar{k}' and \bar{z} all have the same length, we have the axiom

$$\forall \bar{y} < j [\psi(\bar{y}; c(\bar{k})) \leftrightarrow \psi(\bar{y}; c(\bar{k}'))].$$

● **Note:** Indiscernibility condition is built into T .

The Theory T

Let c_1, c_2, \dots be a countable sequence of new constant symbols and add them to the language of PA. Define a theory T in this new language with the following axioms:

- (i) Usual recursive axioms for $+$, \times , $<$ and induction axioms for formulas with bounded quantifiers.
- (ii) The axiom $c_i^2 < c_{i+1}$ for each $i \in \omega$.
- (iii) For each finite subset $\bar{i} = i_1, \dots, i_r \subseteq \omega$, let $c(\bar{i}) = c_{i_1}, \dots, c_{i_r}$. For each $j < \min(\bar{k}, \bar{k}')$ and each Δ_0 formula $\psi(\bar{y}; \bar{z})$ where \bar{k}, \bar{k}' and \bar{z} all have the same length, we have the axiom

$$\forall \bar{y} < j [\psi(\bar{y}; c(\bar{k})) \leftrightarrow \psi(\bar{y}; c(\bar{k}'))].$$

● **Note:** Indiscernibility condition is built into T .

$Con(T)$ **implies** $Con(PA)$

$Con(T)$ **implies** $Con(PA)$

Proof. Let $\mathcal{A} \models T$ and let \mathcal{I} be the initial segment of \mathcal{A} of those elements $a < c_i$ for some $i \in \omega$. \mathcal{I} is closed under $+$ and \times . So it suffices to show that $\mathcal{I} \models PA$.

$Con(T)$ **implies** $Con(PA)$

Proof. Let $\mathcal{A} \models T$ and let \mathcal{I} be the initial segment of \mathcal{A} of those elements $a < c_i$ for some $i \in \omega$. \mathcal{I} is closed under $+$ and \times . So it suffices to show that $\mathcal{I} \models PA$.

Given a formula $\theta(\bar{y})$ in prenex form (in the language of PA), say $\theta(\bar{y}) = \exists x_1 \dots \forall x_r \varphi(\bar{x}, \bar{y})$ define another formula $\theta^*(\bar{y}; \bar{z}) = \exists x_1 < z_1 \dots \forall x_r < z_r \varphi(\bar{x}, \bar{y}; \bar{z})$.

$Con(T)$ implies $Con(PA)$

Proof. Let $\mathcal{A} \models T$ and let \mathcal{I} be the initial segment of \mathcal{A} of those elements $a < c_i$ for some $i \in \omega$. \mathcal{I} is closed under $+$ and \times . So it suffices to show that $\mathcal{I} \models PA$.

Given a formula $\theta(\bar{y})$ in prenex form (in the language of PA), say $\theta(\bar{y}) = \exists x_1 \dots \forall x_r \varphi(\bar{x}, \bar{y})$ define another formula $\theta^*(\bar{y}; \bar{z}) = \exists x_1 < z_1 \dots \forall x_r < z_r \varphi(\bar{x}, \bar{y}; \bar{z})$.

Now, given $\theta(\bar{y})$, $c_{i_1} < \dots < c_{i_k}$ of same length as \bar{y} , and any tuple \bar{a} , each member of which is $< c_{i_1}$, show

$$\mathcal{I} \models \exists x_1 \dots \forall x_r \varphi(\bar{x}, \bar{a}) \Leftrightarrow \mathcal{A} \models \exists x_1 < c_{i_1} \dots \forall x_r < c_{i_k} \varphi(\bar{x}, \bar{a}).$$

$Con(T)$ implies $Con(PA)$

Proof. Let $\mathcal{A} \models T$ and let \mathcal{I} be the initial segment of \mathcal{A} of those elements $a < c_i$ for some $i \in \omega$. \mathcal{I} is closed under $+$ and \times . So it suffices to show that $\mathcal{I} \models PA$.

Given a formula $\theta(\bar{y})$ in prenex form (in the language of PA), say $\theta(\bar{y}) = \exists x_1 \dots \forall x_r \varphi(\bar{x}, \bar{y})$ define another formula $\theta^*(\bar{y}; \bar{z}) = \exists x_1 < z_1 \dots \forall x_r < z_r \varphi(\bar{x}, \bar{y}; \bar{z})$.

Now, given $\theta(\bar{y})$, $c_{i_1} < \dots < c_{i_k}$ of same length as \bar{y} , and any tuple \bar{a} , each member of which is $< c_{i_1}$, show

$$\mathcal{I} \models \exists x_1 \dots \forall x_r \varphi(\bar{x}, \bar{a}) \Leftrightarrow \mathcal{A} \models \exists x_1 < c_{i_1} \dots \forall x_r < c_{i_k} \varphi(\bar{x}, \bar{a}).$$

Follows by induction on θ , using the third set of axioms for T . \diamond

$Con(T)$ implies $Con(PA)$

Proof. Let $\mathcal{A} \models T$ and let \mathcal{I} be the initial segment of \mathcal{A} of those elements $a < c_i$ for some $i \in \omega$. \mathcal{I} is closed under $+$ and \times . So it suffices to show that $\mathcal{I} \models PA$.

Given a formula $\theta(\bar{y})$ in prenex form (in the language of PA), say $\theta(\bar{y}) = \exists x_1 \dots \forall x_r \varphi(\bar{x}, \bar{y})$ define another formula $\theta^*(\bar{y}; \bar{z}) = \exists x_1 < z_1 \dots \forall x_r < z_r \varphi(\bar{x}, \bar{y}; \bar{z})$.

Now, given $\theta(\bar{y})$, $c_{i_1} < \dots < c_{i_k}$ of same length as \bar{y} , and any tuple \bar{a} , each member of which is $< c_{i_1}$, show

$$\mathcal{I} \models \exists x_1 \dots \forall x_r \varphi(\bar{x}, \bar{a}) \Leftrightarrow \mathcal{A} \models \exists x_1 < c_{i_1} \dots \forall x_r < c_{i_k} \varphi(\bar{x}, \bar{a}).$$

Follows by induction on θ , using the third set of axioms for T . \diamond

Remark. Above proof can be formalized and carried out in PA.

Model-Theoretic Proof of PH

Theorem:(Bovykin) The statement “for all e, s, c there exists u such that for every $P : [0, u]^e \rightarrow c$ there is $H \subseteq [0, u]$ homogeneous for P with $\max(s, e \cdot (2^{e \cdot \min(H)} + 1)) < |H|$ ” is not a theorem of PA.

Model-Theoretic Proof of PH

Theorem:(Bovykin) The statement “for all e, s, c there exists u such that for every $P : [0, u]^e \rightarrow c$ there is $H \subseteq [0, u]$ homogeneous for P with $\max(s, e \cdot (2^{e \cdot \min(H)} + 1)) < |H|$ ” is not a theorem of PA.

Proof. WLOG, let $c = 2$, $\mathcal{M} \models PA$ be non-standard, pick d a non-standard integer. Let

$\varphi_1(x_1, \dots, x_d, y), \dots, \varphi_d(x_1, \dots, x_d, y)$ be an enumeration of the first d Δ_0 formulas in at most the free variables shown.

Model-Theoretic Proof of PH

Theorem:(Bovykin) The statement “for all e, s, c there exists u such that for every $P : [0, u]^e \rightarrow c$ there is $H \subseteq [0, u]$ homogeneous for P with $\max(s, e \cdot (2^{e \cdot \min(H)} + 1)) < |H|$ ” is not a theorem of PA.

Proof. WLOG, let $c = 2$, $\mathcal{M} \models PA$ be non-standard, pick d a non-standard integer. Let

$\varphi_1(x_1, \dots, x_d, y), \dots, \varphi_d(x_1, \dots, x_d, y)$ be an enumeration of the first d Δ_0 formulas in at most the free variables shown.

Let u be the minimal point in \mathcal{M} such that for every

$P : [0, u]^{2d+1} \rightarrow 2$ there is a P -homogeneous $H \subseteq [0, u]$ with $|H| > d \cdot (2^{d \cdot \min(H)} + 1)$.

Model-Theoretic Proof of PH

Theorem:(Bovykin) The statement “for all e, s, c there exists u such that for every $P : [0, u]^e \rightarrow c$ there is $H \subseteq [0, u]$ homogeneous for P with $\max(s, e \cdot (2^{e \cdot \min(H)} + 1)) < |H|$ ” is not a theorem of PA.

Proof. WLOG, let $c = 2$, $\mathcal{M} \models PA$ be non-standard, pick d a non-standard integer. Let

$\varphi_1(x_1, \dots, x_d, y), \dots, \varphi_d(x_1, \dots, x_d, y)$ be an enumeration of the first d Δ_0 formulas in at most the free variables shown.

Let u be the minimal point in \mathcal{M} such that for every

$P : [0, u]^{2d+1} \rightarrow 2$ there is a P -homogeneous $H \subseteq [0, u]$ with $|H| > d \cdot (2^{d \cdot \min(H)} + 1)$.

Note that exponent e in this case is really $2d + 1$.

Model-Theoretic Proof of PH

Model-Theoretic Proof of PH

Proof. cont'd. Define our coloring P' by setting
 $P'(a, b_1, \dots, b_d, c_1, \dots, c_d) = 0$ if for all $x < a$,

$$\{i \leq d : \varphi_i(b_1, \dots, b_d, x)\} = \{i \leq d : \varphi_i(c_1, \dots, c_d, x)\}$$

and 1 otherwise.

Model-Theoretic Proof of PH

Proof. cont'd. Define our coloring P' by setting $P'(a, b_1, \dots, b_d, c_1, \dots, c_d) = 0$ if for all $x < a$,

$$\{i \leq d : \varphi_i(b_1, \dots, b_d, x)\} = \{i \leq d : \varphi_i(c_1, \dots, c_d, x)\}$$

and 1 otherwise.

Using the definition of u , extract a P' -homogeneous set $H \subseteq [0, u]$ of size greater than $d \cdot (2^{d \cdot \min(H)} + 1)$.

Model-Theoretic Proof of PH

Proof. cont'd. For every increasing d -tuple $b_1 < \dots < b_d$ in $H - \{\min(H)\}$ define the following sequence of d -many subsets of $[0, \min(H))$:

$$\langle \{x < \min(H) : \varphi_1(b_1, \dots, b_d, x)\}, \dots, \dots, \{x < \min(H) : \varphi_d(b_1, \dots, b_d, x)\} \rangle.$$

Model-Theoretic Proof of PH

Proof. cont'd. For every increasing d -tuple $b_1 < \dots < b_d$ in $H - \{\min(H)\}$ define the following sequence of d -many subsets of $[0, \min(H))$:

$$\langle \{x < \min(H) : \varphi_1(b_1, \dots, b_d, x)\}, \dots, \dots, \{x < \min(H) : \varphi_d(b_1, \dots, b_d, x)\} \rangle.$$

There are at most $2^{d \cdot \min(H)}$ such sequences, so by Pigeonhole Principle, there are $b_1 < b_2 < \dots < b_d < c_1 < \dots < c_d$ in $H - \{\min(H)\}$ such that

$$P'(\min(H), b_1, \dots, b_d, c_1, \dots, c_d) = 0.$$

Model-Theoretic Proof of PH

Proof. cont'd. For every increasing d -tuple $b_1 < \dots < b_d$ in $H - \{\min(H)\}$ define the following sequence of d -many subsets of $[0, \min(H))$:

$$\langle \{x < \min(H) : \varphi_1(b_1, \dots, b_d, x)\}, \dots, \dots, \{x < \min(H) : \varphi_d(b_1, \dots, b_d, x)\} \rangle.$$

There are at most $2^{d \cdot \min(H)}$ such sequences, so by Pigeonhole Principle, there are $b_1 < b_2 < \dots < b_d < c_1 < \dots < c_d$ in $H - \{\min(H)\}$ such that

$$P'(\min(H), b_1, \dots, b_d, c_1, \dots, c_d) = 0.$$

By homogeneity of H , P' must be constantly 0 on $[H]^{2d+1}$.

Model-Theoretic Proof of PH

Proof. cont'd. For every increasing d -tuple $b_1 < \dots < b_d$ in $H - \{\min(H)\}$ define the following sequence of d -many subsets of $[0, \min(H))$:

$$\langle \{x < \min(H) : \varphi_1(b_1, \dots, b_d, x)\}, \dots, \dots, \{x < \min(H) : \varphi_d(b_1, \dots, b_d, x)\} \rangle.$$

There are at most $2^{d \cdot \min(H)}$ such sequences, so by Pigeonhole Principle, there are $b_1 < b_2 < \dots < b_d < c_1 < \dots < c_d$ in $H - \{\min(H)\}$ such that

$$P'(\min(H), b_1, \dots, b_d, c_1, \dots, c_d) = 0.$$

By homogeneity of H , P' must be constantly 0 on $[H]^{2d+1}$.

Model-Theoretic Proof of PH

Proof. cont'd. Let $\bar{H} = \{h_1, \dots, h_d\}$ denote the last d elements of H . Then for any $a < b_1 < \dots < b_d < c_1 < \dots < c_d$ in $H - \bar{H}$ we have for every $x < a$:

$$\begin{aligned} \{i \leq d : \varphi_i(b_1, \dots, b_d, x)\} &= \{i \leq d : \varphi_i(h_1, \dots, h_d, x)\} \\ &= \{i \leq d : \varphi_i(c_1, \dots, c_d, x)\}. \end{aligned}$$

Model-Theoretic Proof of PH

Proof. cont'd. Let $\bar{H} = \{h_1, \dots, h_d\}$ denote the last d elements of H . Then for any $a < b_1 < \dots < b_d < c_1 < \dots < c_d$ in $H - \bar{H}$ we have for every $x < a$:

$$\begin{aligned} \{i \leq d : \varphi_i(b_1, \dots, b_d, x)\} &= \{i \leq d : \varphi_i(h_1, \dots, h_d, x)\} \\ &= \{i \leq d : \varphi_i(c_1, \dots, c_d, x)\}. \end{aligned}$$

This is the indiscernibility condition we need.

Model-Theoretic Proof of PH

Proof. cont'd. Let $\bar{H} = \{h_1, \dots, h_d\}$ denote the last d elements of H . Then for any $a < b_1 < \dots < b_d < c_1 < \dots < c_d$ in $H - \bar{H}$ we have for every $x < a$:

$$\begin{aligned} \{i \leq d : \varphi_i(b_1, \dots, b_d, x)\} &= \{i \leq d : \varphi_i(h_1, \dots, h_d, x)\} \\ &= \{i \leq d : \varphi_i(c_1, \dots, c_d, x)\}. \end{aligned}$$

This is the indiscernibility condition we need.

Let \mathcal{I} contain all points of \mathcal{M} below some element in the first ω -many elements of H .

Model-Theoretic Proof of PH

Proof. cont'd. Let $\bar{H} = \{h_1, \dots, h_d\}$ denote the last d elements of H . Then for any $a < b_1 < \dots < b_d < c_1 < \dots < c_d$ in $H - \bar{H}$ we have for every $x < a$:

$$\begin{aligned} \{i \leq d : \varphi_i(b_1, \dots, b_d, x)\} &= \{i \leq d : \varphi_i(h_1, \dots, h_d, x)\} \\ &= \{i \leq d : \varphi_i(c_1, \dots, c_d, x)\}. \end{aligned}$$

This is the indiscernibility condition we need.

Let \mathcal{I} contain all points of \mathcal{M} below some element in the first ω -many elements of H .

Then $\mathcal{I} \models PA$ and also satisfies the negation of our statement. \diamond

Model-Theoretic Proof of PH

Proof. cont'd. Let $\bar{H} = \{h_1, \dots, h_d\}$ denote the last d elements of H . Then for any $a < b_1 < \dots < b_d < c_1 < \dots < c_d$ in $H - \bar{H}$ we have for every $x < a$:

$$\begin{aligned} \{i \leq d : \varphi_i(b_1, \dots, b_d, x)\} &= \{i \leq d : \varphi_i(h_1, \dots, h_d, x)\} \\ &= \{i \leq d : \varphi_i(c_1, \dots, c_d, x)\}. \end{aligned}$$

This is the indiscernibility condition we need.

Let \mathcal{I} contain all points of \mathcal{M} below some element in the first ω -many elements of H .

Then $\mathcal{I} \models PA$ and also satisfies the negation of our statement. \diamond

End

Thank You