

Math 2135 - Assignment 5

Due October 4, 2024

Problems 1-5 are review material for the first midterm on October 2. Solve them before Wednesday!

(1) Let

$$A = \begin{bmatrix} 0 & 3 & 1 & 2 \\ 1 & 4 & 0 & 7 \\ 2 & -1 & -3 & 8 \end{bmatrix}, b = \begin{bmatrix} 6 \\ 5 \\ -8 \end{bmatrix}$$

(a) Give the solution for $Ax = b$ in parametrized vector form.

(b) Give vectors that span the null space of A .

(2) Let $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a linear transformation with

$$T\left(\begin{bmatrix} 1 \\ 2 \end{bmatrix}\right) = \begin{bmatrix} 2 \\ -1 \\ 1 \end{bmatrix} \text{ and } T\left(\begin{bmatrix} 3 \\ 4 \end{bmatrix}\right) = \begin{bmatrix} 0 \\ 1 \\ -2 \end{bmatrix}.$$

What is the standard matrix of T ?

(3) Let $T: \mathbb{R}^n \rightarrow \mathbb{R}^n, x \mapsto Ax$, be a surjective linear map. Show that T is injective as well.

(4) True or false? Explain your answer.

(a) If $Ax = b$ is inconsistent for some vector b , then A cannot have a pivot in every column.

(b) If vectors $\mathbf{v}_1, \mathbf{v}_2$ are linearly independent and \mathbf{v}_3 is not in the span of $\mathbf{v}_1, \mathbf{v}_2$, then $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$ is linear independent.

(c) The range of $T: \mathbb{R}^n \rightarrow \mathbb{R}^m, x \mapsto Ax$, is the span of the columns of A .

(5) (a) Give examples of square matrices A, B such that neither A nor B is 0 (the matrix with all entries 0) but $AB = 0$.

(b) If the first two columns of a matrix B are equal, what can you say about the columns of AB ?

(c) We can view vectors in \mathbb{R}^n as $n \times 1$ matrices. For $\mathbf{u} = \begin{bmatrix} 2 \\ -1 \\ 3 \end{bmatrix}, \mathbf{v} = \begin{bmatrix} 0 \\ 2 \\ 1 \end{bmatrix}$

compute $\mathbf{u}^T \cdot \mathbf{v}$ and $\mathbf{u} \cdot \mathbf{v}^T$. Interpret the results.

(6) Prove for $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ with $ad - bc \neq 0$ that

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

(7) Are the following invertible? Give the inverse if possible.

$$A = \begin{bmatrix} 2 & 1 \\ 4 & -9 \end{bmatrix}, \quad B = \begin{bmatrix} 2 & -3 \\ 4 & -6 \end{bmatrix}, \quad C = \begin{bmatrix} 0 & 1 & 3 \\ 0 & 0 & 1 \\ 0 & -1 & -1 \end{bmatrix}$$

(8) A **diagonal matrix** A has all entries 0 except on the diagonal, that is,

$$A = \begin{bmatrix} a_{11} & 0 & \dots & 0 \\ 0 & a_{22} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & a_{nn} \end{bmatrix}.$$

Under which conditions is A invertible and what is A^{-1} ?