

Reminder

- Handouts are due Friday.
- Check WebAssign for online homework.
- Written homework is due Thursday.
- Syllabus last page sign and return by Friday.

Daily Quiz

- Go to [Socrative.com](https://www.socrative.com) and complete the quiz.
- Use your full name.
- Room Name: HONG5824

5.6 Integration by Parts - Your choice matters

Integrating by parts Find $\int x \sin x \, dx$.

Recall that last time, we found that $\int x \sin x \, dx = -x \cos x + \sin x + C$.

What if we choose a different u and dv ?

Yesterday, we chose $u = x$ and $dv = \sin x \, dx$.

Today, let's try $u = \sin x$ and $dv = x \, dx$.

$$\begin{array}{l|l} u = \sin x & v = \frac{x^2}{2} \\ \hline du = \cos x \, dx & dv = x \, dx \end{array}$$

$$\int u \, dv = uv - \int v \, du$$

$$\int x \sin x \, dx = \sin x \frac{x^2}{2} - \int \frac{x^2}{2} \cos x \, dx$$

$$= \sin x \frac{x^2}{2} - \frac{1}{2} \int x^2 \cos x \, dx$$

The new integrand $x^2 \cos x$ has a higher degree of x than we started with. This is no good because if we integrate by parts again on $x^2 \cos x$, we'll get $\int x^3 \sin x \, dx$

observation: choice of u and dv matters a lot!

5.5 How to choose your U and dV : LIATE

Integration by parts requires us to know the derivative of u and the anti-derivative of dv . Since some functions are harder to integrate than others, we let dv be a function that is **easier to integrate** while we let u be the function that is **harder to integrate**.

Following this heuristic, we have a rule that helps us pick the right u and we let the remainder be dv : **LIATE**.


When choosing u , follow the below priority list.

1. **L**ogarithmic functions (e.g. $\log x$)
2. **I**nverse Trig functions (e.g. $\arctan x$)
3. **A**lgebraic functions (e.g. $x^2, \frac{1}{x^7}$)
4. **T**rig functions (e.g. $\tan x$)
5. **E**xponential functions (e.g. $2^x, e^x$)

5.6 Integration by Parts

Guidelines for selecting u and dv

LIATE



Choose u to be the function that comes first in this list.

Evaluate $\int \ln x \, dx$.

$$\begin{array}{l|l} u = \ln x & v = x \\ \hline du = \frac{1}{x} dx & dv = dx \end{array}$$

$$\begin{aligned} uv - \int v du &= x \ln x - \int x \frac{1}{x} dx \\ &= x \ln x - x + C \end{aligned}$$

5.6 Integration by Parts

exponential

LIATE

algebraic

Integrating by parts twice

Find $\int \underbrace{t^2}_{\text{algebraic}} \underbrace{e^t}_{\text{exponential}} dt.$

$$\begin{array}{l|l} u = t^2 & v = e^t \\ \hline du = 2t dt & dv = e^t dt \end{array}$$

$$\begin{aligned} uv - \int v du &= t^2 e^t - \int e^t 2t dt \\ &= t^2 e^t - 2 \underbrace{\int t e^t dt}_I \end{aligned}$$

$I = \int \underbrace{t}_{\text{algebraic}} \underbrace{e^t}_{\text{exponential}} dt$ (Integ by parts again) LIATE

$$\begin{array}{l|l} u = t & v = e^t \\ \hline du = dt & dv = e^t dt \end{array}$$

$$\begin{aligned} uv - \int v du &= t e^t - \int e^t dt \\ &= t e^t - e^t + C = I \end{aligned}$$

$$\int t^2 e^t dt = t^2 e^t - 2[t e^t - e^t + C]$$

What happens if you don't follow LIATE?

Integrating by parts twice Find $\int t^2 e^t dt$. ^{algebraic}

Suppose we used LIATE the first time but not the second time.

$$\frac{u = t^2 \quad | \quad v = e^t}{du = 2t dt \quad | \quad dv = e^t dt} \quad \int t^2 e^t dt = t^2 e^t - \int e^t 2t dt = t^2 e^t - 2 \int t e^t dt$$

$I = \int t e^t dt$ We are supposed to pick $u = t$ and $dv = e^t dt$ following LIATE. What if we forgot? Let's try $u = e^t$ and $dv = t dt$.

$$\frac{u = e^t \quad | \quad v = \frac{t^2}{2}}{du = e^t dt \quad | \quad dv = t dt} \quad \int t e^t dt = e^t \frac{t^2}{2} - \int \frac{t^2}{2} e^t dt = \frac{t^2 e^t}{2} - \frac{1}{2} \int t^2 e^t dt$$

Observe that we now have t^2 instead of t . This is bad because if we try to simplify, we get back to where we started, making no progress.

$$t^2 e^t - 2I = t^2 e^t - 2 \left[\frac{t^2 e^t}{2} - \frac{1}{2} \int t^2 e^t dt \right] = t^2 e^t - t^2 e^t + \int t^2 e^t dt = 0 + \int t^2 e^t dt$$

Conclusion:
Always follow LIATE

5.6 Integration by Parts (Substitution before By-Parts)

$$\int \cos \sqrt{x} \, dx$$

Simple substitution
 $w = \sqrt{x}$
 $dw = \frac{1}{2} x^{-\frac{1}{2}} dx$

$$\sqrt{x} \, dw = \frac{1}{2} dx$$
$$2w \, dw = dx$$

$$\int \cos(w) (2w \, dw)$$
$$= 2 \int w \cos(w) \, dw$$

Now use integration by parts
on the above integral.

LI(A)TE

w is algebraic

$u = w$	$v = \sin(w)$
$du = dw$	$dv = \cos(w) \, dw$

$$\int w \cos(w) = uv - \int v \, du$$
$$= w \sin(w) - \int \sin(w) \, dw$$
$$= w \sin(w) + \cos(w) + C$$

Hence $\int \cos \sqrt{x} \, dx$

$$= 2 \int w \cos(w) \, dw$$
$$= 2 [w \sin(w) + \cos(w) + C]$$
$$= 2 [\sqrt{x} \sin(\sqrt{x}) + \cos(\sqrt{x}) + C]$$

5.6 Integration by Parts (Definite Integrals)

$$\int_a^b f(x)g'(x) dx = f(x)g(x)\Big|_a^b - \int_a^b g(x)f'(x) dx$$

5.6 Integration by Parts

$$\int_0^1 \underbrace{\arctan(x)}_{\text{Inverse Trig}} dx$$

L(I)ATE

$u = \arctan x$	$v = x$
$du = \frac{1}{1+x^2} dx$	$dv = dx$

$$u \Big|_0^1 - \int_0^1 v du = x \arctan x \Big|_0^1 - \underbrace{\int_0^1 x \frac{1}{1+x^2} dx}_I$$

$$I = \int_0^1 \frac{x}{1+x^2} dx \quad (\text{simple } u\text{-sub})$$

$$u = 1+x^2$$

$$u(0) = 1+0 = 1$$

$$du = 2x dx$$

$$u(1) = 1+1^2 = 2$$

$$\begin{aligned} \int_0^1 \frac{x}{1+x^2} dx &= \int_{u(0)}^{u(1)} \frac{du/2}{u} = \frac{1}{2} \int_1^2 \frac{du}{u} \\ &= \frac{1}{2} \ln|u| \Big|_1^2 = \frac{1}{2} [\ln(2) - \ln(1)] \end{aligned}$$

$$\int_0^1 \arctan(x) dx = x \arctan x \Big|_0^1 - I$$

$$= [\arctan 1 - 0] - \frac{1}{2} \ln(2)$$

$$= \frac{\pi}{4} - \frac{\ln(2)}{2}$$