

### Exercise 13.47

#### Abstract Algebra 1

#### MATH 3140

SEBASTIAN CASALAINA

ABSTRACT. This is Exercise 13.47 from Fraleigh [Fra03, §13]:

**Exercise 13.47.** Show that any group homomorphism  $\phi : G \rightarrow G'$  where  $|G|$  is a prime must either be the trivial homomorphism or a one-to-one map.

*Solution.* Let  $\phi : G \rightarrow G'$  be a group homomorphism where  $|G|$  is a prime. Let  $e' \in G'$  be the identity element. The problem asks us to show that  $\phi(g) = e'$  for all  $g \in G$ , or that  $\phi$  is one-to-one.

To prove this, let us consider  $\ker \phi$ . The kernel of a homomorphism is a subgroup of  $G$ , and, since  $|G|$  is finite,  $|\ker \phi|$  divides  $|G|$  (Theorem of Lagrange [Fra03, p.100]). By virtue of the fact that  $|G|$  is prime, it follows that either  $|\ker \phi| = 1$  or  $|\ker \phi| = |G|$ . That is, either  $\ker \phi = \{e\}$ , where  $e$  is the identity element of  $G$ , or  $\ker \phi = G$ .

In the former case (i.e.,  $\ker \phi = \{e\}$ ),  $\phi$  is one-to-one (a homomorphism is one-to-one if and only if the kernel is trivial [Fra03, Corollary 13.18, p.131]). In the latter case,  $\phi(g) = e'$  for all  $g \in G$ , from the definition of the kernel. □

## REFERENCES

[Fra03] John Fraleigh, *A First Course in Abstract Algebra*, Seventh edition, Addison Wesley, Pearson, 2003.

UNIVERSITY OF COLORADO, DEPARTMENT OF MATHEMATICS, CAMPUS BOX 395, BOULDER, CO 80309

*Email address:* casa@math.colorado.edu