

### Exercise 6.4.23

#### Linear Algebra MATH 2130

SEBASTIAN CASALAINA

ABSTRACT. This is Exercise 6.4.23 from Lay [LLM21, §6.4]:

**Exercise 6.4.23.** Suppose  $A = QR$ , where  $Q$  is  $m \times n$  and  $R$  is  $n \times n$ . Show that if the columns of  $A$  are linearly independent, then  $R$  must be invertible.

*Solution.* The columns of  $A$  are linearly independent if and only if  $\ker(A) = \{\mathbf{0}\}$ . It follows that  $\ker(R) = \{\mathbf{0}\}$ , since if  $\mathbf{x} \in \ker(R)$ , then  $A\mathbf{x} = QR\mathbf{x} = Q\mathbf{0} = \mathbf{0}$ , so that  $\mathbf{x} \in \ker(A) = \{\mathbf{0}\}$ . A square matrix is invertible if and only if its kernel is trivial; therefore  $R$  is invertible.  $\square$

## REFERENCES

[LLM21] David Lay, Stephen Lay, and Judi McDonald, *Linear Algebra and its Applications*, Sixth edition, Pearson, 2021.

UNIVERSITY OF COLORADO, DEPARTMENT OF MATHEMATICS, CAMPUS BOX 395, BOULDER, CO 80309

*Email address:* `casa@math.colorado.edu`