

EXAM 1: SOME PRACTICE PROBLEMS

1. Prove the following.

Proposition.

- (a) If $a, m \in \mathbb{Z}$ and $m|a$, then $m|(an)$ for any $n \in \mathbb{Z}$.
(b) If $a, b, m \in \mathbb{Z}$, and $m|a$ and $m|b$, then $m|(a + b)$ and $m|(a - b)$.

(Use the definition of “ $|$ ” from your fact sheet.)

2. Use both parts of the previous problem to show that, if $m, a, b, x, y \in \mathbb{Z}$ and $m|a$ and $m|b$, then $m|(ax + by)$.

3. Prove the following.

Proposition. If $a, b, c, d, m \in \mathbb{Z}$, $m|(a - b)$, and $m|(c - d)$, then $m|(a + c - (b + d))$. (Use the definition of “ $|$ ” from your fact sheet.)

4. Prove the following.

Proposition. If $a, b \in \mathbb{Z}$, a is odd, and b is even, then ab is even. (Use the definitions of even and odd integers: an even integer is one of the form $2k$ for some $k \in \mathbb{Z}$; an odd integer is one of the form $2k + 1$ for some $k \in \mathbb{Z}$.)

5. Use proof by contrapositive to prove the following.

Proposition. If n^2 is odd, then n is odd. (Use the definitions of even and odd integers: an even integer is one of the form $2k$ for some $k \in \mathbb{Z}$; an odd integer is one of the form $2k + 1$ for some $k \in \mathbb{Z}$.)

6. Use proof by contrapositive to prove the following.

Proposition. Let $a, b \in \mathbb{Z}$. If ab is a multiple of 3, then either a or b is a multiple of 3. (Hint: an integer that is *not* a multiple of 3 is of the form $3k + r$ where $k \in \mathbb{Z}$ and r equals either 1 or 2.)

7. Prove that, if $m \in \mathbb{Z}$ is even, then $4|m^2$, and if m is odd, then $4|(m^2 - 1)$.

8. Prove by contradiction that, if $a, b \in \mathbb{Z}$ are odd, then there does not exist $c \in \mathbb{Z}$ with $a^2 + b^2 = c^2$. Hint: assume $a, b \in \mathbb{Z}$ are odd and that there does exist such an integer c . Show that $4|(c^2 - 2)$. Use this together with the previous exercise to derive a contradiction.

9. Identify each of the following statements as true or false (circle “**T**” or “**F**”). **Please explain your answers:** If a statement is true, explain why (you don’t need to provide a complete proof; just a sentence or two will do). If a statement is false, provide a counterexample to the statement, and explain why it’s a counterexample.

(a) $\forall m \in \mathbb{Z}, \exists n \in \mathbb{Z}: m|n.$

T **F**

(b) $\exists n \in \mathbb{Z}: \forall m \in \mathbb{Z}, m|n.$

T **F**

(c) $\exists n \in \mathbb{Z}: \forall m \in \mathbb{Z}, n m.$	T	F
(d) $\forall m \in \mathbb{Z}, \forall n \in \mathbb{Z}, \exists k \in \mathbb{Z}: (m - n) k.$	T	F
(e) $\exists k \in \mathbb{Z}: \forall m \in \mathbb{Z}, \forall n \in \mathbb{Z}, (m - n) k.$	T	F
(f) $\exists k \in \mathbb{Z}: \forall m \in \mathbb{Z}, \forall n \in \mathbb{Z}, k (m - n).$	T	F
(g) $\exists n \in \mathbb{Z}: \forall m \in \mathbb{Z}, \forall k \in \mathbb{Z}, k (m - n).$	T	F
(h) $\sim(\forall m \in \mathbb{Z}, \exists k \in \mathbb{Z}: \forall n \in \mathbb{Z}, (m - n) k).$	T	F

10. Let F_n be the n th Fibonacci number, defined by

$$F_1 = F_2 = 1, \quad F_{n+2} = F_{n+1} + F_n \quad (n \geq 1).$$

Use mathematical induction to prove that, for any $n \in \mathbb{N}$, the sum of the first n Fibonacci numbers with *even* index equals $F_{2n+1} - 1$. That is: prove that, $\forall n \in \mathbb{N}$,

$$F_2 + F_4 + F_6 + \cdots + F_n = F_{2n+1} - 1.$$

11. Use the principle of mathematical induction to prove that, for any $n \in \mathbb{N}$,

$$1 \cdot 1! + 2 \cdot 2! + 3 \cdot 3! + \cdots + n \cdot n! = (n + 1)! - 1.$$

(Hint: $(k + 2)(k + 1)! = (k + 2)!$.) Please clearly identify your base step, induction hypothesis, inductive step, and the conclusion of your proof.

12. Use the principle of mathematical induction to prove that, for any $n \in \mathbb{N}$, the sum of the first n *odd* positive integers is n^2 . In other words prove that, $\forall n \in \mathbb{N}$,

$$1 + 3 + 5 + 7 + \cdots + 2n - 1 = n^2.$$

13. Use the principle of mathematical induction to prove that, for any $n \in \mathbb{N}$,

$$1 \cdot 3 + 2 \cdot 4 + 3 \cdot 5 + 4 \cdot 6 + \cdots + n(n + 2) = \frac{n(n + 1)(2n + 7)}{6}.$$

14. Use mathematical induction to prove that, for any $n \in \mathbb{Z}_{\geq 0} = \{0, 1, 2, 3, \dots\}$,

$$5|(n^5 - n).$$

15. Form the negation of each of the following statements. Your negated statement should not contain the symbol “ $\sim$ ” (or any symbol that means “not” or “negation”) in it anywhere. (Don’t worry about whether these statements are true or false.) Remark: the negation of $a < b$ is $a \geq b$.

- (a) $\forall z \in \mathbb{R}, |25 - 37| < z.$
- (b) $\exists y \in \mathbb{R}: \forall z \in \mathbb{R}, |25 - y| < z.$
- (c) $\sim(\forall x \in \mathbb{R}, \exists y \in \mathbb{R}: \forall z \in \mathbb{R}: |x - y| < z).$

(d) $\forall x \in \mathbb{R}, \sim(\exists y \in \mathbb{R} : \sim(\forall z \in \mathbb{R}, |x - y| < z))$.

16. Identify each of the following statements as true or false, by putting a “T” or “F” in the space to the *left* of the statement. Then, in the space to the *right* of the statement, put the *number* of the statement that is the *negation* of the statement in question. For example, if the negation of statement 2 is statement 7, then put a “7” in the space to the right of statement 2.

One of the statements has no negation present, so leave the space to the right of that statement blank.

(Recall that $\mathbb{R}^+$ denotes the set of positive real numbers.)

- | | | | |
|----|-------|---|-------|
| 1. | _____ | $\forall w \in \mathbb{R}^+, \forall x \in \mathbb{R}, \forall y \in \mathbb{R}, \forall z \in \mathbb{R}^+, w + z \leq x - y$ | _____ |
| 2. | _____ | $\exists w \in \mathbb{R}^+, \exists x \in \mathbb{R}, \forall y \in \mathbb{R}, \forall z \in \mathbb{R}^+, w + z \leq x - y$ | _____ |
| 3. | _____ | $\exists w \in \mathbb{R}^+, \exists x \in \mathbb{R}, \exists y \in \mathbb{R}, \exists z \in \mathbb{R}^+, x < w + y + z$ | _____ |
| 4. | _____ | $\sim(\sim(\forall w \in \mathbb{R}^+, \exists x \in \mathbb{R}, \forall y \in \mathbb{R}, \exists z \in \mathbb{R}^+, x - y < w + z))$ | _____ |
| 5. | _____ | $\exists w \in \mathbb{R}^+, \forall x \in \mathbb{R}, \exists y \in \mathbb{R}, \forall z \in \mathbb{R}^+, w + z \leq x - y$ | _____ |
| 6. | _____ | $\forall w \in \mathbb{R}^+, \exists x \in \mathbb{R}, \forall y \in \mathbb{R}, \exists z \in \mathbb{R}^+, w + z \leq x - y$ | _____ |
| 7. | _____ | $\forall w \in \mathbb{R}^+, \forall x \in \mathbb{R}, \sim(\forall y \in \mathbb{R}, \forall z \in \mathbb{R}^+, w + z \leq x - y)$ | _____ |
| 8. | _____ | $\sim(\forall w \in \mathbb{R}^+, \forall x \in \mathbb{R}, \sim(\forall y \in \mathbb{R}, \forall z \in \mathbb{R}^+, x - y < w + z))$ | _____ |
| 9. | _____ | $\forall w \in \mathbb{R}^+, \forall x \in \mathbb{R}, \exists y \in \mathbb{R}, \exists z \in \mathbb{R}^+, w + z \leq x - y$ | _____ |

17. (a) Explain, intuitively, why the negation of the statement $P \Rightarrow Q$ is the statement $P \wedge \sim Q$ (meaning “ P and not Q ”).

(b) Negate the statement

$$\forall \varepsilon > 0, \exists \delta > 0: 0 < |x - a| < \delta \Rightarrow |f(x) - L| < \varepsilon.$$

18. Prove that $5^{2n} - 1$ is a multiple of 8 for all $n \in \mathbb{N}$.

19. What’s wrong with the following proof?


Theorem. All sneakers are identical.

Proof. Let $A(n)$ be the statement that all sneakers in any set of n sneakers are identical.

Is $A(1)$ true? Yes, any one sneaker is identical to itself.

Now assume $A(k)$: any k sneakers are identical to each other. To deduce $A(k+1)$, line all $k+1$ sneakers up in a row. The first k sneakers in that row are identical, by the induction

hypothesis. So are the last k , by the induction hypothesis. But the second sneaker in the row belongs to both the first k and the last k , so all sneakers are identical to the second one.



So $A(k+1)$ follows. So $A(k) \Rightarrow A(k+1)$. So by induction, $A(n)$ is true for all $n \in \mathbb{N}$.

In particular, let n be the total number of sneakers in existence. Then all of these are identical. $\square$