

CLS Solutions to Selected Exercises: Individual Homework Assignment #1

Assignment. Part 3: A Measles Epidemic (pages 23–24): Exercises 14–20. **Part 4: Other Diseases** (page 24): Exercises 21ab, 22.

Exercise 14.

One stays infected for 14 days. This is because, as discussed in Section 1.2, the recovery coefficient b equals the reciprocal of the number of days k that one stays infected. Here we have $b = 1/14 \text{ day}^{-1}$, so $k = 14$ days.

Exercise 15.

$$S_T = b/a = (1/14)/0.00001 = 7,142.9 \text{ persons.}$$

Exercise 16.

$$S(1) \approx 4,4446.6, \quad I(1) \approx 2,903.4, \quad R(1) \approx 2,650.0; \text{ all units are persons.}$$

Exercise 17.

$$S(2) \approx 43,156.1, \quad I(2) \approx 3,986.5, \quad R(2) \approx 2,857.4; \text{ all units are persons.}$$

Exercise 18.

$$S(3) \approx 41,435.7, \quad I(3) \approx 5,422.1, \quad R(3) \approx 3,142.1; \text{ all units are persons.}$$

Exercise 19.

$$S(2) \approx 43,494.2, \quad I(2) \approx 3,706.8, \quad R(2) \approx 2800.0; \text{ all units are persons.}$$

Exercise 20.

(a) The new transmission coefficient is half the old one. So the new coefficient is $a = 0.5 \times 0.00001 = 0.000005$.

(b) $S_T = b/a = (1/14)/(0.000005) \approx 14,286$.

Another way to see this is: by cutting the transmission coefficient in half, we double the threshold value S_T of S . (Think about why this makes intuitive sense.)

- (c) At the outset of the disease, we have

$$\begin{aligned} I'(0) &= aS(0)I(0) - bI(0) = I(0)(aS(0) - b) \\ &= (2,100)(0.000005 \times 45400 - 1/14) = 326.7. \end{aligned}$$

That is, I' is initially positive, so I is initially increasing, so the quarantine does *not* eliminate the epidemic.

Exercise 21.

- (a) Since $b = 0.08 \text{ day}^{-1}$, we have that the infection lasts for $k = 1/0.08 = 12.5$ days.
- (b) We need I' to be positive; that is, $aSI - bI > 0$. Factoring gives $I(aS - b) > 0$. Since $I > 0$, this means we need $aS - b > 0$, or $aS > b$, or $S > b/a = 0.08/0.00002 = 4,000.0$. So there must be more than 4,000 susceptible individuals for the illness to take hold.

Exercise 22.

- (a) Since the illness lasts for 4 days, we know that our recovery coefficient, b , should be $b = \frac{1}{4 \text{ days}} = 0.25 \frac{1}{\text{days}}$. Since a typical susceptible person meets only about 0.3% of infection population each day, we know that $p = 0.3\% = .003$. Finally, since the infection is transmitted in only one contact out of 6, we know that $q = \frac{1}{6} = 0.167$. Hence our SIR model for this measles-like disease looks like:

$$\begin{aligned} S' &= -aSI = -qpSI = -(0.167)(.003)SI = -0.0005SI \\ I' &= aSI - bI = qpSI - bI = (0.167)(.003) - 0.25I = 0.0005SI - 0.25I \\ R' &= bI = 0.25I. \end{aligned}$$

- (b) We want to know: at what point does I start decreasing – that is, at what point does I' become negative? But

$$\begin{aligned} I' &= 0.0005SI - 0.25I \\ &= I(0.0005S - 0.25). \end{aligned}$$

Since $I \geq 0$ always, I' can only be negative if $0.0005S - 0.25 < 0$, or $0.0005S < 0.25$, or $S < 0.25/0.0005 = 500$. In sum: if $S < 500$, then $I' < 0$, and hence the illness will fade away without becoming an epidemic.