

Modeling a disease using SIR.

## I) Initial set-up.

$S$ : # of susceptibles  
 $I$ : # of infected  
 $R$ : # of recovered

$S'$ : rate of change of  $S$   
 $I'$ : rate of change of  $I$   
 $R'$ : rate of change of  $R$

These are all variables: they vary with time  $t$ .

## Assumptions:

- Everyone infected recovers eventually.
- The duration of infection is the same for everyone.
- Once recovered, you're immune and can't infect.
- Only a fraction of contacts with disease cause infection.
- The units are:
  - days for time  $t$ ;
  - individuals for  $S$ ,  $I$ , and  $R$ ;
  - individuals/day for  $S'$ ,  $I'$ , and  $R'$ .

II) Thinking about the rates of change  $S'$ ,  $I'$ , and  $R'$ .(a)  $R'$ .

Say the disease lasts  $k$  days. Then each day, on average, the number recovered will increase by  $1/k$  times the size of the infected population. So

$$R' = \frac{1}{k} I = bI \quad \text{where } b = 1/k.$$

Note:  $b$  is constant (it doesn't change with  $t$ ). We say  $R'$  is proportional to  $I$ .

(b)  $S'$ .

Suppose:

(i) Each susceptible has contact with a fraction, call it  $p$ , of the infected population on a given day. Since the number of possible S-to-I contacts on a given day is  $S \cdot I$ , this means the number of actual S-to-I contacts on a given day is  $p \cdot S \cdot I$ .

(ii) A fraction, call it  $q$ , of the actual contacts lead to infection.

Together, (i) and (ii) mean  $q \cdot p \cdot S \cdot I$  new infections each day, meaning  $S$  decreases by  $q \cdot p \cdot S \cdot I$  each day.  $\therefore$

$$S' = -qpSI = -aSI \quad \text{where } a = qp.$$

(The minus sign reflects the decrease in  $S$ .)

(c) Assuming the total population  $S+I+R$  stays constant, the changes in  $S, I$ , and  $R$  must cancel, meaning  $S' + I' + R' = 0$ , so

$$I' = -S' - R'$$

or, by (a) and (b) above,

$$I' = aSI - bI$$

III)

SUMMARY

Under the assumptions described above, we have

$$\begin{aligned} S' &= -aSI \\ I' &= aSI - bI \\ R' &= bI \end{aligned}$$

SIR equations

Here:

•  $b (>0)$  is the recovery coefficient. Units:  $1/\text{day}$ , or  $\text{day}^{-1}$

•  $a (>0)$  is the transmission coefficient. Units:  $1/(\text{person} \cdot \text{day})$ , or  $(\text{person} \cdot \text{day})^{-1}$

→  $a$  and  $b$  are called parameters.

Question: so what?

Answer: prediction. More on this soon.