

CLS Monomers, Dimers, Trimers, and Rates of Change (SOLUTIONS)

A beaker contains three types of molecules, called monomers, dimers, and trimers. We use M , D , and T to stand for the quantities of each of the three respective types. Suppose these quantities are changing over time, according to the following “rate equations:”

$$M' = -4M^2 - 0.8MD,$$

$$D' = 2M^2 - 0.8MD,$$

$$T' = 0.8MD.$$

Let's suppose that, *initially*, there are equal (nonzero) quantities of monomers and dimers.

1. Is D initially increasing or decreasing? Please explain.

Initially, we have $D = M$, by the above note. So initially, by the above equation for D' , we have

$$\begin{aligned} D' &= 2M^2 - 0.8MD \\ &= 2M^2 - 0.8M \cdot M \\ &= 2M^2 - 0.8M^2 \\ &= (2 - 0.8)M^2 = 1.2M^2 > 0. \end{aligned}$$

Since D' is initially positive, we see that D is initially increasing.

2. At a certain point in the process, D changes from increasing to decreasing (if D is initially increasing), or from decreasing to increasing (if D is initially decreasing). At the point where this happens, what is the value of the ratio M/D ? Please explain. (We might call this value the “threshold value” of M/D .)

To say that D changes from increase to decrease, or vice versa, is to say that $D' = 0$. Let's examine where this happens, by setting the above formula for D' equal to zero:

$$2M^2 - 0.8MD = 0.$$

Factor out an M :

$$M(2M - 0.8D) = 0.$$

Divide through by M (assuming $M \neq 0$):


$$2M - 0.8D = 0.$$

Solve for M/D :


$$M/D = 0.8/2 = 0.4.$$

So the threshold value of M/D is 0.4.


3. Which of the four graphs on the following page could possibly be a graph of the quantities M , D , and T modeled by the above rate equations? Please explain your reasoning carefully, and on the correct graph, label which curve is M , which is D , and which is T . Hint: start by thinking about increase and decrease.




(i)



(ii)



(iii)



(iv)

Since $T' = 0.8MD$ is always positive (as long as neither M nor D equals zero), at least one of our three curves must be steadily increasing. This eliminates graph (i). We can also eliminate graph (iv) because, by exercise 2 above, D' must also increase initially, and graph (iv) does not include two curves that are initially increasing.

To distinguish between graphs (ii) and (iii) we note that, by exercise 2 above, D peaks when M/D equals 0.4. This clearly eliminates graph (iii) – in that graph, at the point (around $t = 2.5$) where D peaks, we see that M is less than 0.1 and D is larger than 0.5, so that M/D is less than $0.1/0.5 = 1/5 = 0.2$.

The remaining graph (ii) must therefore be the correct graph. The labeling of each of the quantities M , D , and T in that graph follows by considering the signs of M' , D' , and T' .

4. Fill in the blanks (try to answer based primarily on quantitative reasoning and mathematics; you shouldn't need any advanced knowledge of chemical reactions):

A monomer may react with another monomer to form a dimer. These monomer-to-monomer reactions cause a decrease in the total quantity of monomers. Moreover, the rate at which this occurs is proportional to M^2 (since each of the M milligrams of monomers present has roughly M milligrams of other monomers with which to react). The monomer-to-monomer reactions therefore correspond to the term $-4M^2$ in the above equation for M' .

Further, whenever two monomers are lost to a monomer-to-monomer reaction, one dimer is gained. That is: the rate at which dimers are gained from such reactions equals half the rate at which monomers are lost to these reactions. Since half of $4M^2$ equals $2M^2$, the monomer-to-monomer reactions account for the term $2M^2$ in the above equation for D' .

A monomer may also react with a dimer to form a trimer. The rate at which this occurs is proportional to the product of the quantity of monomers and the quantity of dimers (since each of the M milligrams of monomers present has D milligrams of dimers with which to react). The decrease in M resulting from these monomer-to-dimer reactions therefore corresponds to the term $-0.8MD$ in the above equation for M' . Analogously, the decrease in D resulting from these monomer-to-dimer reactions corresponds to the term $-0.8MD$ in the above equation for D' .

Finally, when a monomer and a dimer are lost to a monomer-to-dimer reaction, one trimer is gained. This accounts for the term $0.8MD$ in the above equation for T' .

5. Use the rate equations on the first page, above, to compute $M' + 2D' + 3T'$. What does this tell you about $M + 2D + 3T$? How would you interpret this result in terms of the chemical reactions taking place?

We readily compute that

$$\begin{aligned} M' + 2D' + 3T' &= -4M^2 - 0.8MD + 2(2M^2 - 0.8MD) + 3(0.8MD) \\ &= (-4 + 2 \cdot 2)M^2 + (-0.8 + 2(-0.8) + 3(0.8))MD \\ &= 0. \end{aligned}$$

The fact that $M' + 2D' + 3T' = 0$ tells us that $M + 2D + 3T$ is constant.

Interpretation: if a monomer is considered a basic unit, a dimer counts as two such units, and a trimer counts as three, then the number of basic units is preserved throughout the reaction.