

More on sets.

Let A, B be sets.

Definitions 1, 2, 3.

We define:

1) The union $A \cup B$ ("A union B") of A and B by

$$A \cup B = \{x : x \in A \text{ or } x \in B\}$$

(in math, "or" is inclusive: it means one or the other, or both).

2) The intersection $A \cap B$ ("A intersect B") of A and B by

$$\begin{aligned} A \cap B &= \{x : x \in A \text{ and } x \in B\} \\ &= \{x \in A : x \in B\} = \{x \in B : x \in A\}. \end{aligned}$$

3) The difference $A - B$ ("A minus B") of A and B by

$$A - B = \{x \in A : x \notin B\}.$$

4) The Cartesian product $A \times B$ ("A cross B") of A and B by

$$A \times B = \{\text{ordered pairs } (x, y) : x \in A, y \in B\}.$$

Example A.

Let

$$\begin{aligned} A &= (-30, 84], \quad B = [12, 157), \quad C = \{e, f, g\}, \\ D &= \{e, m\}, \quad E = \{\{e\}, \{e, f, g\}, e, f, g\} \end{aligned}$$

Then:

$$A \cup B = (-30, 157), \quad A \cap B = [12, 84),$$

$$A - B = (-30, 12), \quad B - A = [84, 157),$$

$$A \times B = \{(x, y) : -30 < x \leq 84, 12 \leq y < 157\}$$

(a rectangle with part of its border missing).

$$C \cup D = \{e, f, g, m\}, \quad C \cap D = \{e\},$$

$$C - D = \{f, g\}, \quad D - C = \{m\},$$

$$C \times D = \{(e, e), (e, m), (f, e), (f, m), (g, e), (g, m)\},$$

$$C \cap E = \{e, f, g\}, \quad C - E = \emptyset, \quad E - C = \{\{e\}, \{e, f, g\}\},$$

etc.

Note: we can generalize Defns 1, 2, 4 to more than two sets, e.g.

$$C \cup D \cup E = \{e, f, g, m, \{e\}, \{e, f, g\}\}$$

$$\mathbb{R} \times \mathbb{R} \times \mathbb{R} \times \mathbb{R} = \{\text{ordered quadruples } (x, y, z, t) : x, y, z, t \in \mathbb{R}\}$$

(also written \mathbb{R}^4), etc.

We can generalize Defn 3 too, but be careful: for sets A, B, C , $A - (B - C)$ need not equal $(A - B) - C$. (E.g. consider $A = \{1, 2, 3, 4, 5, 6\}$, $B = \{1, 2, 3, 4\}$, $C = \{3, 4, 5, 6\}$.)

We can also mix operations, e.g.

$$(C \cap D) \cup E = E, \quad C \cap (D \cup E) = C,$$

etc.

Definition 5.

Given a set S , we define the power set $\mathcal{P}(S)$ to be the set of all subsets of S .

Example B.

For C and D as above,

$$\mathcal{P}(D) = \{\emptyset, \{e\}, \{m\}, \{e, m\}\},$$

$$\mathcal{P}(C) = \{\emptyset, \{e\}, \{f\}, \{g\}, \{e, f\}, \{f, g\}, \{e, g\}, \{e, f, g\}\}.$$

Note that $|\mathcal{P}(D)| = 4$, $|\mathcal{P}(C)| = 8$.

In general, we have

Theorem. If S is a finite set (that is, $|S| = n$ for some $n \in \mathbb{N}$), then

$$|\mathcal{P}(S)| = 2^{|S|}.$$

(Proof later.)

For example, for E as above,

$$|\mathcal{P}(E)| = 2^{|E|} = 2^5 = 32.$$

Note: if S is infinite, then $\mathcal{P}(S)$ is huge - even larger than S , in a certain sense.

E.g. $|\mathcal{P}(\mathbb{Z})| = |\mathbb{R}|$, in a sense we'll discuss later.