

PART I. Basic probability.**1. Permutations.**

- (a) The number of ways of arranging n objects in order is

$$n! = n \cdot (n-1) \cdot (n-2) \cdots 2 \cdot 1.$$

- (b) The number of ways of arranging r objects (in order) out of n objects is

$$n \cdot (n-1) \cdot (n-2) \cdots (n-r+1) = \frac{n!}{(n-r)!}.$$

- (c) The number of ways of arranging n objects, where n_1 of them are the same, n_2 of them are the same, \dots n_r of them are the same, is

$$\binom{n}{n_1, n_2, \dots, n_r} = \frac{n!}{n_1! n_2! \cdots n_r!}.$$

2. Combinations.

- (a) (Combinations.) The number of ways of choosing r objects out of n objects, without keeping track of order, is

$$\frac{n \cdot (n-1) \cdot (n-2) \cdots (n-r+1)}{r!} = \frac{n!}{r!(n-r)!}.$$

This number is sometimes called “ n choose r ,” written $\binom{n}{r}$. Note that this is also the number of r -element subsets of a set with n elements.

- (b) The number of ways of placing n objects into r distinct groups, of size n_1, n_2, \dots, n_r , is

$$\binom{n}{n_1, n_2, \dots, n_r} = \frac{n!}{n_1! n_2! \cdots n_r!}$$

(same number as in 1(c) above).

3. Probability axioms.

- (a) $P(A) \geq 0$ for any event A .
 (b) $P(S) = 1$, where S is the sample space.
 (c) If the events $A_1, A_2, A_3, A_4, \dots$ are mutually exclusive (no two of them can happen together), then

$$P(A_1 \cup A_2 \cup A_3 \cup A_4 \cup \cdots) = P(A_1) + P(A_2) + P(A_3) + P(A_4) + \cdots.$$

(The list $A_1, A_2, A_3, A_4, \dots$ could be finite or infinite.)

4. Basic probability rules and formulas.

- (a) If all outcomes in a sample space S are equally likely, and $|E|$ denotes the number of outcomes in the event E , then

$$P(E) = \frac{|E|}{|S|}$$

(assuming the sample space is a finite set).

- (b) For any event E ,

$$P(E) = 1 - P(E^c),$$

where E^c denotes the complement of E (meaning all outcomes in the sample space except those in E).

- (c) For any events E and F (not necessarily mutually exclusive), we have

$$P(E \cup F) = P(E) + P(F) - P(EF).$$

- (d) For any events E, F , and G (not necessarily mutually exclusive), we have

$$P(E \cup F \cup G) = P(E) + P(F) + P(G) - P(EF) - P(EG) - P(FG) + P(EFG).$$

PART II. Conditional probability.

1. Formulas for $P(E|F)$.

- (a) Given any events E and F , we have

$$P(E|F) = \frac{P(EF)}{P(F)}.$$

- (b) Suppose all events in the sample space are equally likely. Then for any events E and F , we have

$$P(E|F) = \frac{|EF|}{|F|}.$$

2. Formulas for $P(EF)$.

- (a) Given any events E and F , we have

$$P(EF) = P(F) \cdot P(E|F).$$

- (b) (Generalization.) Given any events A_1, A_2 , and A_3 , we have

$$P(A_1 A_2 A_3) = P(A_1) \cdot P(A_2|A_1) \cdot P(A_3|A_1 A_2).$$

- (c) (Further generalization.) Given any finite or infinite list of events, we have

$$P(\text{all events happen}) = P(\text{first one happens}) \cdot P(\text{second happens given that first does}) \\ \cdot P(\text{third does given that first two do}) \cdot P(\text{fourth does given that first three do}) \cdots$$

3. Independent events.

- (a) If events E and F are independent ($P(E) = P(E|F)$), we have

$$P(EF) = P(E) \cdot P(F).$$

- (b) (Generalization.) If events $A_1, A_2, A_3, A_4, \dots$ are independent (they don't affect each other), then

$$P(A_1 A_2 A_3 A_4 \cdots) = P(A_1) \cdot P(A_2) \cdot P(A_3) \cdot P(A_4) \cdots$$

(The list $A_1, A_2, A_3, A_4, \dots$ could be finite or infinite.)

4. Bayes's Formula (also known as the law of total probability).

- (a) For any events E and F ,

$$P(E) = P(F)P(E|F) + P(F^c)P(E|F^c)$$

(again, F^c denotes the complement of F).

- (b) (Generalization.) Suppose $F_1, F_2, F_3, \dots, F_n$ are mutually exclusive and exhaustive events. Then

$$P(E) = P(F_1)P(E|F_1) + P(F_2)P(E|F_2) + P(F_3)P(E|F_3) + \cdots + P(F_n)P(E|F_n).$$